

## Week 1 Examples

**Three Examples.** Solve differential equations without a book. Three basic examples used throughout a course in differential equations, which require only a calculus background.

**Growth-Decay:**  $\frac{dA(t)}{dt} = k A(t), A(0) = A_0.$

The unique solution is  $A(t) = A_0 e^{kt}$ . Radioactive decay. Jeweler's bench light experiment. Malthusian population dynamics. RC and LR circuits. Drug elimination. First-order chemical reactions, law of mass-action. Compound continuous bank interest.

**Newton Cooling:**  $\frac{du(t)}{dt} = -h(u(t) - u_1), u(0) = u_0.$

The solution is  $u(t) = u_1 + (u_0 - u_1) e^{-ht}$ . Hot chocolate at initial temperature  $u_0$  with room thermometer reading  $u_1$ . Symbol  $u(t)$  = time-varying hot chocolate dial thermometer temperature.

**Verhulst Dynamics:**  $\frac{dP(t)}{dt} = (a - bP(t))P(t), P(0) = P_0.$

The solution is  $P(t) = \frac{aP_0}{bP_0 + (a - bP_0)e^{-at}}$ . Fish population  $P(t)$  in **Cecret Lake at Alta**. Carrying capacity. Stocking and re-stocking. Harvesting.

**Example 1:** Exercise 1.2-2: Solve  $dy/dx = (x - 2)^2, y(2) = 1.$

Method of quadrature. Answer Check details. Non-reversible steps, false proof for  $0 = 1.$

<http://www.math.utah.edu/~gustafso/s2015/2250/FTC-Method-of-Quadrature.pdf>

**Example 2:** Exercise 1.2-4:  $dy/dx = 1/x^2, y(1) = 5.$

Power rule in Newton calculus. Answer check shortcuts.

**Example 3:** Exercise 1.2-10:  $dy/dx = x e^{-x}, y(0) = 1.$

Integral tables and integration by parts. Jennifer Lahti's solution:

<http://www.math.utah.edu/~gustafso/s2015/2250/2250Week1exercises-JenniferLahti-1.2-5+8+10.pdf>

**Example 4:** Exercise 1.3-8:  $dy/dx = x^2 - y$

Thread edge-to-edge solutions through the direction field at each blue dot. JPEG image source:

<http://www.math.utah.edu/~gustafso/s2015/2250/exercise1.3-8-EdwardsPenney.jpg>

**Example 5:** Exercise 1.3-14:  $dy/dx = y^{1/3}, y(0) = 0$

Explain application of the Peano and Picard theorems.

Computer numerical methods fail on this example. Why?

**Example 6:** Exercise 1.4-6: Solve  $y' = 3\sqrt{xy}$

Three answers. Book reports only one answer.

**Example 7:** Exercise 1.4-10: Solve  $(1 + x^2)y' = (1 + y)^2$

Two answers. Book reports only one answer.

**Example 8:** Exercise 1.4-18: Solve  $x^2y' = 1 - x^2 + y^2 - x^2y^2$  [See Example 11 *infra*]

**Example 9:** Exercise 1.4-22: Solve  $y' = 4x^3y - y, y(1) = -3$

**Example 10:** Show that  $y' = x + y$  is not separable.

TEST I.  $f_x/f$  depends on  $y$  implies  $y' = f(x, y)$  not separable.

**Example 11:** Find a factorization  $f(x, y) = F(x)G(y)$  given

(1)  $f(x, y) = 2xy + 4y + 3x + 6$

(2)  $f(x, y) = (1 - x^2 + y^2 - x^2y^2)/x^2$

Answers: (1)  $F = x + 2, G = 2y + 3$ ; (2)  $F = (1 - x^2)/x^2, G = 1 + y^2$ . Main idea: Choose  $y = 0$  in  $F(x) = f(x, y)/G(y)$  to find  $F(x) = (3x + 6)/G(0)$  in equation (1). How to find  $G$ ? **Warning:** Divide by zero is not allowed. Choose  $y = 0, y = 1$ , etc, until no divide by zero error.

**Example 12:** Midterm 1 examples:  $y' = x + y, y' = x + y^2, y' = x^2 + y^2$