

1220-90 Final Exam  
Fall 2013

Name \_\_\_\_\_

**Instructions.** Show all work and include appropriate explanations when necessary. Answers unaccompanied by work may not receive credit. Please try to do all work in the space provided and circle your final answers. The last page contains some useful formulas.

1. (10pts) Compute the following derivatives:

(a) (5pts)  $D_x(\ln(x^3 + x))$

(b) (5pts)  $D_x(x^{\sin x})$

2. (10pts) Use L'Hôpital's Rule to compute the following limits:

(a) (5pts)  $\lim_{x \rightarrow 0} \frac{e^{2x} - 1}{x}$

(b) (5pts)  $\lim_{x \rightarrow 0} \frac{x^2}{\sin^2 x}$

3. (5pts) Find a formula for the inverse of the function

$$f(x) = \sqrt[3]{6 - x}$$

4. (24pts) Evaluate the following indefinite integrals. Remember  $+C$ !

(a) (6pts)  $\int x e^{-3x} dx$

(b) (6pts)  $\int \sin^4 x \cos^3 x dx$

(c) (6pts)  $\int \frac{1}{x^2 - 3x + 2} dx$      **Hint:**  $x^2 - 3x + 2 = (x - 2)(x - 1)$ .

(d) (6pts)  $\int \frac{1}{x^2 + 4x + 5} dx$      **Hint:**  $x^2 + 4x + 5 = (x + 2)^2 + 1$ .

5. (12pts) Determine, by whatever method you wish, whether the following series are convergent or divergent. Circle 'C' if the series is convergent or 'D' if the series is divergent. No work is necessary.

C    D     $\sum_{n=1}^{\infty} \frac{1}{n^3}$

C    D     $\sum_{n=1}^{\infty} \frac{n^2 + 1}{5n^2 + 9}$

C    D     $\sum_{n=1}^{\infty} 3 \left(\frac{1}{8}\right)^n$

C    D     $\sum_{n=1}^{\infty} (-1)^{n-1} e^{-n}$

C    D     $\sum_{n=1}^{\infty} \sin(n\pi)$

C    D     $\sum_{n=1}^{\infty} \frac{n!}{n^2}$

6. (12pts) Perform the integral test in two steps:

- (a) (6pts) Use **integration by parts** to find

$$\int \frac{\ln x}{x^2} dx.$$

**Hint:** Set  $u = \ln x$  and  $dv = x^{-2} dx$ .

- (b) (6pts) Use the above and the **integral test** to determine whether the following series converges or diverges

$$\sum_{n=1}^{\infty} \frac{\ln n}{n^2}.$$

7. (10pts) Find the **interval** of convergence of the power series

$$\sum_{n=1}^{\infty} \frac{(x-1)^n}{n^2}$$

8. (9pts) Let

$$f(x) = \ln(1+x)$$

(a) (6pts) Find the following derivatives of  $f(x)$  evaluated at  $x=0$ :

$$f(0) = \underline{\hspace{2cm}}$$

$$f'(0) = \underline{\hspace{2cm}}$$

$$f''(0) = \underline{\hspace{2cm}}$$

(b) (3pts) Use your computations above to write out  $P_2(x)$ , the second degree MacLaurin polynomial for  $f(x)$ .

9. (10pts) Match the function with its Maclaurin series by writing the letter in the blank provided.

\_\_\_\_\_  $f(x) = e^x$

A.  $\sum_{n=0}^{\infty} x^n$

\_\_\_\_\_  $f(x) = \frac{1}{1-x}$

B.  $\sum_{n=1}^{\infty} (-1)^n 2^{n+1} x^n$

\_\_\_\_\_  $f(x) = x \sin x$

C.  $\sum_{n=0}^{\infty} \frac{x^n}{n!}$

\_\_\_\_\_  $f(x) = \cos x$

D.  $\sum_{n=0}^{\infty} \frac{(-1)^n x^{2n+2}}{(2n+1)!}$

\_\_\_\_\_  $f(x) = \frac{2}{1+2x}$

E.  $\sum_{n=0}^{\infty} \frac{(-1)^n x^{2n}}{(2n)!}$

10. (8pts) Consider the conic section

$$4x^2 + y^2 + 8x - 6y + 4 = 0.$$

(a) (2pts) What type of conic section is this? Circle one answer below:

Parabola

Ellipse

Hyperbola

(b) (4pts) The center of the conic section is the point ( \_\_\_\_\_ , \_\_\_\_\_ ).

(c) (2pts) The shortest distance from the center to a point on the conic section is \_\_\_\_\_.

11. (16pts) Match the polar equation to its graph by writing the letter in the blank provided. Every answer will be used exactly once.

\_\_\_\_\_  $r = \cos 3\theta$

\_\_\_\_\_  $r = \sin \theta$

\_\_\_\_\_  $r = 1 + 2 \cos \theta$

\_\_\_\_\_  $r = \sin 2\theta$


\_\_\_\_\_  $r = 2$

\_\_\_\_\_  $r = \theta$


\_\_\_\_\_  $r = 2 + 1 \cos \theta$

\_\_\_\_\_  $r = \cos \theta$


**A**




**B**




**C**




**D**




**E**




**F**



**G**



**H**



12. (6pts) Match the Cartesian coordinates  $(x, y)$  on the left to the corresponding polar coordinates  $(r, \theta)$  on the right by writing the letter in the blank provided. Every answer will be used exactly once.

\_\_\_\_\_  $(1, 1)$

A.  $(1, \frac{3\pi}{2})$

\_\_\_\_\_  $(-\sqrt{3}, 1)$

B.  $(\sqrt{2}, \frac{\pi}{4})$

\_\_\_\_\_  $(0, -1)$

C.  $(2, \frac{5\pi}{6})$

13. (6pts) Match the polar coordinates  $(r, \theta)$  on the left to the corresponding Cartesian coordinates  $(x, y)$  on the right by writing the letter in the blank provided. Every answer will be used exactly once.

\_\_\_\_\_  $(4, \pi)$

A.  $(-4, 0)$

\_\_\_\_\_  $(2, \frac{4\pi}{3})$

B.  $(-1, -\sqrt{3})$


\_\_\_\_\_  $(\sqrt{2}, \frac{7\pi}{4})$

C.  $(1, -1)$

14. (12pts) Consider the curve determined by the polar equation  $r = e^{\theta/3}$ . A graph of this polar curve, between  $\theta = 0$  and  $\theta = \pi$  is given at the bottom of the page.

- (a) (6pts) Find the area of the region inside the curve and above the  $x$ -axis.

- (b) (6pts) Find the length of the curve.



**Trigonometric Formulas:**

$$\sin^2 x + \cos^2 x = 1$$

$$1 + \tan^2 x = \sec^2 x$$

$$\sin^2 x = \frac{1}{2}(1 - \cos 2x)$$

$$\cos^2 x = \frac{1}{2}(1 + \cos 2x)$$

$$\sin 2x = 2 \sin x \cos x$$

$$\sin x \cos y = \frac{1}{2}(\sin(x - y) + \sin(x + y))$$

$$\sin x \sin y = \frac{1}{2}(\cos(x - y) - \cos(x + y))$$

$$\cos x \cos y = \frac{1}{2}(\cos(x - y) + \cos(x + y))$$

**Inverse Trigonometric Formulas:**

$$D_x \sin^{-1} x = \frac{1}{\sqrt{1-x^2}} \quad -1 < x < 1$$

$$D_x \cos^{-1} x = -\frac{1}{\sqrt{1-x^2}} \quad -1 < x < 1$$

$$D_x \tan^{-1} x = \frac{1}{1+x^2}$$

$$D_x \sec^{-1} x = \frac{1}{|x|\sqrt{x^2-1}} \quad |x| > 1$$

**Calculus with Polar Curve  $r = f(\theta)$ :**

$$A = \int_a^b \frac{1}{2} [f(\theta)]^2 d\theta$$

$$L = \int_a^b \sqrt{[f(\theta)]^2 + [f'(\theta)]^2} d\theta$$

$$m = \frac{f'(\theta) \sin \theta + f(\theta) \cos \theta}{f'(\theta) \cos \theta - f(\theta) \sin \theta}$$