

Square-Triangular Numbers

Jim Carlson

April 26, 2004

Contents

1	Introduction	2
2	Existence	2
3	Finding an equation	3
4	Solutions by brute force	4
5	Speeding things up	5
6	Solutions by algebraic numbers	7
7	Recursion relations	9
8	Fundamental solutions, continued fractions	10
9	More about continued fractions	13
10	More Diophantine equations	14

1 Introduction

We begin with a puzzle. Its solution will lead us deep into the land of number theory:

Problem 1 *Is it possible to arrange a collection of stones in both a square and a triangular shape?*

To make sure we understand the problem, let's imagine that we have four stones. They can certainly be arranged in a square:

```
* *  
* *
```

But can we arrange them as a triangle? It turns out we can't, as the next figure shows:

```
*           *           *           etc.  
           * *         * *  
           * * *       * * *  
                   * * *
```

But perhaps there is some special number of stones — let's call it N — that can be arranged in both a square and a triangular shape:

Problem 2 *Is there a number N that is both square and triangular?*

Comments: *Make two lists, compare them. Try brute-force search. Do square-triangular numbers exist? How frequent are square-triangular numbers? Are there infinitely many of them?*

2 Existence

One way to answer the previous question — *do square-triangular numbers exist?* — is to make two lists. One list is for the square numbers 1, 4, 9, 16, etc., and another is for the triangular numbers 1, 3, 6, 10, etc. A number that appears in both lists solves the puzzle. You probably found 36 in both lists. This is the first square-triangular number. We write it as

$$N_1 = 36.$$

This raises the question: *are there any other square-triangular numbers?* So let's set a small challenge:

Problem 3 *Is there a square-triangular number $M > 100$?*

You may be able to find more square-triangular numbers by comparing lists, but we should pause to ask: *Are there a better methods?* Better methods will be needed to solve more difficult questions. Indeed, let us set the following challenge:

Problem 4 *(a) Is there a square triangular number $M > 10^6$? (b) What about $M > 10^{12}$? (c) And $M > 10^{24}$? (d) How many square-triangular numbers are there?*

One way to find square-triangular numbers might be to find a formula for them. Another might be to find an equation whose solutions are exactly the square-triangular numbers. Still another way is to write a computer program to search for square-triangular numbers. We will explore each these ideas below, and will eventually answer the fundamental question (d) *how many square-triangular numbers are there?*

3 Finding an equation

We now ask if there is a more algebraic way of approaching the problem of square-triangular numbers:

Problem 5 *(a) Is there an equation whose solution gives all square triangular numbers. (b) Is there a formula for these numbers?*

Let's see if we can solve part (a). A number N is square if it can be written as $N = m^2$ for some m . A number is triangular if it can be written as $n(n+1)/2$ for some n . This statement requires a proof:

Problem 6 *Show that every triangular number can be written as $n(n+1)/2$ for some n .*

If N is both square and triangular, then it can be written as m^2 for some m and as $n(n+1)/2$ for some n . Thus $N = m^2 = n(n+1)/2$ is given by solutions to the equation

$$m^2 = n(n+1)/2 \tag{1}$$

Problem 7 *Find several positive integer solutions to the equation $m^2 = n(n+1)/2$. What are the corresponding square-triangular numbers?*

An equation like $m^2 = n(n+1)/2$ for which we demand *integer* solutions is called a *Diophantine equation*. Diophantus was a Greek mathematician who

worked in Alexandria on questions like this around 250 AD. To solve our equation we will put it in simpler form. Begin by clearing denominators:

$$2m^2 = n^2 + n.$$

Now it is an equation with integer coefficients, the usual form of a Diophantine equation. Next, complete the square:

$$2m^2 = (n + 1/2)^2 - 1/4$$

Clearing denominators again, we find

$$8m^2 = (2n + 1)^2 - 1,$$

or

$$(2n + 1)^2 - 2(2m)^2 = 1.$$

Introduce new variables

$$x = 2n + 1 \text{ and } y = 2m. \tag{2}$$

Then our equation becomes

$$x^2 - 2y^2 = 1 \tag{3}$$

This is Pell's equation, named for John Pell (1611-1685). In fact Mr. Pell had almost nothing to do with the equation that bears his name. Archimedes' Cattle Problem can be reduced to the solution of Pell's equation, so perhaps the problem was known as early as 250 BC. However, there is no direct evidence for this speculation. In any case, by 628 AD, the Indian mathematician Brahmagupta not only explicitly mentions the problem but shows amazing progress in understanding it. See [1]. As mentioned in the introduction, this Pell's equation is the equation of a hyperbola. A solution is a *lattice point* on this hyperbola. A lattice point is a point with integer coefficients, like (5, 7).

Theorem 1 *If x, y is a solution of Pell's equation, then $N = (y/2)^2$ is a number that is both square and triangular. The square has $y/2$ dots on a side, the triangle has $(x - 1)/2$ dots on the bottom row.*

Problem 8 *How many solutions of Pell's equation $x^2 - 2y^2 = 1$ can you find (in five minutes)? How many solutions do you think there are?*

Problem 9 *Draw a picture of the lattice points in the box $|x| < 5, |y| < 5$ and of the hyperbola $x^2 - 2y^2 = 1$. Indicate the solutions of Pell's equation.*

4 Solutions by brute force

Let us now find square-triangular numbers by finding solutions to Pell's equation. One strategy is to inspect all pairs of integers x, y , with $0 \leq x \leq N$,

$0 \leq y \leq N$. If a pair solves the equation, we write it down on our list of solutions. If it does not, we go on to the next pair. This is called the *brute-force method*.

Problem 10 How many pairs of “candidate solutions” x, y for Pell’s equation are there satisfying $0 \leq x \leq N, 0 \leq y \leq N$?

Problem 11 Following the notation of the previous problem, find all nonnegative solutions of Pell’s equation with $N < 100$. What are the corresponding square-triangular numbers?

Problem 12 Discuss the feasibility of using brute-force search when $N = 1000$. Is there a way of improving our search?

Let’s talk about strategy. In the previous problem we could list all the points x, y with integer coordinates that lie in the box $0 \leq x, y < N$. There are N^2 such points. So when $N = 100$, there are 10,000 points to inspect. Already a large number! Consider how much larger it is for $N = 1000$. The number of points to inspect grows quadratically in the magnitude of the input N . Can we do better? That is, can we speed things up?

Problem 13 Suppose that a computer program can inspect all candidate solutions with $0 \leq x, y < 10^3$ in one second. How long do you expect it to take if $x, y < 10^6$? What if we try an upper bound of 10^9 or 10^{12} ?

Computer scientists usually measure (or estimate) the time taken for a computation to produce its output in terms of the number of bits in the input. If the input is an integer, like 5, then the number of bits in the input is the number of binary digits needed to express the number. Since the binary representation of 5 is 101, the number of bits of input for an input “5” is three.

Problem 14 Express 17 in binary form. What is the decimal form of the binary number 11011011?

Problem 15 Suppose that inspection of a lattice point takes T seconds. If the “inspection box” is defined by $0 < x < N, 0 < y < N$. How many seconds S will it take to inspect all lattice points in the box? Your answer should express T as a function of N . Now let b be the number of bits needed to write the number N . Express S as a function of b .

5 Speeding things up

Two ways of speeding up the search for solutions to Pell’s equation are: (a) use a computer, (b) be more clever. Of course, if we combine (a) and (b), we will do even better.

First let's discuss using a computer. Using the Python language (see xxx), we will design a function `search(N)` that prints out all solutions to Pell's equation satisfying $0 \leq x < N$, $0 \leq y < N$. Thus we can do the following:

```
>>> search(100)
1 0
3 2
17 12
99 70
```

The function `search` finds four solutions in the twinkling of an eye. Here is how the function is defined:

```
def search(N):
    for x in range(0,N):
        for y in range(0,N):
            if x*x - 2*y*y == 1:
                print x, y
```

The code is almost self explanatory. Numbers x and y in the range $[0, N)$ are generated and for each pair (x, y) we test whether the equation holds. If it does, we print out the solution. For more information on Python, see xxx.

The function `search(N)` works quite well when $N = 1000$, but already when $N = 10000$ it requires considerable time. We could use a faster computer or a faster language, but the real difficulty is that the amount of work to be done grows quadratically in N .

Problem 16 *Use the Python program `search` with $N = 100, 1000, 100000$. Comment on what you find. Can you find a solution of Pell's equation with $y > 10^6$ using `search`?*

Now let's try to be more clever. To begin, rewrite Pell's equation as

$$x^2 = 1 + 2y^2.$$

Notice that we can systematically inspect values of y , accepting only those for which $1 + 2y^2$ is a perfect square. A table of squares helps to do this quickly, even if we are doing the inspection manually.

Problem 17 *Find all solutions of Pell's equation with $y < 100$ using "linear search on y ."*

Problem 18 *Now that you have tried linear search by hand to find solutions of Pell's equation, design a computer program which uses the same method. Do you find more solutions? How big of a solution can you find?*

For the last problem it is useful to have a function that determines whether a number is a square or not. The following function works well if the numbers are not too big. For larger numbers one should use an integer version of Newton's method (xxx).

```
def issquare(n):
    isqrt = int(sqrt(n))
    if n == isqrt*isqrt:
        return 1
    else:
        return 0
```

The function `issquare` works by computing the integer part of \sqrt{n} . Thus, if $n = 5$, $isqrt = int(sqrt(5)) = int(2.23606\dots) = 2$. Then we check if the square of this “integer square root” is equal to n . If it is, we return 1 (true). Otherwise we return 0 (false). Since floating point arithmetic is inaccurate for very large numbers, this function will fail when n is too large. One really needs a true integer square root function. See xxx.

6 Solutions by algebraic numbers

In the last section you found that 1, 0 and 3, 2 are solutions to Pell's equation. These give the “trivial” square-triangular numbers 0 and 1. You also have found the solution 17, 12 which gives the square triangular number 36. Finally, you found larger solutions using a combination of brute force search and a somewhat more clever search, possibly aided by a computer. But there is a much better way of finding solutions which is based on a little theory that comes from a factored form of Pell's equation:

$$(x - \sqrt{2}y)(x + \sqrt{2}y) = 1. \tag{4}$$

The key idea is that every solution of Pell's equation gives a factorization of 1. Thus $x, y = 3, 2$ gives the factorization

$$(3 - 2\sqrt{2})(3 + 2\sqrt{2}) = 1.$$

We claim that from an “old” factorization one can produce a “new” one. To see this, square the preceding equation:

$$((3 - 2\sqrt{2})(3 + 2\sqrt{2}))^2 = 1.$$

It can be rewritten as

$$(3 - 2\sqrt{2})^2(3 + 2\sqrt{2})^2 = 1.$$

Doing the algebra, we find

$$(17 - 12\sqrt{2})(17 + 12\sqrt{2}) = 1$$

Thus $x, y = 17, 12$ should be a solution of Pell's equation — and it is, as you can easily check. It corresponds to the square-triangular number 36.

Problem 19 *What is the solution of Pell's equation corresponding to $(3 + 2\sqrt{2})^3$? What about $(3 + 2\sqrt{2})^4$? Check your results. What are the corresponding square-triangular numbers?*

Problem 20 *What is the solution of Pell's equation corresponding to $(3 + 2\sqrt{2})^{16}$? Check your result. What is the corresponding square-triangular number?*

Problem 21 *Are there finitely or infinitely many square-triangular numbers? Explain.*

Problem 22 *How many integer solutions are there to the equation $x^2 - y^2 = 1$?*

Problem 23 *Investigate integer solutions of quadratic equations of the form $x^2 - Ny^2 = 1$ for various values of N . What conclusions can you draw?*

Notes

It is worth reflecting on what we have done. Our problem is to find integer solutions to an equation with integer coefficients. But the best way so far to do this is to use irrational numbers. We solve a problem in one area of mathematics by using ideas from another area.

The numbers used, by the way, are not random irrational numbers. First, they have the form $a + b\sqrt{2}$ where a and b are integers. Secondly these numbers form what is called a *ring*: the sum any two such numbers is again such a number, as is the product. What about the quotient of two such numbers?

We denote the ring of numbers $\{ a + b\sqrt{2} \mid a, b \in \mathbf{Z} \}$ by the symbol $\mathbf{Z}[\sqrt{2}]$ — “the integers \mathbf{Z} with $\sqrt{2}$ adjoined.” All numbers in $\mathbf{Z}[\sqrt{2}]$ have a remarkable property: each satisfies a quadratic equation with integer coefficients.

Problem 24 *Show that $\mathbf{Z}[\sqrt{2}]$ is a ring.*

Problem 25 *Describe the set of numbers obtained by taking quotients of elements of $\mathbf{Z}[\sqrt{2}]$. What properties does it have?*

Problem 26 *What is the quadratic equation with integer coefficients satisfied by $5 + 7\sqrt{2}$?*

Problem 27 Show that every number in $\mathbf{Z}[\sqrt{2}]$ satisfies a quadratic equation with integer coefficients.

There is one final observation worth making. Some of the numbers U in $\mathbf{Z}[\sqrt{2}]$ have a *multiplicative inverse*. This means that there is a number V , also in $\mathbf{Z}[\sqrt{2}]$, such that $UV = 1$. Can you determine these numbers? They are called the *units* of $\mathbf{Z}[\sqrt{2}]$. The units form a *group* under multiplication: elements can be multiplied, and also divided.

Problem 28 Describe the units of $\mathbf{Z}[\sqrt{2}]$.

7 Recursion relations

In the last section we used powers of the “magic” irrational numbers $\mu = 3 + 2\sqrt{2}$ to find integer solutions to Pell’s equation. We’ll now show how to use μ to compute solutions even more efficiently. The idea is that the n -th solution is given by

$$\mu^n = x_n + y_n\sqrt{2}.$$

Our goal is to compute x_{n+1} and y_{n+1} from x_n and y_n . To this end notice that

$$x_{n+1} + y_{n+1} = (3 + 2\sqrt{2})(x_n + y_n\sqrt{2}).$$

If we multiply out the right hand side (RHS) and collect terms, we get

$$RHS = (3x_n + 4y_n) + (2x_n + 3y_n)\sqrt{2}.$$

Comparing with the left-hand side, we find that

$$x_{n+1} = (3x_n + 4y_n) \tag{5}$$

$$y_{n+1} = (2x_n + 3y_n) \tag{6}$$

These so-called *recursion relations* allow us to quickly compute large solutions to Pell’s equation, as well as to solve many other problems.

Problem 29 Let $U_n = (x_n, y_n)$ be the n -th solution of Pell’s equation. Compute U_8 , as well as the associated square-triangular number M_8 .

Problem 30 Find a square-triangular number $M > 10^{20}$.

Problem 31 Let $U_n = (x_n, y_n)$ be the n -th solution of Pell’s equation. How much bigger than x_n is x_{n+1} ? What can you say about the y coordinates?

Problem 32 Let M_n be the n -th square-triangular number. What can you say about its size as a function of n ?

Problem 33 What is the (approximate) number of solutions to Pell's equation with $x < k$? What is the (approximate) number of square-triangular numbers with $M < k$?

Problem 34 Study the ratio $r_n = x_n/y_n$ as n becomes large. What can you conclude? Can you explain your conclusion?

8 Fundamental solutions, continued fractions

By following the pattern used to solve the equation $x^2 - 2y^2 = 1$ it is not hard to find infinitely many solutions to the equation $x^2 - Ny^2 = 1$ given a “nontrivial” solution (a, b) with $a > 0$ and $b > 0$. When N is small, we can do this by trial and error, if the nontrivial solution exists.

Problem 35 Find a nontrivial solution of $x^2 - Ny^2 = 1$ for the prime numbers $N = 2, 3, 5, 7, 11, 13, 17$. Any observations? For what nonprime coefficients N can we find nontrivial solutions?

For general N the trial and error (or systematic search) strategy may not be efficient enough to produce a solution in a reasonable amount of time. However, there is another method, based on the idea of *continued fractions*, that works quite well.

A continued fraction is an expression like

$$4 + \frac{1}{2 + \frac{1}{7 + \frac{1}{3}}}$$

Note that the numerators are all equal to 1. A more compact notation for the above continued fraction is $[4; 2, 7, 3]$.

Problem 36 What is the simple fraction corresponding to $[4; 2, 7, 3]$? What is the value of the continued fraction $[1; 2, 3, 4]$?

We can also convert any simple fraction into a continued fraction. For this we use three ideas — the integer part of a number, its remainder mod 1, and unit fractions:

1. Given a number α , let $[\alpha]$ be the “largest integer” in α . Thus $[3.1416] = 3$.
2. The “remainder mod 1” of a number, which we write as $\{\alpha\} = \alpha - [\alpha]$. Thus $\{3.1416\} = 0.1416$. The remainder of a number mod 1 is always less than 1 and greater than or equal to zero. Notice that

$$\alpha = [\alpha] + \{\alpha\}, \tag{7}$$

3. A *unit fraction* is a number of the form $1/x$.

Now let's find the continued fraction of $1281/401$. The first step is to apply (7) to write the given number as an integer and a remainder mod 1:

$$1281/401 = 3 + 78/401.$$

Then express the last term as a unit fraction:

$$1281/401 = 3 + 1/(401/78)$$

Repeat this two-step process until the denominator of the unit fraction is an integer. Thus

$$1281/401 = 3 + 1/(5 + 11/78) = 3 + 1/(5 + 1/(78/11)).$$

and then

$$1281/401 = 3 + 1/(5 + 1/(7 + 1/11)).$$

The process stops at this point, since the denominator of the unit fraction is an integer. The result is the continued fraction $[3; 5, 7, 11]$.

Problem 37 Find the continued fraction of $1193/532$.

Problem 38 Find the continued fraction of $\sqrt{2} \sim 1.4142135623730951$. Comment on your results.

The same process can be used to compute the continued fraction of $\sqrt{2}$. This can be done using the decimal expansion and a bit of guessing and a leap of faith, or exact algebra. Let's try it with the decimal expansion first. Step one yields

$$1.4142135623730951 = 1 + 0.4142135623730951,$$

Since

$$1/0.4142135623730951^{-1} = 1/(2.41421356454765235)$$

step two yields

$$1.414213562 = 1 + 1/(2.41421356454765235)$$

Applying steps one and two to 2.41421356454765235 , we find

$$1.414213562 = 1 + 1/(2 + 1/(2.41421354969884616))$$

Notice that the denominators of the last two expressions are almost all the same — they agree through the seventh decimal place. If we used infinite precision arithmetic, there would be exact agreement of the remainders no matter how far out we carry the expansion. With exact agreement we would find the infinite continued fraction

$$\sqrt{2} = [1, 2, 2, 2, \dots].$$

Problem 39 Find the continued fractions of $\sqrt{3}$ and $\sqrt{7}$ using a decimal approximation to each.

There is, of course, a way of using full precision: just work with radicals. We illustrate the method for $\sqrt{2}$. First, write $\sqrt{2}$ as an integer part and a remainder mod 1:

$$\sqrt{2} = 1 + (\sqrt{2} - 1).$$

Next, write the remainder as a unit fraction:

$$\sqrt{2} = 1 + 1/(\sqrt{2} - 1)^{-1}.$$

Rationalize the denominator to get

$$\sqrt{2} = 1 + 1/(1 + \sqrt{2}).$$

We could proceed as before (this is recommended for practice), but at this point we notice that the formula is self-similar: we can substitute for the $\sqrt{2}$ on the right-hand side the entire right-hand side:

$$\sqrt{2} = 1 + 1/(2 + 1/(1 + \sqrt{2})) = 1 + 1/(2 + 1/(2 + 1/(1 + \sqrt{2}))) = \dots$$

Repeating this argument, we *prove* that

$$\sqrt{2} = [1, 2, 2, 2, \dots]$$

Problem 40 Find the continued fraction expansions of $\sqrt{3}$ and $\sqrt{7}$ without using decimal approximations — or approximations of any kind. Prove that your result is correct.

We can now begin to see what the relation of continued fractions is to Pell's equation $x^2 - Ny^2 = 1$. Given a continued fraction $[a; b, c, d, \dots]$ we can form its *convergents*. These are shortened continued fractions like $[a; b]$, $[a; b, c]$, etc.

Problem 41 Compute the first three convergents of the continued fraction expansion of $\sqrt{2}$. How do they relate to the solutions of Pell's equation? Do you detect a general pattern?

Problem 42 Investigate continued fractions as a tool for solving other equations of Pell's type, $x^2 - Ny^2 = 1$. Be sure to try $N = 3$, $N = 7$. Look for patterns.

A *fundamental* solution to Pell's equation is a solution (a, b) , with both a and b positive, from which all other positive solutions can be found using the expression $(a + b\sqrt{N})^k$. A fundamental solution is smaller than all other positive solutions.

The *size* of a fundamental solution is $\log(a)/\log(N)$. It is roughly the number of digits of a divided by the number of digits in N .

Problem 43 (Contest) Find the largest fundamental solution you can.

9 More about continued fractions

Let's look more closely at our method for finding continued fractions. Define a *continued fraction with remainder* to be an expression of the form

$$F = [a_0, a_1, \dots, a_n, \theta],$$

where $0 < \theta < 1$. We expand such a fraction by replacing θ by $[1/\theta], \{1/\theta\}$, where $[x]$ denotes the integer part of x and $\{x\}$ denotes its fractional part — the part between 0 and 1. Thus F becomes

$$F' = [a_0, a_1, \dots, a_n, [1/\theta], \{1/\theta\}],$$

Let's try this rule on $\sqrt{7} \sim 3.6055512754639891$. We have

```
[3, 0.6055512754639891] <== (a)
[3, 1, 0.65138781886599784]
[3, 1, 1, 0.53518375848799526]
[3, 1, 1, 1, 0.86851709182133385]
[3, 1, 1, 1, 1, 0.15138781886599184]
[3, 1, 1, 1, 1, 6, 0.60555127546422849] <== (b)
[3, 1, 1, 1, 1, 6, 1, 0.65138781886534503]
```

The remainders at (a) and at (b) are 0.6055512754639891 and 0.60555127546422849 disagree in the 12th place. If you use a calculator that works to a higher precision, you will find that the agreement is higher. If you work with exact expressions you will find that the agreement is perfect. This proves that the continued fraction is

$$[3, 1, 1, 1, 1, 6, 1, 1, 1, 1, 6, \dots]$$

with the pattern $[1, 1, 1, 1, 6]$ repeating infinitely often.

Problem 44 *Compute the continued fraction expansion of $\sqrt{11}$, $\sqrt{12}$, $\sqrt{13}$, $\sqrt{14}$. Do you notice a pattern? Can you prove that it is correct? What do you conclude about fundamental solutions to Pell's equation?*

The Python code below is useful in computing continued fractions:

```
def expand( C ):
    x = C[-1] # last element
    x = 1.0/x
    a = int(x)
    b = x - a
    return C[:-1]+[a,b]
```

To work with exact expressions, we use numbers of the form $a + b\sqrt{7}$ where a and b are rational numbers. The set of all such expressions is closed under

addition, subtraction, multiplication and division by nonzero numbers. It is a so-called *number field*, and is written $\mathbf{Q}(\sqrt{7})$. For some purposes (such as writing computer program), it is better to write elements of this field as $(a + b\sqrt{7})/c$, where a , b , and c are integers. Then the field operations are given by integer formulas involving triples of integers $[a, b, c]$ and $[a', b', c']$.

Problem 45 Find integer formulas for the four field operations in $\mathbf{Q}(\sqrt{D})$.

Problem 46 Find an algorithm for computing the continued fraction expansion of \sqrt{D} using only integer operations on lists $[a, b, c]$

Problem 47 Find the continued fraction expansion of $\sqrt{61}$. Find a fundamental solution to $x^2 - 61y^2 = 1$. What is the length of this solution?

Problem 48 Investigate the continued fraction expansion of $2^{1/3}$.

10 More Diophantine equations

Here is another Diophantine problem, one that comes from geometry:

Problem 49 Find all positive integer solutions of the equation (1) $x^2 + y^2 = z^2$. What about the related equation (2) $x^3 + y^3 = z^3$? What about equations with more variables, like (3) $x^3 + y^3 + z^3 = w^3$?

Solutions of the Diophantine equation (1) are called *Pythagorean triples*, and of course we know one: 3, 4, 5. These are the lengths of a right triangle with whole-number sides. Once we have one Pythagorean triple, it is easy to find others, such as 6, 8, 10. But these are not interesting, since they correspond to similar triangles. What we really seek are *primitive* Pythagorean triples. These are triples of numbers satisfying the equation which have no common factors.

A basic question about primitive Pythagorean triplets is: (1) how many of them are there? One can ask other questions. (2) Is there a formula for them? (3) Are there interesting patterns that we can find in the triples? By studying some “experimental data” — lists of triples — one can make a start on (3) and perhaps even say something about (1) and (2).

Fermat’s theorem

One of the most famous mathematical problems of all time is “Fermat’s last theorem.” Pierre Fermat (1601–1665) was a judge in Toulouse, France and also a very serious amateur mathematician. One evening reading a copy of Diophantus’ *Arithmetica*, newly rediscovered and translated from Greek to Latin,

he came on a theorem about Pythagorean triplets. In the margin of the book he wrote “I have found a marvelous proof that there are no positive integer solutions to the equation

$$x^n + y^n = z^n \tag{8}$$

for $n > 2$. Despite much progress in special cases, the problem remained unsolved until Andrew Wiles, a British mathematician working at Princeton University, announced his solution in 1994.

Wiles’ proof, not only settled an old problems, but it also opened the doors to new areas of research through the introduction of new ideas and techniques. It relies on a great deal of advanced mathematics that was developed since Fermat’s time, but there is one part that we can mention here. Given a (hypothetical) solution a, b, c to Fermat’s equation, one can consider the cubic equation

$$y^2 = (x - a)(x - b)(x - c) \tag{9}$$

Cubic Diophantine equations are not completely understood, but they are much better understood than those of higher degree. They played a huge role in Wiles’ proof. Indeed, the unsuspected connection between cubic equations and Fermat’s equation, discovered in 1982 by the German mathematician Gerhard Frey, and proved in 1986 by the American mathematician Kenneth Ribet, was one of the milestone events that led to the solution Fermat.

References

- [1] <http://www-groups.dcs.st-and.ac.uk/~history/HistTopics/Pell.html>