

Rigidity of Harmonic Maps of Maximum Rank

James A. Carlson and Domingo Toledo

June 24, 1992

Contents

1. Introduction.	1
2. Outline of the proof	5
3. Existence of adapted root systems	11
4. Root systems.	14
5. Combinatorial arguments	18
6. Centralizers of semi-simple elements.	29
7. Rigidity	34
8. Existence.	37
9. Concluding Remarks	40

§ 1. Introduction.

Fix an irreducible symmetric space $X = G/K$ of non-compact type which is not Hermitian symmetric, and where the group G is a classical group. The purpose of this paper is to study the “largest” possible mappings of compact Kähler manifolds to quotients of X by discrete groups of isometries.

To make the meaning of “largest” precise, let us define the *homotopical rank* of a continuous map between compact manifolds to be the smallest dimension of the image of a map in the same homotopy class. Our goal is then to find the maximum homotopical rank of maps $f : M \rightarrow N$, where M is a compact Kähler manifold and N is a compact, locally symmetric manifold covered by X . Since it is known that this number is strictly smaller than the dimension of X , cf. Cor. (3.3) of [5], it remains to find the sharp upper bound, and to classify the maps that attain this bound. A reasonable way to do this is to answer the following.

Suppose $X_1 = G_1/K_1$ is a Hermitian symmetric space that can be totally geodesically embedded in X , and is of maximum dimension among all such Hermitian symmetric spaces.

Question 1: Is the homotopical rank of f at most the dimension of X_1 ?

Question 2: If $f : M \rightarrow N$ has homotopical rank equal to the dimension of X_1 , must X_1 have a compact locally symmetric quotient N_1 which admits a totally geodesically

Research partially supported by the National Science Foundation

immersion $g : N_1 \rightarrow N$ so that f factors as $f = gh$ for some holomorphic map $h : M \rightarrow N_1$?

Both questions have affirmative answers when the symmetric space X_1 is embedded in X as the fixed point set of an involution, provided that a small list of exceptions is excluded. More precisely, we prove the following:

1.1. Theorem. *Let $X = G/K$ where G is one of the groups $SL(2n, \mathbb{R})$, $n \geq 3$, $SO^+(2p, 2q)$, $p, q \geq 3$, $(p, q) \neq (3, 3)$, $SU^*(2n)$, $n \geq 3$ and $Sp(p, q)$, $p, q \geq 2$. Then the answer to both questions is affirmative, where $X_1 = G_1/K_1$ is embedded in X by the natural inclusion of $G_1 = Sp(n, \mathbb{R})$, $U(p, q)$, $SO^*(2n)$ and $U(p, q)$ in G , respectively.*

Question 1 has an affirmative answer for all X as in the first paragraph, provided that $\dim(X_1)$ is replaced by $\dim(X_1) + 2$. Then the sharp bound is either $\dim(X_1)$ or $\dim(X_1) + 2$. Just which alternative holds cannot, however, be decided by the methods of this paper. Problems of a global nature which bear on this question are discussed in §9. In any case, the bound that we obtain is much lower than the dimension of X , usually about $1/2 \dim(X)$, and so is a considerable improvement on [5], Cor. (3.2).

Observe that if in Theorem (1.1) we had also assumed that M is locally symmetric, then the conclusion could easily have been derived from the rigidity theorem of Margulis [16]. The point of Theorem (1.1) is therefore that the only assumption on M is that it is a compact Kähler manifold. Consequently this theorem is in the same relation to the Margulis rigidity theorem as Siu's rigidity theorem is to Mostow's rigidity theorem, cf. [18]. [23].

The method of proof follows the pattern pioneered by Siu in the proof of his rigidity theorem [23]. Since N has non-positive curvature, the existence theorem of Eells and Sampson [9] applies and the map f may be assumed to be harmonic. Its rank therefore gives an upper bound on the homotopical rank. Since M is compact Kähler and the curvature tensor of N is rather special, any harmonic map of M to N is actually pluriharmonic, i.e., the restriction of f to any local complex curve in M is harmonic [19]. If $\dim(M) > 1$, the resulting system of partial differential equations satisfied by f is overdetermined, and can be very effectively studied by just considering the two-jets of a solution. The technical content of this paper is in fact mostly the study of the two-jet of a pluriharmonic mapping of a complex manifold to a symmetric space whose one-jet has maximum possible rank. The results of our study will then have immediate topological consequences to the study of homotopical rank of continuous maps.

A key first-order consequence of pluriharmonicity is the following [19]. For each $x \in M$, $df(T_x^{1,0}M)$ is an isotropic subspace of $T_{f(x)}^{\mathbb{C}}N$ with respect to the complex multilinear extension of the curvature tensor R of N , i.e., $R(X, Y, X, Y) = 0$ for all $X, Y \in df(T_x^{1,0}M)$. Thus, if $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$ is a Cartan decomposition of the Lie algebra of G , where \mathfrak{k} is the isotropy algebra at $f(x)$, then \mathfrak{p} is naturally isomorphic to $T_{f(x)}N$, and $df(T_x^{1,0}M)$ corresponds to a subspace W of $\mathfrak{p}^{\mathbb{C}}$. The above isotropy condition is then equivalent to

the assertion that W is abelian: $[W, W] = 0$. It is therefore natural to study pluriharmonic maps of maximum rank via the following three-step program:

- 1) For each real simple Lie algebra \mathfrak{g} with Cartan decomposition $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$, find the maximum dimension of $W \subset \mathfrak{p}^{\mathbb{C}}$, W abelian.
- 2) Classify, up to automorphism of \mathfrak{g} , all abelian subalgebras of maximum dimension.
- 3) Use (2) to find normal forms for the pluriharmonic maps of maximum possible rank.

We carry out step 1 for all the classical real simple Lie algebras for which the problem has not been previously solved. These are the classical real simple Lie algebras whose complexification is also simple (i.e., corresponding to the symmetric spaces of type III in Helgason's terminology, [13], p. 379), which means that the corresponding symmetric space is not Hermitian symmetric and has rank at least two. These are the Lie algebras of the groups $SL(n, \mathbb{R})$, $n > 2$, $SO(p, q)$, $p, q > 2$, $SU^*(2n)$, $n > 2$, and $Sp(p, q)$, $p, q > 1$.

It turns out that for some of these algebras an abelian subalgebra of $\mathfrak{p}^{\mathbb{C}}$ of maximum dimension is obtained in a very geometric way: it is the space of $(1, 0)$ tangent vectors to a Hermitian symmetric subspace G_1/K_1 which is fixed by an involution of G/K . (The integrability condition for the complex structure on G_1/K_1 is equivalent to assertion that the space of $(1, 0)$ -tangent vectors is abelian). Such algebras can be locally realized by pluriharmonic maps, and can be realized globally by taking suitable quotients by discrete subgroups, where for this last step one uses [2]. We call these "algebras of Type I" (cf. §2) and carry out steps 2 and 3 only for the subclass listed in the statement of Theorem 1.1.

The remaining algebras in our list we call "algebras of Type II", cf. §2 for the precise definition. For these algebras (with the exception of $SO(6, 6)$) the largest dimension of an abelian subalgebra of $\mathfrak{p}^{\mathbb{C}}$ is one more than the largest dimension of a Hermitian symmetric subspace, and in most cases this subspace is no longer fixed by an involution. The question of realizability of the largest abelian subalgebra by the tangent space of a pluriharmonic map is discussed in §9.

So far we have stated the consequences of our local results in terms of mappings between compact manifolds. However, the present state of the existence theory of harmonic mappings (cf. §2 of [8] for a recent survey) makes it clear that the compactness of the target is not essential, and, more generally, that the map can be replaced by a harmonic section of a flat G/K -bundle. In particular, the existence theorem of Corlette for harmonic sections [6], combined with our results, gives immediate applications to representations of fundamental groups of compact Kähler manifolds into the Lie groups in question. Thus, if $\rho : \pi_1(M) \rightarrow G$ is a representation, where G is as in Theorem 1 and the harmonic section of ρ has rank = $\dim(X_1)$, then the image of ρ must be contained in an embedding of an appropriate subgroup G_1 .

In this context it is natural to use characteristic classes to give lower bounds on the rank of a harmonic section (even though a notion of homotopical rank still applies). We treat an example for the group $SO^+(2p, 2q)$: given $\rho : \pi_1(M) \rightarrow SO^+(2p, 2q)$, the associated flat bundle E with fibre \mathbb{R}^{2p+2q} is isomorphic, as a vector bundle, to a direct sum

$E_+ \oplus E_-$ of bundles with fiber \mathbb{R}^{2p} and \mathbb{R}^{2q} respectively, namely the maximal positive and negative sub-bundles for the indefinite inner product. Neither of these bundles is necessarily flat. In particular, if $e_+ \in H^{2p}(M)$ denotes the Euler class of E_+ , then e_+ does not necessarily vanish, and its powers can be used to give lower bounds for the rank of a harmonic section:

1.2. Theorem. *Let M be a compact Kähler manifold, let $\rho : \pi_1(M) \rightarrow SO^+(2p, 2q)$, $p, q \geq 3$, $(p, q) \neq (3, 3)$, be a representation, and let $e_+ \in H^{2p}(M)$ be the cohomology class associated to ρ as above. If $e_+^q \neq 0$, then the image of ρ is contained in some embedding of $U(p, q)$ in $SO^+(2p, 2q)$.*

The reason we restrict ourselves to the algebras listed above is that all the other classical cases are known. If the Lie algebra is that of the automorphism group of a Hermitian symmetric space then the type of problems that we study have all been answered by Siu [23] and are equivalent to his rigidity theorem. If \mathfrak{g} is simple but $\mathfrak{g}^{\mathbb{C}}$ is not simple, then \mathfrak{g} is a simple complex Lie algebra regarded as a real Lie algebra. The corresponding symmetric spaces are those of the form G/G_u , where G is a simple complex Lie group and G_u is its compact real form, i.e., the symmetric spaces of type IV ([13], p. 379). In this case the abelian subalgebras of $\mathfrak{p}^{\mathbb{C}}$ are the same as the abelian subalgebras of the complex simple Lie algebra \mathfrak{g} . In this case problems 1 and 2 have long been solved, namely by Schur [20] for $\mathfrak{sl}(n, \mathbb{C})$ and by Malcev [15] for all the complex simple Lie algebras. The solution of problem 3 above is contained in the paper of Corlette [7] (for the classical \mathfrak{g} and two of the exceptional ones). Finally, if \mathfrak{g} is classical, of real rank one, and not already included in the above cases, then either $G = SO^+(1, n)$, $n \geq 4$, or $G = Sp(1, n)$, $n \geq 2$, which are treated in [19], [5].

From the point of view of thoroughness, there remain only some of the exceptional Lie algebras. For the twelve exceptional real Lie algebras whose complexification is simple, two are Hermitian symmetric, hence included in [24], while real form of F_4 corresponding to the Cayley plane is treated in [3]. Consequently there remain nine algebras for which the questions 1, 2, 3 above have not been studied. For the exceptional simple complex Lie algebras there remain three for which question 3 has not been studied.

We emphasize that in this paper we study only the pluriharmonic mappings which have maximum possible rank. Other classes of pluriharmonic maps are: (1) those that factor through any totally geodesic Hermitian subspace, (2) those that factor through Riemann surfaces, and (3) those that factor through totally geodesic real tori. It is natural to ask if there are additional, nonobvious classes. It turns out that there is at least one other construction, which comes from algebraic geometry: the theory of variations of Hodge structure gives natural examples of mappings of algebraic surfaces into quotients of symmetric spaces of suitable groups of type $SO(p, q)$ that do not fall into any of the above categories.

Our interest in variations of Hodge structure originates with our observation in [5], §6, that they are a substitute, for non-Hermitian symmetric targets, for holomorphic maps to Hermitian symmetric targets. In particular, for each non-Hermitian X one

would expect a number $r(X)$ so that every pluri-harmonic map to X of rank larger than $r(X)$ lifts to a variation of Hodge structure, extending our result in [5], Theorem (6.1) for X the quaternionic hyperbolic space. The reason for this expectation is its analogy with similar rank results of Siu for Hermitian targets, conjectured in §6 of [24] and proved in [25]. We present a precise result of this nature in §9.

Finally, we mention that Simpson [22] has recently constructed other classes of harmonic mappings that are neither variations of Hodge structure nor fall into any of the three obvious classes mentioned above. The resulting abelian spaces consist of semi-simple elements and are therefore quite different in nature from the cases we study.

The paper is organized as follows. In §2 we give a detailed outline of the proofs of steps 1-3, and carry out the details in §3–7. In §8 we present the example from Hodge theory, and in the final section §9 we discuss further results and open problems.

As we have already mentioned, this paper and Corlette’s paper [7] have complementary results concerning the classification of harmonic mappings. We have also borrowed methods from each other in obtaining and presenting the results, and we are very grateful to Corlette for his contributions to this paper. We thank O. Gabber for explaining to us the relevance cohomological methods and Kostant’s theorem to rigidity. We also thank H. Hecht and P. Trombi for help in Lie algebra matters, A. Beauville, P. Deligne, M. Gromov, and R. Varley for enlightening remarks concerning the examples in §8, and Carlos Simpson and Kevin Corlette for their insightful comments regarding Theorem 9.3. Finally we are grateful to the IHES for its hospitality while a good portion of this work was in progress.

§ 2. Outline of the proof

Let \mathfrak{g} be a real simple Lie algebra as in the introduction. In particular, $\mathfrak{g}^{\mathbb{C}}$ is simple and \mathfrak{g} is not the Lie algebra of the automorphism group of a symmetric domain. In this section we sketch the methods that we use to carry the steps 1 to 3 described in the introduction.

Step 1: Dimension bounds. Let W be an abelian subalgebra of $\mathfrak{g}^{\mathbb{C}}$ which lies in $\mathfrak{p}^{\mathbb{C}}$. Without loss of generality we may assume that W is also maximal with respect to inclusion among such algebras. The problem is to bound the dimension of W , and this can be accomplished by an elaboration of the general outline of Malcev [15], which consists of three steps:

Part a: Reduction to nilpotents. The space W has a Jordan decomposition

$$W = W_{\sigma} \oplus W_{\nu},$$

where the first space consists of semisimple elements and the second of nilpotents. In addition, we may assume, after possible conjugation by an element of $K^{\mathbb{C}}$, that W_{σ} is

defined over the reals, cf. [5], Lemma (4.2). Then the centralizer of W_σ is the complexification reductive Lie subalgebra of \mathfrak{g} , invariant under the Cartan involution, which decomposes as $\mathfrak{z} \oplus \mathfrak{l}$, where \mathfrak{z} is the center and where \mathfrak{l} is semisimple. Furthermore $W_\sigma \subset \mathfrak{z}^\mathbb{C}$, and if $\mathfrak{l} = \oplus \mathfrak{l}_i$ is a decomposition into simple ideals, then each \mathfrak{l}_i is the Lie algebra of a sub-symmetric space, with Cartan decomposition $\mathfrak{l}_i = \mathfrak{k}_i \oplus \mathfrak{p}_i$, and $W_\nu = \oplus W_i$, where $W_i = W_\nu \cap \mathfrak{p}_i^\mathbb{C}$. We then have

$$W = W_\sigma \oplus W_1 \oplus \cdots \oplus W_k$$

where each $W_i \subset \mathfrak{p}_i^\mathbb{C}$ consists of nilpotent elements, cf. §4 of [5] for details.

In §6 we find all the possible subalgebras \mathfrak{l}_i that can occur for an algebra \mathfrak{g} in our list. It turns out that possible subalgebras \mathfrak{l}_i are also essentially in our list (we only have to add some of the cases excluded as “already known”). Thus, the problem is reduced to bounding the dimension of the abelian subalgebras of $\mathfrak{p}^\mathbb{C}$ consisting entirely of nilpotent elements for the algebras \mathfrak{g} of the following groups: $SL(n, \mathbb{R})$, $n \geq 2$, $SU^*(2n)$, $n \geq 2$, $SO(p, q)$ and $Sp(p, q)$, p, q arbitrary.

Part b: Reduction to commutative systems of non-compact roots. Let us assume, therefore, that W consists entirely of nilpotent elements. By Lemma 3.2, there is a Cartan subalgebra \mathfrak{t} of \mathfrak{k} with the following properties. First the eigenvalues of the adjoint action of \mathfrak{t} on $\mathfrak{g}^\mathbb{C}$ form a root system $\Delta \subset \mathfrak{t}^*$. For each $\alpha \in \Delta$ let \mathfrak{g}_α denote the corresponding eigenspace. Then there is a system of positive roots $\Delta^+ \subset \Delta$ such that W is contained in the sum of the positive root spaces:

$$W \subset \sum_{\alpha \in \Delta^+} \mathfrak{g}_\alpha.$$

A root system Δ with this property will be called *adapted to W* . Because the dimension of \mathfrak{t} may be smaller than the rank of $\mathfrak{g}^\mathbb{C}$, the adapted root system may be a restricted one. In particular, the spaces \mathfrak{g}_α may have dimension greater than one. However, since \mathfrak{t} is invariant under the Cartan involution, so are the \mathfrak{g}_α , and consequently these spaces decompose into compact and noncompact parts: $\mathfrak{g}_\alpha = \mathfrak{k}_\alpha \oplus \mathfrak{p}_\alpha$, where $\mathfrak{k}_\alpha \subset \mathfrak{k}^\mathbb{C}$ and $\mathfrak{p}_\alpha \subset \mathfrak{p}^\mathbb{C}$. Roots have multiplicity at most two, and when the multiplicity equals two, both terms in the above decomposition of \mathfrak{g}_α are nonzero. In any case, there are sets

$$\begin{aligned} \Delta(\mathfrak{k}) &= \{ \alpha \in \Delta \mid \mathfrak{k}_\alpha \neq 0 \} \\ \Delta(\mathfrak{p}) &= \{ \alpha \in \Delta \mid \mathfrak{p}_\alpha \neq 0 \}, \end{aligned}$$

where the first is a root system for the compact subalgebra \mathfrak{k} , and where

$$\mathfrak{p}^\mathbb{C} = \sum_{\alpha \in \Delta(\mathfrak{p})} \mathfrak{p}_\alpha.$$

The sets $\Delta(\mathfrak{k})$ and $\Delta(\mathfrak{p})$ are disjoint if and only if the all roots have multiplicity one. We shall use the term “compact” and “noncompact” to refer to elements of either set, bearing in mind that a root can be both compact and noncompact. Let $\Delta^+(\mathfrak{p}) = \Delta^+ \cap \Delta(\mathfrak{p})$, and let

$$\mathfrak{p}^+ = \sum_{\alpha \in \Delta^+(\mathfrak{p})} \mathfrak{p}_\alpha.$$

Then $W \subset \mathfrak{p}^+$.

Now choose a total ordering of the roots which is compatible with addition and with the partial ordering defined by Δ^+ . Using Gaussian elimination, one can construct a basis $\{Z_i, i = 1, \dots, d\}$ for W which has the form $Z_i = X_i + Y_i$, where

- i) $X_i \in \mathfrak{p}_{\alpha(i)}$,
- ii) $\alpha(i) < \alpha(j)$ for $i < j$.
- iii) Y_i is a linear combination of vectors $X_\beta \in \mathfrak{p}_\beta$ with $\beta > \alpha(i)$ and $\beta \notin \{\alpha(1), \dots, \alpha(d)\}$.

Let $C = \{\alpha(i) : i = 1, \dots, d\}$. The elements of C are called the *leading roots* of W ; they depend upon the choice of a total order.

Now form the Lie bracket of two basis vectors:

$$[Z_i, Z_j] = [X_i, X_j] + ([X_i, Y_j] + [Y_i, X_j]) + [Y_i, Y_j]. \quad (2.1)$$

By construction, the first term in the right-hand side belongs to the space $\mathfrak{p}_{\alpha(i)+\alpha(j)}$, while the remaining terms belong to the space

$$\sum_{\beta > \alpha(i)+\alpha(j)} \mathfrak{p}_\beta.$$

Therefore the vanishing of $[Z_i, Z_j]$ implies the vanishing of $[X_i, X_j]$. Thus if we let W_C denote the space spanned by the X_i , then W_C is an *abelian* subspace of $\mathfrak{p}^{\mathbb{C}}$ of the same dimension as W . Moreover, W_C , being a direct sum of the root spaces \mathfrak{p}_α , $\alpha \in C$, is abelian if and only if the set C of leading roots is *commutative*, meaning that the sum of any two elements of C is not a root. Thus we are reduced to the purely combinatorial problem of *finding commutative systems of non-compact roots of maximum cardinality*.

In tables 1 and 2 of §4 we list the root systems Δ and positive roots Δ^+ for all the algebras listed in (a), and in §5 we solve the problem of finding the commutative systems of maximum cardinality for each $\Delta^+(\mathfrak{p})$. There is only one technical complication in §4,5. Given a root system Δ , all choices Δ^+ of a system of positive roots are equivalent by an automorphism of Δ . But this automorphism may not preserve the decomposition of Δ into compact and non-compact roots, ie, the corresponding automorphism of \mathfrak{g} need not commute with the Cartan involution. Thus for each root system Δ we have to consider all the choices of Δ^+ modulo automorphisms that preserve the Cartan decomposition. The equivalence classes of choices are parametrized by $Aut(\Delta)/Aut(\Delta(\mathfrak{k}))$. This set has cardinality one for $SL(n, \mathbb{R})$ and $SU^*(2n)$, which we list in Table 1; but has

larger cardinality for $SO(p, q)$ and $Sp(p, q)$, which we list in Table 2. The main technical difficulty in §5 arises from the fact that, for the groups $SO(p, q)$ and small values of p or q , the maximum cardinality of C is attained in non-obvious cosets, which do not follow the same pattern as for p, q large.

Part c: Combination of (a) and (b). Given these results one can compute a bound on the dimension of any abelian space W which decomposes as

$$W = W_\sigma \oplus W_1 \oplus \cdots \oplus W_k,$$

where the first term consists of semisimple elements and the remaining ones consist of nilpotents for an irreducible symmetric pair (G_i, K_i) . The required bound is the maximum of these over all possible decompositions. What one finds *a posteriori* is that the maximum is always achieved by a space consisting entirely of nilpotents in the single irreducible symmetric pair G/K . Moreover, this maximum is achieved only in this way except for $SL(4, \mathbb{R})$, $SL(2n+1, \mathbb{R})$ and $SO(2p+1, 2q+1)$. In these cases there is also a space with a single semi-simple element achieving the maximum. In the case of $SL(3, \mathbb{R})$ there is, in addition, a two-dimensional space of semi-simple elements which achieves the maximum dimension.

To explain Steps 2 and 3, we first need to define a division of the root systems into two types, which we call types I and II, defined as follows. Let C be a commutative system of maximum cardinality, and let W_C be the corresponding commutative space. Let \mathfrak{g}' denote the subalgebra of \mathfrak{g} generated by the real points of the subspace $W_C + \bar{W}_C \subset \mathfrak{p}^{\mathbb{C}}$. Then a case by case verification shows that there are only two possibilities, with all the properties now listed:

Type I. \mathfrak{g}' corresponds to a Hermitian symmetric subspace. This happens precisely when $W_C + \bar{W}_C = \mathfrak{p}_1^{\mathbb{C}}$, where $\mathfrak{p}_1 \subset \mathfrak{p}$ is a Lie triple system, thus corresponds to a totally geodesic subspace, which is necessarily Hermitian symmetric, since $\mathfrak{p}_1^{\mathbb{C}}$ contains an abelian subalgebra of half its dimension, namely W_C . Thus for type I, $W_C = \mathfrak{p}_1^{1,0}$, the space of (1,0)-tangent vectors to an invariant complex structure on \mathfrak{p}_1 , which for ease of notation we denote by \mathfrak{p}_1^+ . Moreover, this symmetric subspace is the fixed point set of an involution. Thus, the Lie algebra \mathfrak{g} has two commuting involutions σ, τ , where the first is the Cartan involution, which give a $\mathbb{Z}/2 \times \mathbb{Z}/2$ -grading of \mathfrak{g} , i.e., a four-term decomposition

$$\mathfrak{g} = \mathfrak{k}_1 \oplus \mathfrak{m} \oplus \mathfrak{p}_1 \oplus \mathfrak{q}, \tag{2.2}$$

refining the Cartan decomposition of \mathfrak{g} where the terms have (σ, τ) -types $(1, 1)$, $(1, -1)$, $(-1, 1)$, and $(-1, -1)$, respectively. If we let $\mathfrak{g}_1 = \mathfrak{k}_1 \oplus \mathfrak{p}_1$, we have $\mathfrak{g}' \subset \mathfrak{g}_1$, with equality except in the cases where $\mathfrak{g}' = \mathfrak{su}(p, q)$, in which case $\mathfrak{g}_1 = \mathfrak{u}(p, q)$. The groups in question are $Sp(p, q)$, $SO(2p, 2q)$ for $p, q \geq 3$, $SL(2n, \mathbb{R})$ for $n \geq 2$, $SU^*(2n)$ for $n \geq 2$.

Type II. \mathfrak{g}' does not correspond to a Hermitian symmetric subspace. In these cases \mathfrak{g}' is very large, usually equal to \mathfrak{g} , and the commutative system C of maximum cardinality contains a subsystem C' of cardinality one less, so that the corresponding commutative

space $W_{C'}$ is of the form \mathfrak{p}_1^+ for a Lie triple system $\mathfrak{p}_1 \subset \mathfrak{p}$, which is necessarily Hermitian symmetric but, in most cases, is no longer the fixed point set of an involution. The groups in question are complementary, in our list, to those listed under Type I, except for the group $SO(6,6)$. The latter has two inequivalent systems of largest cardinality 9, one which falls under Type I, the other under Type II.

In the remaining steps we assume that the algebra is of Type I.

Step 2: First order rigidity. Let $W \subset \mathfrak{p}^{\mathbb{C}}$ be an abelian subalgebra of maximum dimension and consisting entirely of nilpotent elements (the second assumption is unnecessary for all groups of Type I except $SL(4, \mathbb{R})$). Let C be its commutative system of leading roots as in Step 1b. Using the notation established there, let $W_C \subset \mathfrak{p}^{\mathbb{C}}$ be the direct sum of the corresponding root spaces, with basis $\{X_i\}$, where $\{Z_i = X_i + Y_i\}$ is a basis for W . Then, as in the definition of Type I, $W_C = \mathfrak{p}_1^+$, and property (iii) of Step 1b implies that $Y_i \in \mathfrak{q}^{\mathbb{C}}$. From the evident multiplicative properties of the $\mathbb{Z}/2 \times \mathbb{Z}/2$ -grading (2.2), it follows that the second term of the right-hand side of (2.1) belongs to $\mathfrak{m}^{\mathbb{C}}$ and the third term belongs to $\mathfrak{k}_1^{\mathbb{C}}$, hence each term vanishes. Thus, if we define a linear map $\phi : \mathfrak{p}_1^+ \rightarrow \mathfrak{q}^+$, where $\mathfrak{q}^+ = \mathfrak{q}^{\mathbb{C}} \cap \mathfrak{p}^+$, by letting $\phi(X_i) = Y_i$, then

$$W = \{X + \phi(X) : X \in \mathfrak{p}_1^+\},$$

and the vanishing of the second term in the right-hand side of (2.1) is equivalent to the following cocycle condition for ϕ :

$$[X, \phi(Y)] + [\phi(X), Y] = 0 \quad \text{for all } X, Y \in \mathfrak{p}_1^+. \quad (2.3)$$

This equation means geometrically that $W_t = \{X + t\phi(X) : X \in \mathfrak{p}_1^+\}$ is a deformation of W to $W_C = \mathfrak{p}_1^+$, and cohomologically means the following. Let $V \subset \mathfrak{g}^{\mathbb{C}}$ be a subspace which is stable under the Cartan involution, is a \mathfrak{p}_1^+ -module, i.e., $[\mathfrak{p}_1^+, V] \subset V$, and which contains the image of ϕ . Then $\phi \in \text{Hom}(\mathfrak{p}_1^+, V) = C^1(\mathfrak{p}_1^+, V)$ and (2.3) is equivalent to ϕ being a cocycle. Suppose that its class in $H^1(\mathfrak{p}_1^+, V)$ is zero. This means that there is and $X_0 \in V$ such that

$$\phi(X) = [X, X_0] \quad \text{for all } X \in \mathfrak{p}_1^+. \quad (2.4)$$

Moreover, it is clear that $X_0 \in V \cap \mathfrak{k}^{\mathbb{C}}$, thus $\exp(X_0)$ gives a $K^{\mathbb{C}}$ -conjugacy between W and \mathfrak{p}_1^+ , i.e., \mathfrak{p}_1^+ is rigid as an abelian subspace of $\mathfrak{p}^{\mathbb{C}}$.

A module V that satisfies these properties is $\mathfrak{m}^{\mathbb{C}} \oplus \mathfrak{q}^{\mathbb{C}}$, and in §7 we use essentially this module together with a well-known theorem of Kostant [10], to prove that for most algebras of Type I the cohomology class of ϕ vanishes. We also show that the $K^{\mathbb{C}}$ -conjugacy provided by (2.4) is actually a K -conjugacy. We conclude that if \mathfrak{g} is of Type I and $f : M \rightarrow G/K$ is a pluri-harmonic map of maximum rank, then for each $x \in M$ there is a totally geodesic Hermitian symmetric subspace $X_x \subset G/K$ containing $f(x)$ so that $df(T_x M) = T_{f(x)} X_x$. We call this *first-order rigidity of f* .

We remark that at this point the restriction to Type I may be mostly for convenience, in order to apply cohomological methods to replace the more involved combinatorial arguments of [15], where algebras analogous to our Type II (e.g., B_n) also occur, and are handled directly. Since the next and final step is out of reach of the present methods, this is at the moment not a serious restriction.

Step 3: Global rigidity. *This is the only step where we use the second order information in the pluri-harmonic equation.* We assume that $f : M \rightarrow G/K$ is a pluriharmonic map of maximum rank for which first-order rigidity has been proved in Step 2. At each point it is tangent to a totally geodesic Hermitian symmetric subspace, so second-order rigidity is equivalent to local rigidity or global rigidity.

We consider first the local problem. Suppose that $U \subset M$ is a connected open set, let $X = G/K$, and suppose that $f : U \rightarrow X$ is a smooth map. We regard X as the manifold of isotropy subalgebras $\{\mathfrak{k}_x \subset \mathfrak{g} : x \in X\}$. Then there are tautological subbundles $\mathfrak{k}_X, \mathfrak{p}_X$ of the trivial bundle $\mathfrak{g}_X = X \times \mathfrak{g}$, with fibre over x the isotropy subalgebra \mathfrak{k}_x and its canonical complement \mathfrak{p}_x respectively. Moreover the flat connection on \mathfrak{g}_X corresponding to the trivialization splits (by projection to the two components) as the direct sum of a K -connection and a one-form with values in \mathfrak{p}_X . The bundle $f^*\mathfrak{g}_X$ splits as $f^*\mathfrak{k}_X \oplus f^*\mathfrak{p}_X$, and the induced flat connection splits accordingly. If U is simply connected, the datum of the induced bundle and its decomposition and the decomposition of the flat connection determine the map f uniquely up to left translation by an element of G . Thus in studying the local problem we will replace the map f by a simply connected open set U , a flat Lie algebra bundle \mathfrak{g}_U over U , a (non-flat) decomposition $\mathfrak{g}_U = \mathfrak{k}_U \oplus \mathfrak{p}_U$, and a decomposition of the flat connection d on \mathfrak{g}_U as

$$d = D + \theta \tag{2.5}$$

where D is a K -connection on \mathfrak{g}_U and $\theta \in A^1(U, \mathfrak{p}_U)$. Then θ replaces df , and the pluri-harmonic equation $D''d'f = 0$, cf. [23], [19], [5], for f becomes

$$D''\theta' = 0, \tag{2.6}$$

where $D = D' + D''$ and $\theta = \theta' + \theta''$ are the decompositions of D and θ into (1,0) and (0,1) components respectively.

Now our assumption is that for some $x \in U$, $\theta'(T_x^{1,0}U)$ has maximum possible dimension. Since, by (2.6), θ' is holomorphic, θ' has maximum possible rank in the complement of a complex analytic subvariety of U . Call this set V and observe that it is connected, open, and dense in U . Thus if G is of Type I, and first-order rigidity holds, for each $x \in V$, $\theta'(T_x^{1,0}V) = \mathfrak{p}_1^+(x)$, where $\mathfrak{p}_1(x)$ is the tangent space to a totally geodesic Hermitian symmetric subspace of G/K fixed by an involution. For each $x \in V$ we get a decomposition (2.2) of \mathfrak{g}_x , and a corresponding decomposition of the bundle \mathfrak{g}_V , denoted by a subscript V under each term of (2.2).

The geometric interpretation of this decomposition is the following. There is a fibration $G/K_1 \rightarrow G/K$, and the fibre over $y \in G/K$ parametrizes the tangent spaces

at y to totally geodesic embeddings of G_1/K_1 in G/K passing through y . The pull-back to G/K_1 of the bundle \mathfrak{g}_X over G/K splits according to (2.2), and the sum of the last three terms is isomorphic to the tangent bundle of G/K_1 . The map $f : V \rightarrow G/K$ lifts to a map $F : V \rightarrow G/K_1$ by letting $F(x) = df(T_x V)$, which makes sense since by assumption $df(T_x V)$ is the tangent space to a totally geodesic embedding of G_1/K_1 passing through $f(x)$, hence is a point of G/K_1 lying over $f(x)$. The connection D in (2.5) splits as $D = D_1 + \mu$, where D_1 is a K_1 -connection and $\mu \in A^1(V, \mathfrak{m}_V)$. Thus the original flat connection d of (2.5) splits further as

$$d = D_1 + \mu + \theta. \quad (2.7)$$

Observe that in this notation, $dF = \mu + \theta$.

Now the orbits of G_1 on G/K_1 foliate this manifold into leaves isomorphic to G_1/K_1 which project to the different geodesic embeddings of G_1/K_1 in G/K . We want to show that $f(U)$ is contained in a single such embedding. Since V is dense in U , this is equivalent to showing that $F(V)$ is contained in a single leaf of the foliation. Since V is connected, this is equivalent to $F(V)$ being tangent to the foliation. By construction the one-form θ has values in $(\mathfrak{p}_1)_V$, which is the tangent bundle to the foliation, so it suffices to show that $\mu = 0$. This is accomplished by the equations that μ must satisfy. First the pluriharmonic equation (2.6) splits into two equations:

$$D_1''\theta' = 0 \quad \text{and} \quad [\mu'', \theta'] = 0 \quad (2.8)$$

and the flatness of the original connection d in (2.5), when decomposed further into types and also according to (2.2) gives the further equation

$$[\mu', \theta'] = 0. \quad (2.9)$$

In §7 we show that the second equation (2.8) together with (2.9) have a cohomological interpretation, and that the vanishing of μ can be proved by appealing again to Kostant's theorem. This shows the local rigidity of f , from which the global rigidity is immediate (cover M by simply connected open sets U as above with connected intersections).

§ 3. Existence of adapted root systems

We now prove the technical result that allows us to put W in upper triangular form in a way that is compatible with the Cartan involution. We owe the following lemma to Henryk Hecht.

3.1. Lemma. *Let \mathfrak{g} be a real semisimple Lie algebra, let θ denote its Cartan involution, and let W be a θ -stable subalgebra of $\mathfrak{g}^{\mathbb{C}}$, each element of which is nilpotent. Then there is a fundamental Cartan subalgebra \mathfrak{h} of \mathfrak{g} and a system of positive roots $\Delta^+ \subset \Delta(\mathfrak{g}, \mathfrak{h})$ such that*

$$W \subset \mathfrak{n}^+ = \sum_{\alpha \in \Delta^+} \mathfrak{g}_{\alpha}.$$

We recall that a Cartan subalgebra \mathfrak{h} of \mathfrak{g} is called *fundamental* if it is stable under the Cartan involution, and, if $\mathfrak{h} = \mathfrak{h}_+ \oplus \mathfrak{h}_-$ denotes its decomposition into ± 1 eigenspaces for θ , then its compact part \mathfrak{h}_+ has maximum possible dimension; i.e., \mathfrak{h}_+ is a Cartan subalgebra for \mathfrak{k} . Similarly, a Cartan subalgebra \mathfrak{h} of $\mathfrak{g}^{\mathbb{C}}$ is called *fundamental* if \mathfrak{h}_+ is a Cartan subalgebra of $\mathfrak{k}^{\mathbb{C}}$.

Let \mathfrak{h} be a fundamental Cartan subalgebra of \mathfrak{g} , let $\mathfrak{t} = \mathfrak{h}_+$, and let $\Delta(\mathfrak{g}, \mathfrak{h})$ denote the roots for the adjoint action of \mathfrak{h} on $\mathfrak{g}^{\mathbb{C}}$. It is a standard and simple fact that the maximality of \mathfrak{h}_+ implies that no root $\alpha \in \Delta(\mathfrak{g}, \mathfrak{h})$ restricts to zero on \mathfrak{t} . From general principles it follows that the set of restrictions to \mathfrak{t} of the elements of $\Delta(\mathfrak{g}, \mathfrak{h})$ forms a root system on \mathfrak{t} , which we will denote by $\Delta(\mathfrak{g}, \mathfrak{t})$. Decomposing the resulting root spaces into compact and noncompact parts as $\mathfrak{g}_\alpha = \mathfrak{k}_\alpha \oplus \mathfrak{p}_\alpha$, and observing that the restriction of a positive system of roots for \mathfrak{h} is a positive system of roots for \mathfrak{t} , we obtain the following related form of the lemma. We state it only for abelian subalgebras of $\mathfrak{p}^{\mathbb{C}}$, rather than more general θ -stable subalgebras, since this is the form that we use.

3.2. Lemma. *Let W be an abelian subspace of $\mathfrak{p}^{\mathbb{C}}$ which consists entirely of nilpotent elements. Then there is a Cartan subalgebra \mathfrak{t} of \mathfrak{k} and a system of positive roots $\Delta^+(\mathfrak{g}, \mathfrak{t})$ such that*

$$W \subset \sum_{\alpha \in \Delta^+(\mathfrak{g}, \mathfrak{t})} \mathfrak{p}_\alpha$$

Proof of Lemma A: Since W is a subalgebra consisting entirely of nilpotent elements, it is contained in the nil-radical $\mathfrak{n} = [\mathfrak{b}, \mathfrak{b}]$ of some Borel subalgebra \mathfrak{b} of $\mathfrak{g}^{\mathbb{C}}$. Applying Theorem 1(i) of Matsuki's paper [17] to the affine symmetric space $(G^{\mathbb{C}}, K^{\mathbb{C}}, \theta)$, it follows easily that there is a Cartan subalgebra \mathfrak{h} of $\mathfrak{g}^{\mathbb{C}}$ in \mathfrak{b} which is θ -stable. Write $\mathfrak{h} = \mathfrak{h}_+ \oplus \mathfrak{h}_-$, where the first term is in $\mathfrak{k}^{\mathbb{C}}$ and the second is in $\mathfrak{p}^{\mathbb{C}}$. Let Δ^+ be the system of positive roots in $\Delta(\mathfrak{g}, \mathfrak{h})$ corresponding to \mathfrak{b} . Since $W \subset \mathfrak{n} = [\mathfrak{b}, \mathfrak{b}]$, one can write a general element $X \in W$ as

$$X = \sum_{\alpha \in S} c_\alpha X_\alpha,$$

where the $S \subset \Delta^+$, $X_\alpha \in \mathfrak{g}_\alpha$, and $c_\alpha \neq 0$ for all $\alpha \in S$. Roots that appear in some such S are said to *belong* to W . We write Δ_W for the collection of all such roots. Now $\theta X \in W \subset \mathfrak{n}$ and

$$\theta X = \sum_{\alpha \in S} c'_\alpha X_{\theta\alpha},$$

where $c'_\alpha \neq 0$ for all $\alpha \in S$. Thus, if α belongs to W , then so does $\theta\alpha$. Define a special collection of positive roots which includes all of those belonging to W by setting

$$(\Delta^+)' = \{ \alpha \in \Delta^+ \mid \theta\alpha \in \Delta^+ \}.$$

We proceed to modify these choices to obtain a new Borel subalgebra of $\mathfrak{g}^{\mathbb{C}}$ containing W which is defined by a fundamental Cartan subalgebra of \mathfrak{g} .

To begin, we observe that there is a θ -stable real form $\mathfrak{h}_{\mathbb{R}}$ of \mathfrak{h} such that $\Delta(\mathfrak{g}, \mathfrak{h}) \subset \mathfrak{h}_{\mathbb{R}}^*$: take $\mathfrak{h}_{\mathbb{R}} = \{ H \in \mathfrak{h} \mid \alpha(H) \in \mathbb{R} \text{ for all } \alpha \in \Delta \}$. Let H be a regular element of $\mathfrak{h}_{\mathbb{R}}$, i.e., one not contained in any root hyperplane, and write $H = H_+ + H_-$, with $H_{\pm} \in (\mathfrak{h}_{\mathbb{R}})_{\pm}$. Choose H so that

$$\alpha(H_+) > 0 \text{ for all } \alpha \in (\Delta^+)' .$$

Recall that a root is *real* if $\theta\alpha = -\alpha$, or, equivalently, if $\alpha(\mathfrak{h}_+) = 0$. Let $\Delta_{\mathbb{R}}$ denote the set of real roots, let $\Delta_{\mathbb{R}}^+ = \Delta_{\mathbb{R}} \cap \Delta^+$ and define a set $P \subset \Delta(\mathfrak{g}, \mathfrak{h})$ by the conditions

$$\begin{aligned} & \text{if } \alpha \in \Delta_{\mathbb{R}}^+ \text{ then } \alpha \in P, \\ & \text{if } \alpha \notin \Delta_{\mathbb{R}} \text{ but } \alpha(H_+) > 0, \text{ then } \alpha \in P. \end{aligned}$$

Then $P \subset \Delta$ is a system of positive roots. Replacing Δ^+ by this new system P of positive roots, we may thus assume that the original Δ^+ satisfies the following conditions:

$$\begin{aligned} & \text{if } \alpha \in \Delta^+ \setminus \Delta_{\mathbb{R}} \text{ then } \theta\alpha \in \Delta^+ \setminus \Delta_{\mathbb{R}}, \\ & \text{if } \alpha \in \Delta^+ \setminus \Delta_{\mathbb{R}}, \beta \in \Delta_{\mathbb{R}}, \text{ and } \alpha + \beta \in \Delta, \text{ then } \alpha + \beta \in \Delta^+ \setminus \Delta_{\mathbb{R}}, \end{aligned} \tag{2.3}$$

in addition to the obvious inclusion $\Delta_W \subset \Delta^+ \setminus \Delta_{\mathbb{R}}$.

Now define θ -stable Lie algebras

$$\begin{aligned} \mathfrak{l} &= \mathfrak{h} \oplus \sum_{\alpha \in \Delta_{\mathbb{R}}} \mathfrak{g}_{\alpha}, \\ \mathfrak{u} &= \sum_{\alpha \in \Delta^+ \setminus \Delta_{\mathbb{R}}^+} \mathfrak{g}_{\alpha}, \end{aligned}$$

where \mathfrak{l} is reductive, \mathfrak{u} is nilpotent and $\mathfrak{q} = \mathfrak{l} \oplus \mathfrak{u}$ is parabolic. Then $W \subset \mathfrak{u}$, and by the second condition (2.3), \mathfrak{l} normalizes \mathfrak{u} . Now, let $\tilde{\mathfrak{h}}$ be a fundamental Cartan subalgebra of \mathfrak{l} . Since $\mathfrak{l}_+ = \mathfrak{l} \cap \mathfrak{k}^{\mathbb{C}}$ and $\mathfrak{k}^{\mathbb{C}}$ have the same rank, $\tilde{\mathfrak{h}}$ is a fundamental cartan subalgebra of $\mathfrak{g}^{\mathbb{C}}$, and since $\tilde{\mathfrak{h}}$ normalizes \mathfrak{u} , \mathfrak{u} is still a sum of root spaces for $\tilde{\mathfrak{h}}$.

Let Δ denote now the root system for $\mathfrak{g}^{\mathbb{C}}$ relative to $\tilde{\mathfrak{h}}$. Then $\Delta = \Delta_0 \cup \Delta_1$, where Δ_0 is the set of roots of $\tilde{\mathfrak{h}}$ belonging to \mathfrak{l} (a sub-system of Δ) and $\Delta_1 = \Delta \setminus \Delta_0$. Define now a positive root system $\Delta^+ \subset \Delta$ by

$$\Delta^+ = \Delta_0^+ \cup \Delta_1^+,$$

where Δ_0^+ is a positive root system for Δ_0 satisfying (2.3), and Δ_1^+ is the set of roots in Δ belonging to \mathfrak{u} . Then Δ^+ also satisfies the conditions(2.3). Let $\tilde{\mathfrak{b}}$ be the Borel subalgebra of $\mathfrak{g}^{\mathbb{C}}$ corresponding to this choice of $\tilde{\mathfrak{h}}$ and Δ^+ . Then $\tilde{\mathfrak{b}}$ is a Borel subalgebra of $\mathfrak{g}^{\mathbb{C}}$ containing W and defined relative to a fundamental Cartan subalgebra of $\mathfrak{g}^{\mathbb{C}}$. It only remains to show that $\tilde{\mathfrak{b}}$ is defined relative to a fundamental Cartan subalgebra of \mathfrak{g} .

To see this last point, observe that $\tilde{\mathfrak{b}}$ is defined relative to a fundamental Cartan subalgebra of $\mathfrak{g}^{\mathbb{C}}$ and a positive system Δ^+ satisfying the first condition (2.3). By Proposition 2 of [17], these are precisely the conditions for the $K^{\mathbb{C}}$ -orbit of $\tilde{\mathfrak{b}}$ to be closed in the flag variety $G^{\mathbb{C}}/B$ of Borel subalgebras of $\mathfrak{g}^{\mathbb{C}}$. But this is equivalent to saying that the K -orbit of $\tilde{\mathfrak{b}}$ is the same as its $K^{\mathbb{C}}$ -orbit. Since by a $K^{\mathbb{C}}$ -equivalence we can make $\tilde{\mathfrak{h}}$ defined over \mathbb{R} , it follows that $\tilde{\mathfrak{b}}$ contains a fundamental Cartan subalgebra of \mathfrak{g} . Finally, let $\mathfrak{h} \subset \mathfrak{g}^{\mathbb{C}}$ now denote this subalgebra, and let Δ^+ denote the positive root system of $\mathfrak{g}^{\mathbb{C}}$ relative to \mathfrak{h} corresponding to $\tilde{\mathfrak{b}}$, and the proof of the lemma is complete.

§ 4. Root systems.

In this section we establish and record in the tables given below the information on the adapted root systems constructed by the procedure of §3 that we shall need for the proof of the main theorem. For technical reasons we divide the systems into two classes, according to whether cardinality of $Aut(\Delta)/Aut(\Delta_{\mathfrak{t}})$ is one or greater than one. The first class, treated in table 1 corresponds to the groups $SL(2n)$, $SL(2n+1)$, and $SU^*(2n)$. The second class, treated in table 2, corresponds to the groups $Sp(p, q)$, $SO(2p, 2q)$, $SO(2p, 2q+1)$, for which $|Aut(\Delta)/Aut(\Delta_{\mathfrak{t}})| = \binom{p+q}{p}$.

The tables are organized as follows. The first line in each entry identifies the group G , the type of root system, and the multiplicities of the roots. The second line gives the type of the compact part of the root system, as well as an explicit representation of the positive compact roots corresponding to a choice of positive system of roots for \mathfrak{g} . The third line lists the positive non-compact roots corresponding to the same system used to obtain the previous line. In the fourth line we exhibit a commutative set C^+ of positive noncompact roots which is, for all but very low values of the parameters, of maximum cardinality; when there is such an exception, conditions are given for C^+ to be maximal. The last line identifies the relevant real form of the group G' generated by $W + \overline{W}$, where W is the abelian space associated to the commutative system C^+ found in the previous line. Here there is an alternative (§2, Step 1b). In the first case (“type I”) G' is the group of a geodesically imbedded Hermitian space and W can be identified with its holomorphic tangent space. In the second case (“type II”), G' is not the group of a Hermitian symmetric space and in fact is equal to G itself, except in the case of $SO(2p+1, 2q+1)$ where it is equal to $SO(2p, 2q+1)$ or $SO(2p+1, 2q)$. The bounds on the cardinality of C will be derived in the following section §5.

The root systems and their orderings can be determined concretely in the following way. Choose a maximal torus T of K , and let V be the “defining” representation space for G , i.e., $V = \mathbb{R}^n$ for $SL(n, \mathbb{R})$, \mathbb{H}^n for $SU^*(2n)$, etc. The complexification of V decomposes into eigenspaces V_{λ} for the action of \mathfrak{t} . The eigenvalues, or weights, are linear functionals on \mathfrak{t} with pure imaginary values, and the set Φ of these weights is stable under complex conjugation, which acts as does multiplication by -1. Choose a vector ξ in $i\mathfrak{t}$, and let Φ^+ , Φ^- , and Φ^0 be the sets of weights which have positive, negative, and zero inner product with ξ . If ξ is chosen generically, Φ^0 is either empty or consists of zero

alone. The set Φ^+ consists of n linearly independent elements which we shall label as $\lambda_1, \dots, \lambda_n$, and $\Phi^- = -\Phi^+$.

The roots of \mathfrak{g} relative to \mathfrak{t} are the linear combinations of the basic weights just described, and have the property that \mathfrak{g}_λ maps V_μ to $V_{\lambda+\mu}$ if $\lambda + \mu$ is a weight in Φ , and to zero in the contrary case. To compute a particular root system it suffices to choose a basis for $V^{\mathbb{C}}$ which diagonalizes T . The standard basis vectors are then weight vectors, and root vectors are matrices with a small number of nonzero entries. It is therefore an elementary, if time consuming problem to determine the roots and their multiplicities. The results are recorded in the two tables already mentioned, and displayed below.

As noted already, the systems in Table 1 are characterized by the condition that $Aut(\Delta)/Aut(\Delta_{\mathfrak{k}})$ has cardinality one. In the case of $SL(2n+1)$ and $SU^*(2n)$ this is evident, since for these the Weyl groups of $G^{\mathbb{C}}$ and $K^{\mathbb{C}}$ relative to \mathfrak{t} coincide. In the case of $SL(2n)$ the two groups may differ by a factor of 2, since when n is odd, the symmetry which reverses the signs of all the λ_i is in the larger of the two Weyl groups, but not the smaller. This transformation is, however, induced by an outer automorphism of $G^{\mathbb{C}}$ preserving $K^{\mathbb{C}}$.

Table 1. Root Systems with $Aut(\Delta)/Aut(\Delta_{\mathfrak{k}}) = 1$		
Group	RS	Remarks
$SL(2n)$	C_n D_n	$m(\text{short}) = 2, m(\text{long}) = 1, \text{long: noncompact}$ $\Delta^+(\mathfrak{k}) = \{ \lambda_i \pm \lambda_j \mid i < j \}$ $\Delta^+(\mathfrak{p}) = \{ \lambda_i \pm \lambda_j \mid i < j \} \cup \{ 2\lambda_i \}$ $C^+ = \{ \lambda_i + \lambda_j \mid i \leq j \}, C \leq n(n+1)/2$ $G' = Sp(n, \mathbb{R})$
$SL(2n+1)$	BC_n B_n	$m(\text{short, medium}) = 2, m(\text{long}) = 1, \text{long: noncompact}$ $\Delta^+(\mathfrak{k}) = \{ \lambda_i \} \cup \{ \lambda_i \pm \lambda_j \mid i < j \}$ $\Delta^+(\mathfrak{p}) = \{ \lambda_i \} \cup \{ \lambda_i \pm \lambda_j \mid i < j \} \cup \{ 2\lambda_i \}$ $C^+ = \{ \lambda_i + \lambda_j \mid i \leq j \} \cup \{ \lambda_1 \}, C \leq n(n+1)/2 + 1$ $G' = SL(2n+1, \mathbb{R})$.
$SU^*(2n)$	C_n C_n	$m(\text{short}) = 2, m(\text{long}) = 1, \text{long: compact}$ $\Delta^+(\mathfrak{k}) = \{ \lambda_i \pm \lambda_j \mid i < j \} \cup \{ 2\lambda_i \}$ $\Delta^+(\mathfrak{p}) = \{ \lambda_i \pm \lambda_j \mid i < j \}$ $C^+ = \{ \lambda_i + \lambda_j \mid i < j \}, C \leq n(n-1)/2$ $G' = SO^*(2n)$

It remains, therefore, to consider the systems with $|Aut(\Delta)/Aut(\Delta_{\mathfrak{k}})| > 1$. These are distinguished by the fact that K is reducible, splitting into two factors $K_{(0)}$ and

$K_{(1)}$. Any maximal torus T splits accordingly, as does the set of weights Φ . To be precise, we say that a weight belongs to one factor of \mathfrak{t} if that factor acts nontrivially while the other one acts trivially. Viewing the indexing on the factors of K as an indexing modulo two, we obtain a parity function on weights, $\epsilon : \Phi \rightarrow \mathbb{Z}/2$. Some care must be taken when 0 is a weight, i.e., when G is $SO(2p, 2q+1)$ or $SO(2p+1, 2q+1)$. In the first case it is a weight of multiplicity 1, and its parity is that assigned to $SO(2q+1)$. In the second case it is a weight of multiplicity 2, with both even and odd weight vectors. We shall therefore view the parity function as multiple-valued. Regardless of the fine points, we there is an induced parity function on roots, which satisfies $\epsilon(\lambda + \mu) = \epsilon(\lambda) + \epsilon(\mu)$. Compact roots have parity 0 and noncompact ones have parity 1. For example, in the case of $SO(2p+1, 2q+1)$, λ_i and 0 are weights, the latter with multiplicity two and both parities. Therefore the root $\lambda_i = \lambda_i + 0$ has multiplicity two and both parities, hence has root vectors of both compact and noncompact types.

It is evident that the Weyl group of $K^{\mathbb{C}}$ acts trivially on parity functions, as does the sign-reversing element which may be an outer automorphism of $K^{\mathbb{C}}$. Therefore parity is an invariant of the equivalence class of the positive systems which we have in mind. Since all the Weyl groups contain the permutations on the index set $I = \{ 1 \dots n \}$, it is clear that the full Weyl group acts transitively on the set of parity functions. Cardinality arguments then imply that parity is a complete invariant. Therefore the systems given in table 2 are representative of all positive systems.

Table 2. Root Systems with $|Aut(\Delta)/Aut(\Delta_{\mathfrak{k}})| > 1$

Group	RS	Remarks
$Sp(p, q)$	C_{p+q} $C_p + C_q$	$m = 1$, long: compact $\Delta^+(\mathfrak{k}) = \{ \lambda_i \pm \lambda_j \mid i < j, \epsilon(\lambda_i) = \epsilon(\lambda_j) \} \cup \{ 2\lambda_i \}$ $\Delta^+(\mathfrak{p}) = \{ \lambda_i \pm \lambda_j \mid i < j, \epsilon(\lambda_i) \neq \epsilon(\lambda_j) \}$ $C^+ = \{ \lambda_i + \lambda_j \mid i \leq p < j \}, C \leq pq$ $G' = SU(p, q)$
$SO(2p, 2q)$	D_{p+q} $D_p + D_q$	$m = 1$ $\Delta^+(\mathfrak{k}) = \{ \lambda_i \pm \lambda_j \mid i < j, \epsilon(\lambda_i) = \epsilon(\lambda_j) \}$ $\Delta^+(\mathfrak{p}) = \{ \lambda_i \pm \lambda_j \mid i < j, \epsilon(\lambda_i) \neq \epsilon(\lambda_j) \}$ $C^+ = \{ \lambda_i + \lambda_j \mid i \leq p < j \}, C \leq pq$ ($p, q \geq 3$) $G' = SU(p, q)$ ($\epsilon(\lambda_1) \neq \epsilon(0)$)
$SO(2p, 2q + 1)$	B_{p+q} $D_p + B_q$	$m = 1$ $\Delta^+(\mathfrak{k}) = \{ \lambda_i \pm \lambda_j \mid i < j, \epsilon(\lambda_i) = \epsilon(\lambda_j) \} \cup \{ \lambda_i \mid \epsilon(\lambda_i) = \epsilon(0) \}$ $\Delta^+(\mathfrak{p}) = \{ \lambda_i \pm \lambda_j \mid i < j, \epsilon(\lambda_i) \neq \epsilon(\lambda_j) \} \cup \{ \lambda_i \mid \epsilon(\lambda_i) \neq \epsilon(0) \}$ $C^+ = \{ \lambda_i + \lambda_j \mid i \leq p < j \} \cup \{ \lambda_1 \}, C \leq pq + 1$ ($p, q \geq 3$) $G' = SO(2p, 2q + 1)$
$SO(2p + 1, 2q + 1)$	B_{p+q} $B_p + B_q$	$m(\text{short}) = 2, m(\text{long}) = 1$ $\Delta^+(\mathfrak{k}) = \{ \lambda_i \pm \lambda_j \mid i < j, \epsilon(\lambda_i) = \epsilon(\lambda_j) \} \cup \{ \lambda_i \}$ $\Delta^+(\mathfrak{p}) = \{ \lambda_i \pm \lambda_j \mid i < j, \epsilon(\lambda_i) \neq \epsilon(\lambda_j) \} \cup \{ \lambda_i \}$ $C^+ = \{ \lambda_i + \lambda_j \mid i \leq p < j \} \cup \{ \lambda_1 \}, C \leq pq + 1$ ($p, q \geq 3$) $G' = SO(2p, 2q + 1) \neq G$

We remark that to give a parity function is to give a decomposition of the interval $I = \{ 1 \dots n \}$ into consecutive subintervals I_j , where $j = 0, \dots, k$, where successive intervals are assigned different parities. The interval I should be thought of as an index set for the positive weights $\{ \lambda_j \}$. To take a concrete example, consider the decomposition

$$I = I_0 \cup I_1 \cup I_2 = \{ 1, 2 \} \cup \{ 3, 4 \} \cup \{ 5, 6 \},$$

then the functional $\lambda_1 - \lambda_2$ is even, hence compact, while $\lambda_2 - \lambda_3$ is odd, hence non-compact. Now consider the structure which the associated parity function defines on the positive roots $\{ \lambda_i \pm \lambda_j \mid 1 \leq i < j \leq 6 \}$. The simple roots are associated to a Dynkin diagram of type of type D_6 . Coloring the noncompact ones black, we obtain the following diagram of type D_6 that describes an adapted root system for $SO(4, 8)$:

Figure 1.

Thus, classes of positive root systems modulo $Aut(\Delta_{\mathfrak{k}})$ are also indexed by certain colored Dynkin diagrams.

§ 5. Combinatorial arguments

We shall now use the information on root systems compiled in the last section to determine the maximum cardinality ν of a commutative set of noncompact roots:

5.1. Proposition. *Let $C \subset \Delta^+(\mathfrak{p}, \mathfrak{k})$ be a commutative set of roots for one of the groups listed in table 3 below. Then its cardinality is at most $\nu(G)$, provided that $p, q \geq 6$ if $G = SO(p, q)$. The bounds are realized by the systems C^+ given in tables 1 and 2. Moreover, any other system of cardinality ν is equivalent to C^+ under the action of $Aut(\Delta_k)$, except that if $G = SO(p, q)$ we require in addition to the previous condition that $(p, q) \neq (6, 6)$. When $(p, q) = (6, 6)$ there are two equivalence classes, each represented by a system of cardinality 9.*

Table 3. The Invariant $\nu(G)$

Group	ν
$SL(2n)$	$n(n+1)/2$
$SL(2n+1)$	$n(n+1)/2 + 1$
$SU^*(2n)$	$n(n-1)/2$
$Sp(p, q)$	pq
$SO(p, q)$	$\lfloor \frac{p}{2} \rfloor \lfloor \frac{q}{2} \rfloor + \epsilon(p, q)$
$\epsilon(p, q)$ is 0 if p and q are even, is 1 otherwise	

We treat first the first three lines of the table, for which $|Aut(\Delta_{\mathfrak{g}}/Aut(\Delta_{\mathfrak{k}})| = 1$, and then treat the more intricate cases in which $|Aut(\Delta_{\mathfrak{g}}/Aut(\Delta_{\mathfrak{k}})| > 1$. In the course of our analysis we shall find that $G = SO(p, q)$, for $\min(p, q) < 6$ admits pathological systems of excess dimension, e.g., of dimension $2q + 1$ for $SO(4, 2q)$. These systems, which constitute more than one equivalence class under the action of $Aut(\Delta_{\mathfrak{k}})$, are mainly responsible for the difficulty of the final part of the proof.

The case $SL(2n)$.

According to the table, the relevant root system is of type C_n , and one system of positive noncompact roots is

$$\Delta^+(\mathfrak{p}) = \{ \lambda_i \pm \lambda_j \mid i < j \} \cup \{ 2\lambda_i \}.$$

Any other system is equivalent to it by an automorphism of G preserving \mathfrak{t} . Consider, therefore, the set of elements free of minus signs, namely,

$$C^+ = \{ \lambda_i + \lambda_j \mid i < j \} \cup \{ 2\lambda_i \}.$$

It is commutative and of cardinality $n(n+1)/2$. Any other commutative subset C must contain an element of the form $\mu = \lambda_a - \lambda_b$. The elements of Δ^+ in which λ_b appears with positive coefficient cannot be in C , since when added to μ they form a root. Therefore C lies in the set

$$C' \cup C'',$$

where C' is free of the index b and where $C'' = \{ \lambda_i - \lambda_b \mid i < b \}$ “contains” the index b with negative coefficient. Arguing by induction, we may assume that $|C'| \leq n(n-1)/2$, so that $|C| < n(n+1)/2$.

This gives the required bound, and shows that any commutative set which attains the bound is equivalent to C^+ . Note, however, that there is an automorphism of G which acts on \mathfrak{t} by the transformation which fixes all roots except $\lambda_i \pm \lambda_n$ and $\pm 2\lambda_n$, for which it interchanges signs. Application of this automorphism to C^+ yields an equivalent commutative system of the same cardinality. The equivalence is not, however, by an inner automorphism of $SL(2n, \mathbb{R})$.

Decompositions of the kind $C = C' \cup C''$ considered above will be used repeatedly in what follows. The characteristic feature is that C' is free of a certain index, while C'' “contains” it with negative coefficient. We call such a decomposition a *reduction of C relative to the given index*.

The case $SL(2n+1)$.

For BC_n we have

$$\Delta^+(\mathfrak{p}) = \{ \lambda_i \} \cup \{ \lambda_i \pm \lambda_j \mid i < j \} \cup \{ 2\lambda_i \}$$

As before all positive systems are equivalent, so that it suffices to consider the one displayed above. We argue by reduction to the previous case, writing $C = C' \cup C''$, where the first term consists of medium and long roots, the second term of short ones. It is clear that C'' has at most one element, since the sum of any two short positive roots is a root. For the other term we already have the bound $|C'| \leq n(n+1)/2$, so that $|C| \leq n(n+1)/2 + 1$.

The case $SU^*(2n)$.

The noncompact roots are $\Delta^+(\mathfrak{p}) = \{ \lambda_i \pm \lambda_j \mid i < j \}$, and the arguments given for $SL(2n)$ apply to show that $C = \{ \lambda_i + \lambda_j \mid i < j \}$ is a system of cardinality $n(n-1)/2$, that this is the maximum cardinality, and that all other systems of maximum cardinality are equivalent to it.

The case $Sp(p, q)$.

We show that $|C| \leq pq$ by induction on the quantity $\min(p, q)$. In the course of the proof we shall see that all commutative systems of maximum cardinality are equivalent.

To begin, we assume $\min(p, q) = 1$, and, without loss of generality we may take $p = 1$. Let $I = \{ 1 \dots q+1 \}$ be the index set, and let r be the index distinguished by the fact that its parity is different from all the others. Then

$$\Delta^+(\mathfrak{p}) = \{ \lambda_i \pm \lambda_r \mid i < r \} \cup \{ \lambda_r \pm \lambda_i \mid i > r \}$$

Let C_i be the subset of C for which the nondistinguished index is i . It is a subset of $\{ \lambda_i \pm \lambda_r \}$ if $i < r$, and a subset of $\{ \lambda_r \pm \lambda_i \}$ in the contrary case. These two sets

are not commutative, so C_i has at most one element. This already establishes the bound $|C| \leq q$.

Finer analysis yields the conjugacy assertion. To begin, note that if C contains $\lambda_i - \lambda_r$, then it cannot contain any element of the form $\lambda_r \pm \lambda_j$, and so $|C| < q$ if $r < q+1$. Thus, if $|C| = q$, then $C = \{ \lambda_i - \lambda_{q+1} \mid i = 1, \dots, q \}$ or $C = \{ \lambda_i + \lambda_{q+1} \mid i = 1, \dots, q \}$. These systems are equivalent under an automorphism of $G^{\mathbb{C}}$ preserving \mathfrak{t} which acts on the fundamental weights by $\lambda_i \mapsto \epsilon_i \lambda_i$, where $\epsilon_i = \pm 1$ and $\epsilon_i = 1$ for $i \leq q+1$.

Consider next the inductive step. Without loss of generality we assume that there are p indices of even parity, and that the first subinterval in the decomposition $I = I_0 \cup \dots \cup I_k$ has even parity. Thus,

$$\begin{aligned} p &= \sum_{j \text{ even}} |I_j| \\ q &= \sum_{j \text{ odd}} |I_j| \end{aligned}$$

The set of positive noncompact roots is

$$\Delta^+(\mathfrak{p}) = \{ \lambda_i \pm \lambda_j \mid i < j, \epsilon(i) \neq \epsilon(j) \}, \quad (5.2)$$

and it has an obvious commutative subset,

$$C^+ = \{ \lambda_i + \lambda_j \mid \epsilon(i) \neq \epsilon(j) \}, \quad (5.3)$$

which has cardinality pq . If C is another commutative set it must contain an element of the form $\lambda_i - \lambda_r$. Arguing as before, we shall reduce C relative to the root with negative coefficient, decomposing it as

$$C = C' \cup C'',$$

where C' is free of the index r , and where

$$C'' \subset \{ \lambda_i - \lambda_r \mid i < r, \epsilon(i) \neq \epsilon(r) \}$$

Let us analyze the case in which $\epsilon(r) = 0$. The other alternative can be handled in the same way, with the same conclusion. The first case to consider is that of $p \leq q$. Then C' is a commutative system with a smaller value of $\min(p, q)$, so, by the induction hypothesis, $C' \leq (p-1)q$. As for C'' , we observe that it is the sum of the quantities $|I_j|$, for j odd and less than b , where $r \in I_b$. Therefore $|C''| \leq q$, with equality possible only if I_b is the rightmost interval. Consequently $|C| \leq pq$. In the case of equality there is an automorphism which carries C to C^+ , namely, one which acts on the fundamental weights by $\lambda_i \rightarrow \epsilon_i \lambda_i$, where $\epsilon_i = 1$ if i is not in the rightmost interval.

For the case in which $p > q$ we use a subsidiary induction argument. Let $C(p, q)$ denote a commutative system with parameters p and q . For each reduction relative to a root $\lambda_i - \lambda_r$ there is an inequality of the form $|C(p, q)| \leq |C(p-1, q)| + q$, with strict

inequality unless r is in the rightmost interval. Apply reduction repeatedly until the resulting system has no roots of the form $\lambda_i - \lambda_j$ or until the quantity $\min(p, q)$ decreases. However this happens one obtains an inequality of the form

$$|C(p, q)| \leq |C(p - k, q)| + kq$$

or

$$|C(p, q)| \leq |C(p - k, q - 1)| + kq + (p - k)$$

In either case the right-hand side can be evaluated, giving the required bound. As before the inequality is strict unless the original system is equivalent to C^+ .

The case $SO(2p, 2q)$.

We now begin the analysis of the most difficult of all the cases, $SO(2p, 2q)$. The plan is to proceed as in the previous case, by induction on $\min(p, q)$. Because of certain unpleasant pathologies, we must consider the cases of $\min(p, q) = 1, 2,$ and 3 separately. If we assume, as we may without loss of generality, that $p \leq q$, these bounds are $2q, 2q + 1,$ and $3q,$ respectively. Given this much, a small modification of the previous induction argument shows that $|C| \leq pq$ for $\min(p, q) \geq 3$. The pathologies alluded to are the systems of excess dimension $2q + 1$ which occur for $SO(4, 2q)$. Their existence is due to the fact that $SO(2p, 2q)$ has fewer compact roots than does $Sp(p, q)$. Indeed, the long roots $2\lambda_i$ are absent, so that certain previously forbidden pairs, e.g., $\lambda_i \pm \lambda_j$, now commute.

Let us begin with the induction argument, assuming the bounds just given for low values of $\min(p, q)$ and assuming $\min(p, q) > 3$ for the system under study. The argument is identical to that used for $Sp(p, q)$, except that the system C'' used in the decomposition relative to a root $\alpha = \lambda_i - \lambda_r$ requires more care. The positive noncompact roots which commute with α and contain the index r are of the form $\beta = \lambda_j - \lambda_r$, with $j \neq i$, or $\gamma = \lambda_i + \lambda_r$. These two possibilities are mutually exclusive, since $\beta + \gamma$ is a root. If a β occurs, then C'' is as before. If a γ occurs, it is a subset of the two-element set $\{\lambda_i \pm \lambda_r\}$. However, because $\min(p, q) > 3$, the sets C'' which arise in this way are always smaller than the other kind. Consequently the argument used for $Sp(p, q)$ still applies, yielding both the bound and the conjugacy assertion. Modulo the proof of the bounds for low values of $\min(p, q)$, we have therefore established the following.

5.4. Proposition. *Let C be a commutative system of positive noncompact roots for $SO(2p, 2q)$, where $p, q > 3$. Then $|C| \leq pq$, and equality holds if and only if $|C|$ is equivalent under the action of $Aut(\Delta_{\mathfrak{f}})$ to the standard commutative system $\{\lambda_i + \lambda_b \mid \epsilon(i) \neq \epsilon(j)\}$.*

We turn next to the special cases, beginning with an analysis of the case in which the decomposition of I given by the parity function has just two subintervals.

5.5. Proposition. *Let C be a commutative subset of the system of positive noncompact roots associated to the decomposition $I = I_0 \cup I_1$, where we assume $p \leq q$. Then $|C| \leq 2q$ if $p = 1$, and $|C| \leq pq$ if $p > 1$. If $p = 1$ and equality holds then C is equivalent to the set*

$$C^* = \{ \lambda_1 \pm \lambda_j \mid j \in I_1 \},$$

and if $p > 2$ and equality holds then C is equivalent to the set

$$C^+ = \{ \lambda_i + \lambda_j \mid i \in I_0, j \in I_1 \}$$

If $p = 2$ there are two equivalence classes, represented by C^* and C^+ . These correspond to geodesic imbeddings of the symmetric spaces for $SO(2, 2q)$ and $SU(2, q)$ in the symmetric space of $SO(4, 2q)$, respectively.

The positive root systems corresponding to the partitions just described are called *Borel-Siebenthal* systems. They are characterized by having a unique noncompact simple root. For $SO(2p, 2q)$ this is the root that bridges the gap between the two intervals, namely, $\lambda_p - \lambda_{p+1}$.

Figure 2. Borel-Siebenthal system

For the proof, as well as for later work, we shall need the following technical results.

5.6. Lemma A. *Let $S = \{ \lambda_i \pm \lambda_r \mid i \in A \}$, where $r \notin A$, and let C be a commutative subset of S . Then C is a subset of one of the following commutative sets:*

- a) $\{ \lambda_i \pm \lambda_r \}$, where i is fixed, or
- b) $\{ \lambda_i + \lambda_r \mid \text{all } i \in A \}$, or
- c) $\{ \lambda_i - \lambda_r \mid \text{all } i \in A \}$.

Proof: If there exist elements $\lambda_i - \lambda_r$ and $\lambda_j + \lambda_r$ in C , then $i = j$ and C is as in case (a). Otherwise C must be as in (b) or (c).

5.7. Lemma B. *Let S and C be as above. Then $|C| \leq \max(2, |A|)$. In particular, if $|C| > 2$, then C must be a subset of one of the sets (b) or (c) above.*

The proof is immediate.

5.8. Lemma C. *Let $S = \{ \lambda_i \pm \lambda_j \mid i \in A, j \in B \}$ be a subset of a root system of type D_n , where the index sets A and B are disjoint. Then*

$$\begin{aligned} |C| &\leq 2|B| \text{ if } |A| = 1, \\ |C| &\leq |A||B| \text{ if } |A| > 1. \end{aligned}$$

Moreover, these bounds are sharp because the sets below are commutative:

- a) $\{ \lambda_1 \pm \lambda_j \mid j \in B \}$
- b) $\{ \lambda_i + \lambda_j \mid i \in A, j \in B \}$

If $|A| = 1$ then a commutative system of maximum cardinality must be of type (a). If $|A| \geq 3$, then a system of maximum cardinality must be equivalent to one of type (b). If $|A| = 2$, then there are 2 equivalence classes.

Proof: Let $C_j = C \cap \{ \lambda_i \pm \lambda_j \mid i \in A \}$ and observe that $|C| = \sum |C_j|$. The inequalities of the proposition follow from the inequalities of the previous lemma applied to the C_j . If $|A| = 1$ then it is clear that system (a) is the only one which attains the bound. If $|A| = 2$ then each C_j must be one of the four sets $\{ \lambda_1 \pm \lambda_j \}$, $\{ \lambda_2 \pm \lambda_j \}$, $\{ \lambda_1 + \lambda_j, \lambda_2 + \lambda_j \}$, $\{ \lambda_1 - \lambda_j, \lambda_2 - \lambda_j \}$. If one C_j has one of the types just listed, then commutativity forces the types of the remaining C_j 's to be the same. This gives four possible commuting sets of maximal dimension, or two equivalence classes. If $|A| > 2$ then each C_j must have cardinality $|A| > 2$, so that it is of the form $\{ \lambda_i + \epsilon_j \lambda_j \mid i \in A \}$. The automorphism which sends λ_j to $\epsilon_j \lambda_j$ and which fixes the remaining λ 's sends the given C to the one displayed in the proposition.

The proposition with which this discussion began is a special case of the last lemma.

Analysis of the case $\min(p, q) = 2$.

We turn at last to the analysis of the difficult case of $\min(p, q) = 2$. Positive root systems satisfying this condition correspond to one of the decompositions of $I = \{ 1, \dots, n \}$ below, where a subinterval written \hat{J} has cardinality one, while $\hat{\hat{J}}$ has cardinality 2.

- 1) $I_0 \hat{I}_1 I_2 \hat{I}_3 I_4$
- 2) $I_0 \hat{I}_1 I_2 \hat{I}_3$
- 3) $\hat{I}_0 I_1 \hat{I}_2 I_3$
- 4) $\hat{I}_0 I_1 \hat{I}_2$
- 5) $I_0 \hat{\hat{I}}_1 I_2$
- 6) $\hat{\hat{I}}_0 I_1$
- 7) $I_0 \hat{\hat{I}}_1$

We shall need one additional technical result:

5.9. Lemma D. *Consider disjoint sets A, A', B and subsets*

$$S = \{ \lambda_i \pm \lambda_j \mid i \in A, j \in B \}$$

$$S' = \{ \lambda_i \pm \lambda_j \mid i \in A', j \in B \}$$

in a root system of type D_n . Suppose further that $|A| = |A'|$, and let $C \subset S \cup S'$ be commutative. Then

$$|C| \leq 2|A||B|.$$

Proof: Let $\phi : A \rightarrow A'$ be a 1-1 correspondence. Define a 1-1 correspondence $\Phi : S \rightarrow S'$ by

$$\Phi(\lambda_i \pm \lambda_j) = \lambda_{\phi(i)} \mp \lambda_j$$

Then $S \cup S'$ splits into equivalence classes $\{ x, \Phi(x) \}$, where $x + \Phi(x)$ is a root. Therefore C can contain at most one element from each class, hence the result.

We must now analyze the commutative systems of positive roots associated to each of the above seven partitions. For the last two there is nothing to prove, since they are of Borel-Siebert type. We take the remaining ones in turn. The first is the most difficult to treat.

Partition 1. In this case $p = |I_0| + |I_2| + |I_4|$ and $q = 2$. Write $I_1 = \{ \alpha \}$ and $I_3 = \{ \beta \}$, and organize the positive noncompact roots as in the table below:

I'_0	I''_0	I'_2	I'_4	I''_2	I''_4
$\lambda_i + \lambda_\alpha$	$\lambda_i + \lambda_\beta$	$\lambda_\alpha + \lambda_j$	$\lambda_\alpha + \lambda_k$	$\lambda_j + \lambda_\beta$	$\lambda_\beta + \lambda_k$
$\lambda_i - \lambda_\alpha$	$\lambda_i - \lambda_\beta$	$\lambda_\alpha - \lambda_j$	$\lambda_\alpha - \lambda_k$	$\lambda_j - \lambda_\beta$	$\lambda_\beta - \lambda_k$

Indices i, j, k will be used systematically to refer to the index sets I_0, I_2 , and I_4 . Roots in the first row of the table are said to have type $I'_0(+), I''_0(+)$, while those in the second row have type $I'_0(-)$, etc. We shall use $\{ I'_r \}, \{ I''_r \}, \{ I_r \} = \{ I'_r \} + \{ I''_r \}$ to refer to the number of roots of the indicated type that lie in the system C under consideration. Applying Lemma 5.9 above, we see that only half the I_4 roots can appear, so that $\{ I_4 \} \leq 2|I_4|$. To bound $\{ I_2 \}$, partition the I_2 roots into sets with j constant:

$$C'_j = C \cap \{ \lambda_\alpha \pm \lambda_j \}$$

$$C''_j = C \cap \{ \lambda_j \pm \lambda_\beta \}$$

$$C_j = C'_j \cup C''_j$$

If $\{ I_2 \} > 2|I_2|$ then there is a j such that $|C_j| > 2$. Call this index j_0 . Commutativity then forces

$$C_{j_0} = \{ \lambda_\alpha + \lambda_{j_0}, \lambda_{j_0} + \lambda_\beta, \lambda_{j_0} - \lambda_\beta \}.$$

Moreover, the presence of $\lambda_{j_0} \pm \lambda_\beta$ in C excludes all roots of type I_0'' , and the presence of $\lambda_\alpha + \lambda_{j_0}$ excludes elements of type $I_0'(-)$. Therefore $\{ I_0 \} \leq |I_0|$. In addition, commutativity forces all sets C_j for $j \neq j_0$ to be subsets of $\{ \lambda_\alpha \pm \lambda_j \}$. Consequently $\{ I_2 \} \leq 2|I_2| + 1$, and so

$$|C| \leq |I_0| + 2|I_2| + 2|I_4| + 1.$$

Since

$$pq = 2|I_0| + 2|I_2| + 2|I_4|,$$

we have $|C| < pq$ unless $|I_0| = 1$, in which case $|C| \leq pq$. When I_0 has cardinality 1, we shall write it as $I_0 = \{ \lambda_0 \}$.

It remains, then, to analyze the case $\{ I_2 \} \leq 2|I_2|$. Note that the relation $\{ I_4 \} \leq 2|I_4|$ obtained in the previous paragraph is still in force. Now if $|I_0| > 1$, then $\{ I_0' \} \leq |I_0|$ and so by Lemma 5.8, $\{ I_0' \} \leq |I_0|$ and $\{ I_0'' \} \leq |I_0|$. Consequently $\{ I_0 \} \leq 2|I_0|$, and so $|C| \leq pq$. If $|I_0| = 1$, then $|C| \leq pq$ unless $\{ I_0 \} > 2$. The roots of type I_0 are

$$\begin{array}{ll} \lambda_0 + \lambda_\alpha & \lambda_0 + \lambda_\beta \\ \lambda_0 - \lambda_\alpha & \lambda_0 - \lambda_\beta \end{array}$$

We must study the cases in which $\{ I_0 \} \geq 3$. If both $\lambda_0 - \lambda_\alpha$ and $\lambda_0 - \lambda_\beta$ are in C , then commutativity forces $\{ I_2' \} = 0$, $\{ I_4' \} = 0$, $\{ I_4'' \} = 0$, and $\{ I_2''(+)\} = 0$. Consequently $|C| \leq 4 + |I_2| \leq pq$. There remain two possibilities: the roots in C of type I_0 are $\{ \lambda_0 + \lambda_\alpha, \lambda_0 + \lambda_\beta, \lambda_0 - \lambda_\alpha \}$ or $\{ \lambda_0 + \lambda_\alpha, \lambda_0 + \lambda_\beta, \lambda_0 - \lambda_\beta \}$. In the first case commutativity forces $\{ I_2' \} = 0$, $\{ I_4' \} = 0$, and $\{ I_2''(-)\} = 0$, so that $|C| \leq 3 + |I_2| + 2|I_4| < pq$. In the second case $\{ I_2' \} = 0$, $\{ I_4' \} = 0$. The biggest set that one can form under these restrictions is the one displayed below, which is commutative. Note that its cardinality is $pq + 1$.

Table 4. Exceptional commutative system for partion 1.

$\lambda_0 + \lambda_\alpha$	$\lambda_0 + \lambda_\beta$	$\lambda_\alpha + \lambda_j$	λ_α	+	λ_k
	$\lambda_0 - \lambda_\beta$	$\lambda_\alpha - \lambda_j$	λ_α	-	λ_k

To summarize, if C is commutative for a system of positive roots coming from partition 1, then $|C| \leq pq + 1$. Moreover, equality is possible only if $|I_0| = 1$ and C has the form given in table 4. This system is not Hermitian.

Partition 2. By ignoring the I_4 roots wherever they appear, one may apply analysis of the preceding case, obtaining the same conclusion, with the obvious modification to table 4 above.

Partition 3. In this case $p = 2$, $q = |I_1| + |I_3|$, and the roots under consideration are as below:

I'_1	I''_1	I'_3	I''_3
$\lambda_\alpha + \lambda_i$	$\lambda_i + \lambda_\beta$	$\lambda_\alpha + \lambda_j$	$\lambda_\beta + \lambda_j$
$\lambda_\alpha - \lambda_i$	$\lambda_i - \lambda_\beta$	$\lambda_\alpha - \lambda_j$	$\lambda_\beta - \lambda_j$

Apply Lemma 5.9 to get $\{ I_3 \} \leq 2|I_3|$. If $\{ I_1 \} \leq 2|I_1|$, then $|C| \leq pq$. Assume, therefore, that $|C| > pq$, hence that $\{ I_1 \} > 2|I_1|$. Let

$$C_i = \{ \lambda_\alpha \pm \lambda_i, \lambda_i \pm \lambda_\beta \} \cap C.$$

The hypothesis implies that there is an i such that $|C_i| > 2$. Without loss of generality we may assume that this $i = 2$ (note $\alpha = 1$), and that

$$C_2 = \{ \lambda_\alpha + \lambda_2, \lambda_2 + \lambda_\beta, \lambda_2 - \lambda_\beta \}.$$

In that case $\{ I''_3 \} = 0$ and

$$C_i \subset \{ \lambda_\alpha + \lambda_2, \lambda_2 - \lambda_\alpha \}$$

for $i > 2$. Therefore C is contained in the set displayed in the table below, where $j \in I_3$ is arbitrary and where $i \in I_1$ satisfies $i > 2$. This system is commutative and of cardinality $pq + 1$. As before, the bound is achieved only the system of the table, and the resulting system is non-Hermitian.

Table 5. Exceptional system.		
$\lambda_\alpha + \lambda_2$	$\lambda_2 + \lambda_\beta$	$\lambda_\alpha + \lambda_j$
	$\lambda_2 - \lambda_\beta$	$\lambda_\alpha - \lambda_j$
$\lambda_\alpha + \lambda_i$		
$\lambda_\alpha - \lambda_i$		

Partition 4. By ignoring the I_3 roots wherever they occur, the analysis of the preceding case applies to give $|C| \leq pq + 1$, with a single system of that cardinality, namely, the one obtained by suppressing the last column of the previous table.

Partition 5.

In this case $p = |I_0| + |I_2|$ and $q = 2$, and the roots to be considered are

I'_0	I''_0	I'_2	I''_2
$\lambda_i + \lambda_\alpha$	$\lambda_i + \lambda_\beta$	$\lambda_\alpha + \lambda_j$	$\lambda_\beta + \lambda_j$
$\lambda_i - \lambda_\alpha$	$\lambda_i - \lambda_\beta$	$\lambda_\alpha - \lambda_j$	$\lambda_\beta - \lambda_j$

Lemma 5.9 implies that $\{ I_2 \} \leq 2|I_2|$. If $I_0 > 1$ then Lemma 5.8 implies that $\{ I_0 \} \leq 2|I_0|$, so that $|C| \leq pq$. If $|I_0| = 1$, then the I_0 roots are $\{ \lambda_1 + \lambda_\alpha, \lambda_1 + \lambda_\beta, \lambda_1 - \lambda_\alpha, \lambda_1 - \lambda_\beta \}$. If $\{ I_0 \} \leq 2$, then we are done. If $\{ I_0 \} = 4$ then $\{ I_2 \} = 0$, since $\lambda_1 - \lambda_\alpha$ kills the $\lambda_\alpha \pm \lambda_j$ and $\lambda_1 - \lambda_\beta$ kills the $\lambda_\beta \pm \lambda_j$. In this case $|C| \leq 4 \leq pq$, with equality only in the case $|I_0| = |I_1| = 1$.

If $\{ I_0 \} = 3$ and both $\lambda_1 - \lambda_\alpha$ and $\lambda_1 - \lambda_\beta$ appear, then $|C| = 3 < pq$. If just one of the roots just discussed does not appear, then there are two possible maximal commutative systems. The first is

$$C' = \{ \lambda_1 + \lambda_\alpha, \lambda_1 - \lambda_\beta, \lambda_1 + \lambda_\beta \} \cup \{ \lambda_\alpha \pm \lambda_j \}.$$

The second is obtained from this one by replacing α with β . In this final case $|C| \leq pq + 1$, realized by the system just given, and systems conjugate to it.

This completes the analysis of the case $\min(p, q) = 2$.

The case $\min(p, q) = 3$.

Consider at last a commutative system C with $\min(p, q) = 3$. The usual arguments based on reduction along a root $\lambda_i - \lambda_r$ apply and function as expected, except that when $\min(p, q)$ decreases, C' may be a system of excess dimension. It suffices to treat this new possibility. If p is the smallest parameter, then the relevant inequality is

$$|C| \leq (p - 1)q + 1 + |C''| = pq + (1 + |C''| - q).$$

Since $p, q \geq 3$, it suffices to show that $|C''| \leq 2$. To do this, consider one of the exceptional systems E described above, and let I be its index set. Let γ be a new index, and let \tilde{E} be a commutative system containing E , built on the index set $I \cup \{ \gamma \}$, and containing a root of the form $\lambda_\gamma - \lambda_r$ or $\lambda_r - \lambda_\gamma$, where $r \in I$. We give a complete argument for the first exceptional system; the others are treated similarly. Referring to Table 4, we see that $\lambda_\gamma - \lambda_r$ cannot occur. In the second case, r must be different from any of the j 's, any of the k 's, and from β . Only the indices 0 and α remain, leaving the roots $\lambda_0 \pm \lambda_\gamma, \lambda_\alpha \pm \lambda_\gamma$ as candidates. The only possible sets C'' are therefore subsets of one of the two-element sets $\{ \lambda_0 - \lambda_\gamma, \lambda_\alpha - \lambda_\gamma \}, \{ \lambda_0 \pm \lambda_\gamma \}, \{ \lambda_\alpha \pm \lambda_\gamma \}$. Thus $|C''| \leq 2$, as required. The other cases are dealt with similarly.

The argument just given in fact proves somewhat more. If $q \geq p = 3$, and if C contains a root of the form $\lambda_i - \lambda_r$ in which a fundamental weight occurs with negative coefficient then we have shown $|C| \leq pq$. However, strict inequality holds if $q > p$. Thus, if $\min(p, q) = 3$ and C has maximum cardinality, it must be “standard”, i.e., equivalent to C^+ , unless $(p, q) = (3, 3)$. In this latter case there are, up to equivalence, just two commutative systems of maximum cardinality, namely C^+ and the system

$$C^* = \{ \lambda_0 + \lambda_1, \lambda_0 \pm \lambda_3, \lambda_0 \pm \lambda_5, \lambda_1 \pm \lambda_2, \lambda_1 \pm \lambda_4 \}$$

The parities of the indices are the natural ones, namely i modulo 2. If W is the abelian space corresponding to C^* , then $W + \bar{W}$ generates $\mathfrak{g}_{\mathbb{C}}$, i.e., $G' = G$. Thus W is not the holomorphic tangent space of any geodesically imbedded hermitian symmetric space.

We note in passing that all of the exceptional systems contain a system of the form

$$E = \{ \lambda_0 + \lambda_\alpha, \lambda_0 \pm \lambda_\beta, \lambda_\alpha \pm \lambda_j \}$$

for a suitable labeling of the indices, and where all indices are fixed except j , which ranges over a set of elements of the same parity.

The cases $SO(2p, 2q + 1)$ and $SO(2p + 1, 2q + 1)$.

In these cases the root system is of type B_n and contains the substem D_n analyzed above together with the short roots λ_i . One sees easily that the commutative system of largest cardinality is that obtained by adjoining a single short root to a system of largest cardinality for $SO(2p, 2q)$.

§ 6. Centralizers of semi-simple elements.

In this section we describe the centralizers of semi-simple elements for the algebras in our list, and use this information to find the maximum dimension μ of an arbitrary abelian space, thus completing all the dimension bounds outlined in Step 1 of §2. In the course of our analysis we will also determine an invariant ν_0 such that if $\dim W > \nu_0$ then W consists entirely of nilpotents. The values of μ and ν_0 are given in table 6 below. The case $\mu = \nu_0$ is interpreted to mean that there are two kinds of abelian subalgebras of dimension μ , one consisting entirely of nilpotents, the other containing a semisimple element.

Table 6. The Invariants μ and ν_0 .

Group	μ	ν_0	Remarks
$SU^*(2n)$	$n(n-1)/2$	$1 + (n-1)(n-2)/2$	$n \geq 3$
$SL(2n)$	$n(n+1)/2$	$1 + n(n-1)/2$	$n \geq 3$
$SL(2n+1)$	$n(n+1)/2 + 1$	μ	
$Sp(p, q)$	pq	$1 + (p-1)(q-1)$	$q \geq p \geq 2$
$SO(p, q)$	$\left[\frac{p}{2}\right] \left[\frac{q}{2}\right] + \epsilon(p, q)$ $p, q \geq 6$	$\left[\frac{p-1}{2}\right] \left[\frac{q-1}{2}\right] + 2$ $p, q \geq 7$	p and q not both odd
$SO(p, q)$	$\left[\frac{p}{2}\right] \left[\frac{q}{2}\right] + 1$	μ	p, q odd ≥ 3

To begin the proof, we recall that by part (b) of Step 1, it suffices to consider centralizers of real elements. These are described as follows:

6.1. Proposition. *Let $\mathfrak{a}_0 \subset \mathfrak{p}$ be an abelian subspace, and \mathfrak{l} be the sum of the non-compact simple subalgebras in the semi-simple part of the centralizer of \mathfrak{a}_0 . Then \mathfrak{l} is as follows:*

1. *If $\mathfrak{g} = \mathfrak{sl}(n, \mathbb{R})$, then $\mathfrak{l} = \mathfrak{sl}(n_1, \mathbb{R}) \oplus \cdots \oplus \mathfrak{sl}(n_k, \mathbb{R})$, where $n_1 + \cdots + n_k \leq n$.*
2. *If $\mathfrak{g} = \mathfrak{su}^*(2n)$, then $\mathfrak{l} = \mathfrak{su}^*(2n_1) \oplus \cdots \oplus \mathfrak{su}^*(2n_k)$, where $n_1 + \cdots + n_k \leq n$.*
3. *If $\mathfrak{g} = \mathfrak{so}(p, p+r)$, $r \geq 0$, then $\mathfrak{l} = \mathfrak{sl}(p_1, \mathbb{R}) \oplus \cdots \oplus \mathfrak{sl}(p_k, \mathbb{R}) \oplus \mathfrak{so}(q, q+r)$, where $p_1 + \cdots + p_k + q \leq p$.*
4. *If $\mathfrak{g} = \mathfrak{sp}(p, p+r)$, $r \geq 0$, then $\mathfrak{l} = \mathfrak{su}^*(2p_1) \oplus \cdots \oplus \mathfrak{su}^*(p_k) \oplus \mathfrak{sp}(q, q+r)$, where $p_1 + \cdots + p_k + q \leq p$.*

Proof: Let \mathfrak{a} be a maximal abelian subspace of \mathfrak{p} containing \mathfrak{a}_0 , and let $\Sigma \subset \mathfrak{a}^*$ denote the root system of \mathfrak{g} relative to \mathfrak{a} . Thus Σ consists of the restrictions to \mathfrak{a} of the roots of $\mathfrak{g}^{\mathbb{C}}$ relative to a Cartan subalgebra of $\mathfrak{g}^{\mathbb{C}}$ containing \mathfrak{a} . This restricted system together with the multiplicities of the simple roots determines \mathfrak{g} (see Table VI, pp 532-534 of [13]). To prove the Proposition, one first observes that from the definition of \mathfrak{l} , the root system of \mathfrak{l} is $\Sigma \cap \mathfrak{a}_0^\perp$. From this it follows that any system of positive roots for $\Sigma \cap \mathfrak{a}_0^\perp$ can be enlarged to a system of positive roots for Σ . Therefore the associated simple roots for $\Sigma \cap \mathfrak{a}_0^\perp$ are also simple roots for Σ . Thus the Dynkin diagram of \mathfrak{l} is a subdiagram of the Dynkin diagram of \mathfrak{g} . To complete the proof, one simply looks in [13], Table VI, at the Dynkin diagrams in question, and verifies that the possibilities listed in the Proposition correspond precisely all the subdiagrams, together with their multiplicities. These are the diagrams A I for case 1, A II for case 2, B II for case 3, r odd, D I for case 3, r even, and finally C II for case 4.

Consider now an abelian space $W = W_\sigma + W_\nu$, where W_σ is the complexification of a real abelian space, and $W_\nu = W_1 + \cdots + W_k$, where $W_i \subset \mathfrak{p}_i^{\mathbb{C}}$, and where $\mathfrak{l}_i = \mathfrak{k}_i \oplus \mathfrak{p}_i$ is the Cartan decompositions of the Lie algebras \mathfrak{l}_i as in the proposition.

The dimension of each of these terms is, by definition, bounded by $\nu(\mathfrak{l}_i)$, the invariant described in Proposition 5.1 and Table 3. To bound the dimension of W_σ , note that the split rank of \mathfrak{z} is the same as the split rank ρ of \mathfrak{g} , a fact which we can write as $\rho(\mathfrak{g}) = \dim(W_\sigma) + \sum_{i=1}^k \rho(\mathfrak{l}_i)$. Therefore

$$\dim W \leq \rho(\mathfrak{g}) - \sum_{i=1}^k \rho(\mathfrak{l}_i) + \sum_{i=1}^k \nu(\mathfrak{l}_i). \quad (6.2)$$

The maximum of the right-hand side over all centralizers is the invariant μ . We must compute it for each of the groups SU^* , SL , Sp , and SO . In so doing we will also determine a minimal integer ν_0 such that $\dim W > \nu_0$ implies that W consists entirely of nilpotent elements. In some cases, namely, $SL(n)$ for n odd or $SO(p, q)$ for both p and q odd, an abelian space of maximal dimension can have semisimple elements. We signal this fact by writing $\nu_0 = \mu$. The relevant information is summarized in Table 6 above.

Case SU .

We begin with this case, since it is the easiest. By the proposition, the summands \mathfrak{g}_i are of the form $\mathfrak{su}^*(2n_i)$, which have split rank $n_i - 1$. Therefore the inequality (6.2) can be written as

$$\dim W \leq k - 1 + \sum_{i=1}^k \nu(\mathfrak{l}_i).$$

Setting $f(n) = \nu(n) + 1$, this reads

$$\dim W \leq -1 + \sum_{i=1}^k f(n_i).$$

It will therefore be sufficient to maximize the expression on the right-hand side, which we write as $F(n_1, \dots, n_k)$, for all partitions of numbers $m \leq n$. In fact, it is enough to maximize $F(n_1, \dots, n_k)$ for all partitions of n . For this it is enough to observe that the function $f(x) = x(x-1)/2 + 1$ is superadditive, i.e., $f(x+y) \geq f(x) + f(y)$, for $x, y \geq 1$. Therefore F , which is a sum of superadditive functions, is also superadditive. Consequently it takes its maximum for the trivial partition (n) , and so $\mu = n(n-1)/2$. Moreover, since f is strictly superadditive if $x > 1$ or $y > 1$, the trivial partition is the only partition for which the maximum is attained, except that $F(2) = F(1, 1)$. Therefore *an abelian space of maximum dimension is necessarily nilpotent, except when $n = 2$* . In the extreme case, $SU^*(4)$, the split rank is one, and $\nu = 1$, so an abelian space of maximal dimension consists entirely of semisimple elements or entirely of nilpotents.

Suppose that W does not consist entirely of nilpotents. Then it lies in a nontrivial centralizer, hence is bounded by the value of F on a nontrivial partition. Superadditivity implies that it is in fact bounded by $F(n-1, 1)$, which is always less than $F(n)$ if $n > 2$. Set

$$\nu_0 = F(n-1, 1) = 1 + (n-1)(n-2)/2.$$

We have shown that *an abelian space of dimension $d > \nu_0$ consists entirely of nilpotent elements*.

Case SL .

The analysis is similar to that of the previous case, except that the formula for $f(n)$ depends on the parity of n . Elementary arguments show that it is superadditive, and strictly superadditive except for the cases

$$\begin{aligned} f(4) &= f(3) + f(1) \\ f(4) &= f(2) + f(2) \\ f(2m+1) &= f(2m) + f(1) \end{aligned}$$

Therefore

$$\mu(n) = \frac{1}{2} \left[\frac{n}{2} \right] \left(\left[\frac{n}{2} \right] + 1 \right) + \epsilon(n),$$

where $\epsilon(n) = 0, 1$ according to whether n is even or odd. When n is odd, there is always an abelian space of maximal dimension which has the form $\mathfrak{a}' + W'$, where \mathfrak{a}' is a one-dimensional space of semisimple elements, and where W' is an abelian space of maximum dimension in a copy of $SL(n-1)$, namely, the centralizer of \mathfrak{a}' in \mathfrak{g} . When n is even, $\mu = \nu$ except for $n = 2, 4$.

Suppose now that W is an abelian subspace for $SL(2n)$ which does not consist entirely of nilpotents. Then it lies in a nontrivial centralizer and so, by superadditivity, is bounded in dimension by $F(2n-1, 1) = n(n-1)/2 + 1$, which quantity we denote by ν_0 . Thus, *if $\dim W > \nu_0$, then W consists entirely of nilpotent elements*.

Case Sp .

We shall write the symplectic group as $Sp(p, p+r)$ as in the proposition. The decomposition of \mathfrak{z} then takes the form

$$\mathfrak{a} + \mathfrak{l}_1 + \cdots + \mathfrak{l}_k + \mathfrak{l}_{k+1}, \quad (6.3)$$

where the \mathfrak{l}_i for $i < k+1$ are copies of $\mathfrak{su}^*(p_i)$, $\mathfrak{g}_{k+1} = \mathfrak{sp}(q, q+r)$, and $p_1 + \cdots + p_k + q \leq p$. Since the split rank of $Sp(p, p+r)$ is p , the bound (6.2) can be written

$$\dim W \leq F(p_1, \dots, p_k) + q(q+r) + 1,$$

where F is the function used in arguing the case SU^* . The relation remains valid when $k = 0$, provided that one defines $F(\emptyset) = -1$. Using the bounds for SU^* , we find that

$$\begin{aligned} g(q) &= (p-q)(p-q-1)/2 + q(q+r) + 1 && \text{for } q < p, \\ g(p) &= p(p+r) \end{aligned}$$

This function is convex on the interval $[0, p-1]$ and so takes its local maxima at endpoints. In fact, it takes its local maxima at the endpoints of the larger interval $[0, p]$, from which we conclude that W is always bounded in dimension by $g(p) = p(p+r)$. This is the value of the invariant μ . Except for the case $\mathfrak{sp}(1, 1) \cong \mathfrak{so}(1, 4)$, the value of g at the right-hand endpoint exceeds the value at the left-hand endpoint. Nontrivial abelian spaces are then one-dimensional, and can be either of semisimple or nilpotent type. Thus, if $(p, p+r) \neq (1, 1)$, then an abelian subspace contained in a nontrivial centralizer must have dimension no larger than $\nu_0 = g(p-1) = (p-1)(p-1+r) + 1$. Any abelian subspace of larger dimension consists entirely of nilpotent elements.

Case SO .

The decomposition of a typical centralizer takes the form (6.3) used above, except that now $\mathfrak{l}_i = \mathfrak{sl}(p_i)$ for $i < k+1$, and $\mathfrak{l}_{k+1} = \mathfrak{so}(q, q+r)$, where $p_1 + \cdots + p_k + q \leq p$. Arguing as before, we find that

$$\dim W \leq F(p_1, \dots, p_k) + \nu(q, q+r) + 1,$$

where now F is the partition function for SL , and $\nu(q, q+r)$ is the dimension of a maximal abelian subalgebra of $\mathfrak{so}(q, q+r)$ which consists entirely of nilpotents. For low values of p there are only a small number of cases to consider. Computing them by hand, we obtain the following for $\mu(p, q)$ with $q \geq p$:

$$\begin{aligned} \mu(1, q) &= 1 \\ \mu(2, q) &= q \\ \mu(3, q) &= q + 1 \\ \mu(4, q) &= 2q + 1 \\ \mu(5, q) &= 2q + 2 \\ \mu(6, q) &= 3 \left\lfloor \frac{q}{2} \right\rfloor + \epsilon(q), \end{aligned}$$

For the remaining cases we proceed by induction, with inductive hypothesis

$$\mu(p, q) = \left\lfloor \frac{p}{2} \right\rfloor \left\lfloor \frac{q}{2} \right\rfloor + \epsilon(p, q),$$

where $\epsilon = 0$ if both p and q are even, and $\epsilon = 1$ otherwise, and where $q \geq p \geq 6$. Because of the superadditivity of the partition function which computes the dimension of the $\mathfrak{sl}(n)$ -type terms, it is enough to argue the cases 1) $\mathfrak{z} = \mathfrak{g}$, 2) $\mathfrak{z} = \mathfrak{a} + \mathfrak{g}_1$, and 3) $\mathfrak{z} = \mathfrak{a} + \mathfrak{g}_1 + \mathfrak{g}_2$, where in each \mathfrak{a} is one-dimensional. Suppose first that $(p, q) = (2a, 2b)$. In case (1) the bound is that for spaces of nilpotents, namely, ab . In case (2) the centralizer is of the form $\mathfrak{a} + \mathfrak{so}(2a - 1, 2b - 1)$, for which the bound is $(a - 1)(b - 1) + 2$. This quantity is strictly less than ab , since for the purposes of induction we assume $b \geq a \geq 4$. For case (3) The centralizer is of the form $\mathfrak{a} + \mathfrak{sl}(2c - 1) + \mathfrak{so}(2a - 2c, 2b - 2c)$ or $\mathfrak{a} + \mathfrak{sl}(2c) + \mathfrak{so}(2a - 2c, 2b - 2c)$. We compute the relevant bound in each case, and see that the second is always larger, namely, $g(c) = (a - c)(b - c) + c(c + 1)/2 + 1$. This function is convex, and so takes its local maxima at endpoints of the relevant interval, which is $[1, a]$. Now $g(1) = (a - 1)(b - 1) + 2$, which is a value already considered, and $g(a) = a(a + 1)/2 + 1$. We find that $ab > g(a)$ for $b \geq a > 2$. The induction step is therefore complete in this case (p and q both even). Moreover, we see that if W is an abelian subspace of dimension greater than

$$\nu_0 = (a - 1)(b - 1) + 2,$$

then it must consist entirely of nilpotent elements.

There is a similar analysis for the other cases. If $(p, q) = (2a, 2b + 1)$, with $a, b \geq 3$, then $\mu = ab + 1$. Moreover, if W has dimension greater than $\nu_0 = (a - 1)b + 2$, then it consists entirely of nilpotents. If $(p, q) = (2a + 1, 2b + 1)$, then $\mu = ab + 1$, but one can make no statement about nilpotents, since there is an abelian space of the form $\mathfrak{a} + W'$ with \mathfrak{a} one-dimensional and semisimple, and W' maximal abelian for a copy of $\mathfrak{so}(2a, 2b)$, the centralizer of \mathfrak{a} .

The induction argument, modulo the verifications indicated above, is now complete. We have also established the following. *Let W be an abelian subspace for $\mathfrak{so}(p, q)$, where $p, q \geq 4$, and at least one of p, q is even. If*

$$\dim W \geq \nu_0 \stackrel{\text{def}}{=} \left\lfloor \frac{p - 1}{2} \right\rfloor \left\lfloor \frac{q - 1}{2} \right\rfloor + 2,$$

then W is abelian.

§ 7. Rigidity

In this section we carry out the arguments necessary to establish infinitesimal and global rigidity that were sketched in §2 for groups of type I. The ultimate goal is to show that f takes values in the image of a quotient of a Hermitian symmetric space G_1/K_1 by a discrete group. Here is a precise statement:

7.1. Rigidity Theorem.. *Let $X = G/K$ be a symmetric space, where G is one of $SL(2n)$, $SU^*(2n)$, $SO(2p, 2q)$, or $Sp(p, q)$, subject to the restrictions $n \geq 3$, in the first two cases, $p, q \geq 3$, $(p, q) \neq (3, 3)$ in the third, and $p, q \geq 2$ in the last. Let $f : M \rightarrow X/\Gamma$ be a harmonic map of maximum rank, as given in Table 6. Then f factors as gh , where $h : M \rightarrow X_1/\Gamma_1$ is holomorphic, $X_1 = G_1/K_1$ is a hermitian symmetric space, Γ_1 is a subgroup of Γ acting discretely on it. Moreover, the map g is a geodesic immersion. Finally, the groups G_1 corresponding to G are given in Table 7 below.*

Proof: According to the sketch presented in §2, the rigidity argument comes in two parts, first order and global (=local). In the last four sections we have completed the arguments sketched in Step 1 of §2, and proved that for the groups G of Theorem (7.1) any maximum dimension abelian subalgebra consists entirely of nilpotent elements, and that its space of leading roots is unique, up to isomorphism, and is \mathfrak{p}_1^+ for the Hermitian symmetric subspace corresponding to G_1 . We conclude the remaining details of Steps 2 and 3 of §2.

Let W be an abelian subalgebra of maximum dimension μ as in Table 6. Then $W = \{X + \phi(X) : X \in \mathfrak{p}_1^+\}$ for some linear map $\phi : \mathfrak{p}_1^+ \rightarrow \mathfrak{q}^+$ satisfying (2.3). We let $\mathfrak{q}' = \mathfrak{q}^+ + \mathfrak{q}^-$, and observe that $\mathfrak{q}' \subset \mathfrak{q}^{\mathbb{C}}$, with equality for the groups SO and Sp , but strict containment for SL and SU . (In the latter cases $\mathfrak{q}^{\mathbb{C}}$ also contains the non-compact part of the fundamental Cartan subalgebra of Lemma (3.1).) We view ϕ as taking values in $\mathfrak{m} + \mathfrak{q}'$, which is a \mathfrak{p}_1^+ -module. From this perspective ϕ is a cocycle for Lie algebra cohomology, with class $[\phi] \in H^1(\mathfrak{p}_1^+, \mathfrak{m} + \mathfrak{q}')$. Observe next that \mathfrak{p}_1^+ is a \mathfrak{k}_1 -module as are the terms of the decomposition $\mathfrak{m}^+ + \mathfrak{m}^- + \mathfrak{q}'$. Therefore the space of 1-cochains

$$C^1 = \text{Hom}(\mathfrak{p}_1^+, \mathfrak{m} + \mathfrak{q}'), \quad (7.2)$$

decomposes into \mathfrak{k}_1 -modules as

$$\text{Hom}(\mathfrak{p}_1^+, \mathfrak{m}^+) \oplus \text{Hom}(\mathfrak{p}_1^+, \mathfrak{m}^-) \oplus \text{Hom}(\mathfrak{p}_1^+, \mathfrak{q}').$$

and this decomposition passes to cohomology. Indeed, one finds that

$$\begin{aligned} \delta(\mathfrak{m}^+) &= 0 \\ \delta : \mathfrak{m}^- &\longrightarrow \text{Hom}(\mathfrak{p}_1^+, \mathfrak{q}') \\ \delta : \mathfrak{q}' &\longrightarrow \text{Hom}(\mathfrak{p}_1^+, \mathfrak{m}^+), \end{aligned}$$

so that

$$H^1 = \text{Hom}(\mathfrak{p}_1^+, \mathfrak{m}^+)_0 / \delta \mathfrak{q}' \oplus \text{Hom}(\mathfrak{p}_1^+, \mathfrak{m}^-)_0 \oplus \text{Hom}(\mathfrak{p}_1^+, \mathfrak{q}')_0 / \delta \mathfrak{m}^-, \quad (7.3)$$

where the subscript denotes the space of cocycles. Using a theorem of Kostant [10], we shall show, under the hypotheses of Theorem 7.1, that only the first term in the preceding decomposition is nonzero: the \mathfrak{m}^- and \mathfrak{q}' components of H^1 vanish. Thus ϕ is a coboundary, so that there is an element X_0 of $\mathfrak{m}^{\mathbb{C}}$ such that $\phi(X) = [X, X_0]$. As noted in §2, this implies that W' is $K^{\mathbb{C}}$ -conjugate to \mathfrak{p}_1^+ .

To obtain the stronger K -conjugacy result, we compare the K and $K^{\mathbb{C}}$ orbits of \mathfrak{p}_1^+ . The first can be identified with K/K_1 and the second with $K^{\mathbb{C}}/F$, where F is a parabolic subalgebra with Lie algebra $\mathfrak{f} = \mathfrak{k}_1^{\mathbb{C}} + \mathfrak{m}^+$. From this last assertion it follows that the two orbits have the same dimension. Indeed, the real tangent space of K/K_1 is the set of real points in \mathfrak{m} , whereas that of $K^{\mathbb{C}}/F$ is \mathfrak{m}^- , viewed as a real vector space. It follows that the inclusion of the K -orbit in the $K^{\mathbb{C}}$ -orbit is open. But the K orbit is compact, so that in view of the connectivity of the $K^{\mathbb{C}}$ -orbit, the inclusion must be an isomorphism.

As explained already Step 3 of §2, from first-order rigidity one obtains local liftings of a harmonic map f of maximum rank from G/K to G/K_1 . Local rigidity follows once this map is known to be horizontal, meaning that $\mu = 0$, where $d = D_1 + \mu + \theta$ gives the reduction of the flat G -connection to a K_1 -connection. Since μ is real-valued, it suffices to establish the vanishing of its (1,0) component μ' . This we do using the equations which μ' satisfies. First, since df maps $T^{1,0}M$ onto \mathfrak{p}_1^+ , we may regard μ' as defined on \mathfrak{p}_1^+ . Equation (2.9) implies that $\mu' : \mathfrak{p}_1^+ \rightarrow \mathfrak{m}$, viewed as is a map from \mathfrak{p}_1^+ to $\mathfrak{m} + \mathfrak{q}'$, defines a class in $H^1(\mathfrak{p}_1^+, \mathfrak{m} + \mathfrak{q}')$. By the application of Kostant's theorem, already referred to above, we see that $[\mu'] = [\mu'_+] + [\mu'_-]$, with $[\mu'_-] = 0$. In fact, μ'_- vanishes as a cocycle, since, by (2.3), there are no coboundaries of type \mathfrak{m}^- . To kill the remaining part of μ , we use the conjugate of the second equation of (2.8), which implies that $\mu'_+(X)$ is a vector in \mathfrak{m}^+ commuting with \mathfrak{p}_1^- . From Table 7 below, we see that any such vector must vanish.

It remains to justify the assertion made earlier, that all $H^1(\mathfrak{p}_1^+, \mathfrak{m} + \mathfrak{q}')$ is entirely of type \mathfrak{m}^+ . For this we use a special case of Theorem 5.14 of [10], which may be stated as follows:

7.4. Theorem (Kostant). *Let $(\mathfrak{g}_1, \mathfrak{k}_1)$ be a Hermitian symmetric pair, and let V be an irreducible \mathfrak{g}_1 -module with highest weight λ and highest weight vector v_λ . Consider a system of simple roots for \mathfrak{g}_1 such that \mathfrak{p}_1^+ is a sum of positive root spaces. Let σ be a simple reflection such that all roots of $\Delta^+(\mathfrak{k}_1)$ are positive. Let ω be an element of the space dual to \mathfrak{p}_1^+ . Then highest weight vectors of $H^1(\mathfrak{p}_1^+, V)$ have the form $\omega \otimes v_{\sigma(\lambda)}$.*

Using the criterion of the theorem and the explicit form of the roots for \mathfrak{g}_1 and \mathfrak{k}_1 given in Table 7, it is easy to determine the (unique) distinguished reflection σ of the preceding theorem. Under the hypotheses of Theorem 7.1 the highest weight vectors of $\mathfrak{m} + \mathfrak{q}'$, which are always of type \mathfrak{m}^+ , are invariant under σ . This completes the proof.

Table 7.

Group	Roots	Remarks
$SL(2n)$	$\Delta^+(\mathfrak{k}_1) = \{ \lambda_r - \lambda_s \mid r < s \}$ $\Delta^+(\mathfrak{p}_1) = \{ \lambda_r + \lambda_s \mid r \leq s \}$ $\Delta^+(\mathfrak{m}) = \{ \lambda_r + \lambda_s \mid r < s \}$ $\Delta^+(\mathfrak{q}) = \{ \lambda_r - \lambda_s \mid r < s \}$ $\mathfrak{m} + \mathfrak{q}' = V^{\lambda_1 + \lambda_2}$ $\sigma = \sigma_{\alpha_n}$	$\mathfrak{k}_1 = \mathfrak{u}(n)$ $\mathfrak{g}_1 = \mathfrak{sp}(n, \mathbb{R})$ $\mathfrak{k} = \mathfrak{so}(2n)$ $\lambda_n \longrightarrow -\lambda_n$
$SU^*(2n)$	$\Delta^+(\mathfrak{k}_1) = \{ \lambda_r - \lambda_s \mid r < s \}$ $\Delta^+(\mathfrak{p}_1) = \{ \lambda_r + \lambda_s \mid r < s \}$ $\Delta^+(\mathfrak{m}) = \{ \lambda_r + \lambda_s \mid r \leq s \}$ $\Delta^+(\mathfrak{q}) = \{ \lambda_r - \lambda_s \mid r < s \}$ $\mathfrak{m} + \mathfrak{q}' = V^{2\lambda_1}$ $\sigma = \sigma_{\alpha_n}$	$\mathfrak{k}_1 = \mathfrak{u}(n)$ $\mathfrak{g}_1 = \mathfrak{so}^*(2n)$ $\mathfrak{k} = \mathfrak{sp}(n)$ $\lambda_n \mapsto -\lambda_n, \lambda_{n-1} \mapsto -\lambda_{n-1}$
$SO(2p, 2q)$	$\Delta^+(\mathfrak{k}'_1) = \{ \lambda_a - \lambda_b \mid a < b \}$ $\Delta^+(\mathfrak{k}''_1) = \{ \lambda_i - \lambda_j \mid i < j \}$ $\Delta^+(\mathfrak{p}_1) = \{ \lambda_a - \lambda_i \}$ $\Delta^+(\mathfrak{m}') = \{ \lambda_a + \lambda_b \mid a < b \}$ $\Delta^+(\mathfrak{m}'') = \{ -\lambda_i - \lambda_j \mid i < j \}$ $\Delta^+(\mathfrak{q}) = \{ \lambda_a + \lambda_i \}$ $\mathfrak{m} + \mathfrak{q}' = V^{\lambda_1 + \lambda_2} \oplus V^{-(\lambda_{n-1} + \lambda_n)}$ $\sigma = \sigma_{\alpha_p} = (p, p+1)$	$\mathfrak{k}'_1 = \mathfrak{u}(p)$ in general, $a, b \leq p$ $\mathfrak{k}''_1 = \mathfrak{u}(q)$ in general, $i, j > p$ $\mathfrak{g}_1 = \mathfrak{u}(p, q)$ $\mathfrak{k}' = \mathfrak{so}(2p)$ $\mathfrak{k}'' = \mathfrak{so}(2q)$ $n = p + q$ $\lambda_p \leftrightarrow \lambda_{p+1}$
$Sp(p, q)$	$\Delta^+(\mathfrak{k}'_1) = \{ \lambda_a - \lambda_b \mid a < b \}$ $\Delta^+(\mathfrak{k}''_1) = \{ \lambda_i - \lambda_j \mid i < j \}$ $\Delta^+(\mathfrak{p}_1) = \{ \lambda_a - \lambda_i \}$ $\Delta^+(\mathfrak{m}') = \{ \lambda_a + \lambda_b \mid a \leq b \}$ $\Delta^+(\mathfrak{m}'') = \{ -\lambda_i - \lambda_j \mid i \leq j \}$ $\Delta^+(\mathfrak{q}) = \{ \lambda_a + \lambda_i \}$ $\mathfrak{m} + \mathfrak{q}' = V^{2\lambda_1} \oplus V^{-2\lambda_n}$ $\sigma = \sigma_{\alpha_p} = (p, p+1)$	$\mathfrak{k}'_1 = \mathfrak{u}(p)$ in general, $a, b \leq p$ $\mathfrak{k}''_1 = \mathfrak{u}(q)$ in general, $i, j > p$ $\mathfrak{g}_1 = \mathfrak{u}(p, q)$ $\mathfrak{k}' = \mathfrak{so}(2p)$ $\mathfrak{k}'' = \mathfrak{so}(2q)$ $n = p + q$ $\lambda_p \leftrightarrow \lambda_{p+1}$

§ 8. Existence.

The purpose of this section is to demonstrate the existence of non-obvious harmonic mappings, namely, ones that do not come from sources 1–3 mentioned in the introduction (geodesic immersions, maps which factor through a curve, and maps which factor through totally geodesic real tori). Our constructions, based on the following lemma, come from Hodge theory:

8.1. Lemma.. *Let $F : M \rightarrow D/\Gamma$ be a variation of Hodge structure, and let $p : D/\Gamma \rightarrow X/\Gamma$ be the canonical projection to the associated locally symmetric space. Then $p \circ F$ is pluriharmonic.*

With this in hand, we can construct pluriharmonic maps whose domain is, in an essential way an algebraic surface:

8.2. Theorem.. *There exist pluriharmonic mappings $f : M \rightarrow X/\Gamma$, where M is an algebraic surface and $X = G/K$ with $G = SO(2p, q)$. The differential of these mappings is generically injective, and the image of the fundamental group is a lattice Γ in G .*

Proof of (8.1). Recall that $D = G/V$ and $X = G/K$, where K is maximal compact and $V \subset K$ is the centralizer of a torus. A map from $f : M \rightarrow D/\Gamma$ is given by a flat G -connection ∇ and a reduction of it to a V -connection, where $D = G/V$. Write $\nabla = D + \theta$, where D is the V -connection and θ is a \mathfrak{g} -valued 1-form representing df . Write the decomposition into forms of type $(1, 0)$ and $(0, 1)$ as $\theta = \theta' + \theta''$. If θ' is $\mathfrak{g}^{-1,1}$ -valued, where $\mathfrak{g}_{\mathbb{C}} = \sum \mathfrak{g}^{p,-p}$ is the natural Hodge decomposition on the Lie algebra, then f is, by definition, a variation of Hodge structure. Now write out the equation $\nabla^2 = 0$, separating it into components relative to the decomposition

$$\mathfrak{g} \otimes A^k = \sum \mathfrak{g}^{-p,p} \otimes A^{r,s}.$$

One finds that $D''^2 = 0$, $D''(\theta') = 0$, and $[\theta', \theta'] = 0$. These same equations hold if one views D as a K -connection and θ as a \mathfrak{p} -valued one-form, i.e., if one views the projection f of F to the associated locally symmetric space. But these are the pluriharmonic equations for $p \circ F$.

It remains to construct variations of Hodge structure with compact base. For this it would suffice to construct a smooth family of algebraic varieties with compact parameter space M ; the “period mapping”, which associates to $t \in M$ the Hodge structure on $H^k(V_t)$ then gives a variation of Hodge structure. Unfortunately, it is difficult to construct nontrivial families with compact parameter space. We therefore construct first a smooth family $V \rightarrow U$, where U is a Zariski open in a projective manifold T , and with the property that the local monodromy representation is finite. To explain this, let $D = T - U$ be the divisor at infinity, let p be a point of D , let B be a small ball centered at p , and let b be a point of $B - D$. “The” local monodromy representation is the natural representation $\rho_p : \pi_1(B - D, b) \rightarrow GL(H^k(V_b))$. By pulling back the original family on U to a suitable unramified cover U' , we obtain a family with trivial local monodromy. The new parameter space sits as a Zariski open in a projective variety M ; appealing to a theorem of Griffiths that asserts that period mappings behave locally as do holomorphic maps to a bounded domain, we find that the lifted map extends (as a variation of Hodge structure) to M . We have therefore reduced the construction to a) filling in the details of the sketch just given and b) constructing nontrivial families with finite local monodromy.

Let us complete the sketch of the proof, then construct the required families. To this end we suppose given a family of algebraic varieties $V \rightarrow T$ which is smooth over a Zariski open set $U = T - D$, where D is a normal crossing divisor, i.e., one whose local equations are of the form $z_1 \cdots z_k = 0$, where z_1, \dots, z_n are local analytic coordinates. Let $\rho : \pi_1(U, b) \rightarrow \Gamma$ be the monodromy representation. Its image is a finitely generated matrix group and so has a normal subgroup Γ' which is torsion-free and of finite index in Γ . We may take this subgroup to be defined by suitable congruences [2]. Let U' be the unramified cover corresponding to the kernel of the natural map $\pi_1(U, b) \rightarrow \Gamma/\Gamma'$, and let $\rho' : \pi_1(U', b') \rightarrow \Gamma'$ be the induced monodromy representation. According to lemma 8.3 below, the cover U' has the form $M - D'$, for a suitable projective manifold M . Now suppose that the local monodromy representations for the family V over T are finite. The same is true for the pullback of this family to M . However, these representations must in fact be trivial, since they take values in a torsion-free group. Griffiths' extension theorem [12] then applies to show that the period map for this family extends from U' to M .

We turn now to the compactification result:

8.3. Lemma. *Let Y be a projective variety, D a hypersurface in Y , and $f' : U' \rightarrow Y - D$ a finite unramified cover. Then there is a projective variety X , and an algebraic map $f : X \rightarrow Y$ which extends f' .*

Proof. By resolution of singularities we may assume that D has normal crossings. Thus, a p is a point of D has a coordinate neighborhood V such that $V^* = V \cap Y - D$ is isomorphic to a product of disks and punctured disks: $V^* = \Delta^p \times \Delta^{*q}$. The unramified covers of these are given by maps

$$(s_1, \dots, s_p, t_1, \dots, t_q) \rightarrow (s_1, \dots, s_p, t_1^{n_1}, \dots, t_q^{n_q})$$

Therefore one can locally form a smooth compactification of the cover in question. The local divisor at infinity is a normal crossing variety. These local compactifications patch together to give a global compactification which, *a priori*, is merely an analytic space. Using the Nakai-Moishezon criterion, however, one sees that it is in fact projective. We give the complete argument in the two-dimensional case, which is all we need. Let L be a bundle on Y which projectively imbeds Y , and let \tilde{L} be its pullback to X . Take two sections of L whose zero sets Z_1 and Z_2 meet transversely have no common points which lie on the branch locus. Then the intersection number $(Z_1 \cdot Z_2)$ is positive. If \tilde{Z}_i denotes the zero set of the pullback, then $(\tilde{Z}_1 \cdot \tilde{Z}_2) = n(Z_1 \cdot Z_2) > 0$. Here n is the degree of the cover. Next, let W be an irreducible curve on X which does not lie in the ramification locus. Its projection to Y meets a suitably chosen zero set Z of a section of L transversely and away from the branch locus. Since W meets the inverse image of Z in the same way, $(W \cdot \tilde{Z})$ is positive. It remains to consider the case of W irreducible and contained in the branch locus. Since that locus is a normal crossing variety, we can still find a section Z of L whose zero set meets the projection of W transversely. Again, we conclude that $(W \cdot \tilde{Z})$ is positive.

Finally, we construct two-parameter families whose global monodromy group is arithmetic, but whose local monodromy groups are finite. To this end consider the family of all hypersurfaces of degree d and dimension n , parametrized by \mathbb{P}^N . Let D be the discriminant locus, i.e., the set $t \in \mathbb{P}^N$ such that the corresponding hypersurface X_t is singular. Let $\tilde{\Gamma} = O(H^n(X_t)_0)$ be the orthogonal group on the primitive integral cohomology which preserves the cup-product. The monodromy group Γ of this family is a subgroup of $\tilde{\Gamma}$ and, according to a result of Beauville [1] Theorem 2, it is of index two, provided that n is even and $d \geq 4$. If $n \geq 4$, it is enough to require that $d \geq 3$. Consider now a generic 2-plane $\mathbb{L} \subset \mathbb{P}^N$, and let $C = \mathbb{L} \cap D$ be its intersection with the discriminant locus. This curve contains a Zariski-open set of smooth points corresponding to hypersurfaces with a single ordinary double point and no other singularities. In suitable local analytic coordinates x_1, \dots, x_{n+1} the equation of such a hypersurface near such a point is $x_1^2 + \dots + x_{n+1}^2 = 0$. It is well-known that the local monodromy group for these singularities is of order two, given by a Picard-Lefschetz reflection. It is also fairly well-known that for when L is generic, the singular locus of C consists of two types of points, those for which X_t has two ordinary double points, and those for which X_t has one A_2 singularity and is otherwise smooth. Recall that an A_2 singularity has local equation $x_1^2 + \dots + x_n^2 + x_{n+1}^3 = 0$. In either case, the local monodromy group is finite. Moreover, the inclusion of $L - C$ in $\mathbb{P}^N - D$ induces an isomorphism on fundamental groups, so that our family restricted to L satisfies both of the local and global conditions we have sought to impose.

We were unable to find a reference for the fact that the singular locus of C , for generic L , has the asserted properties, and so reproduce here an argument due to Robert Varley. To begin, we note that the singular locus of D is the union of two pieces D' and D'' , the first consisting of points t for which X_t has more than one singular point, the second consisting of points t for which X_t has a singular point whose tangent cone is nongeneric. To understand D' , consider the locus $L(p, q)$ of points t in D for which X_t has two singularities at fixed locations p and q in \mathbb{P}^{n+1} . This is a linear subspace of \mathbb{P}^N , since it is defined by the equations $\nabla F(p) = 0$, $\nabla F(q) = 0$, where F is a homogeneous form with undetermined coefficients, and where ∇F is its gradient. The linear system of hypersurfaces parametrized by $L(p, q)$ has a base locus B , namely the set of points common to all of the hypersurfaces of the system. By Bertini's theorem, a general member of this system of hypersurfaces is smooth outside B . However, if d is large enough, e.g., larger than 3, B consists of p and q alone. (One may argue with reducible hypersurfaces to establish this point). Similar arguments apply to D'' . Consider the locus $L(p, v)$ consisting of hypersurfaces $\{ F = 0 \}$ where $\nabla F(p) = 0$ and $H(F, p)v = 0$, where $H(F, p)$ is the (Hessian) matrix of second partials at p . This set is again a projective linear subspace, and Bertini's theorem applies in the same way: if d is large enough, then the base locus consist of p and nothing else.

It remains to show that the differential of the pluriharmonic map f is generically injective. By Griffiths' local Torelli theorem [11], dF is generically injective if the 2-plane L is generically transverse to the orbits of $PGL(n)$ acting on \mathbb{P}^N , which is the case if L is chosen generically. Since F is horizontal, the result follows.

§ 9. Concluding Remarks

In this section we prove some of the results that were mentioned in the introduction and which follow from our methods. We also discuss some open problems.

First we recall the statement of theorem 1.2 from the introduction, concerning representations into $SO^+(2p, 2q)$:

9.1. Theorem. *Let M be a compact Kähler manifold, let $\rho : \pi_1(M) \rightarrow SO^+(2p, 2q)$, $p, q \geq 3$, $(p, q) \neq (3, 3)$, be a representation, and let $e_+ \in H^{2p}(M)$ be the Euler class of a maximal positive sub-bundle of the flat vector bundle associated to ρ . If $e_+^q \neq 0$, then the image of ρ is contained in some embedding of $U(p, q)$ in $SO^+(2p, 2q)$.*

For the proof, we note that if the representation ρ is reductive, then by a Theorem of Corlette [6], it follows that the associated flat bundle has a harmonic section, equivalently, that there is a ρ -equivariant harmonic map $f : \tilde{M} \rightarrow X$, where X is the symmetric space for $SO^+(2p, 2q)$. The characteristic class in question is represented by the pull-back by f of an invariant form on X of degree $2pq$, hence its non-vanishing implies that f has rank at least $2pq = \dim(X_1)$, where X_1 is the symmetric space for $U(p, q)$. The compactness of M implies that f is pluriharmonic, hence by the analogue of Theorem 1.1 for sections, it follows that the image of f is contained in an embedded copy of X_1 . This implies that the image of ρ is contained in the subgroup of $SO^+(2p, 2q)$ which leaves X_1 invariant, and this group is isomorphic to $U(p, q)$. Thus it suffices to prove that ρ is reductive, and this follows from the following Lemma:

9.2. Lemma. *Let $\rho : M \rightarrow SO^+(2p, 2q)$ be a representation such that the characteristic class e_+ satisfies $e_+^2 \neq 0$. Then the Zariski closure of the image of ρ is a reductive subgroup of $SO^+(2p, 2q)$.*

Proof: Otherwise, the Zariski image of the image of ρ is contained in a proper \mathbb{R} -parabolic subgroup of $SO^+(2p, 2q)$, and ρ can be deformed into a representation into the (semi-simple) Levi factor H of this subgroup. Now the possibilities for H can be read off from the third line of Proposition 6.1. Geometrically this means that the indefinite inner product space $\mathbb{R}^{2p, 2q}$ splits as an orthogonal direct sum

$$\mathbb{R}^{2p, 2q} = \mathbb{R}^{a_0, a_0} \oplus \mathbb{R}^{a_1, a_1} \oplus \dots \oplus \mathbb{R}^{a_k, a_k} \oplus \mathbb{R}^{b, b+c},$$

where $a_0 + a_1 + \dots + a_k + b = 2p$, $c = 2q - 2p$, H acts trivially on \mathbb{R}^{a_0, a_0} , acts by the standard embedding of $SL(a_i, \mathbb{R})$ in $SO^+(a_i, a_i)$ on \mathbb{R}^{a_i, a_i} , and acts as $SO^+(b, b+c)$ on $\mathbb{R}^{b, b+c}$. The bundle E associated to ρ splits accordingly, and maximal positive and negative sub-bundles E_+ , E_- of E can be chosen to also split accordingly:

$$E_+ = E_+^0 \oplus E_+^1 \oplus \dots \oplus E_+^k \oplus E'_+,$$

similarly for E_- . Now the Euler class e is multiplicative, so

$$e_+ = e(E_+) = e(E_+^0)e(E_+^1) \dots e(E_+^k)e(E'_+).$$

For H to be a proper subgroup of $SO^+(2p, 2q)$ we must have that for at least one $i \in \{0, \dots, k\}$ the summand $E_+^i \neq 0$. If the summand $E_+^0 \neq 0$, since it corresponds to a trivial sub-bundle, $e(E_+^0) = 0$, and consequently $e_+ = 0$. If one of the summands $E_+^i \neq 0$, since the bundle $E_+^i \oplus E_-^i$ is flat and $E_+^i \approx E_-^i$ as real vector bundles (because the embedding of $SL(a_i)$ in $SO^+(a_i, a_i)$ preserves a symplectic form on \mathbb{R}^{a_i, a_i} which dually pairs E_+^i and E_-^i), we see that $e(E_+^i)^2 = 0$ and consequently $e_+^2 = 0$, and the proof is complete.

From the discussion of §8 we note that harmonic maps which arise from variations of Hodge structure are nilpotent, meaning that $df(T_x^{1,0}M) \subset \mathfrak{p}^{\mathbb{C}}$ is nilpotent for all $x \in M$. As noted in the introduction there is a partial converse to this. The number $\nu_0 = \nu_0(G/K)$, is as in §6, Table 6, and $2\nu_0(X) = r(X)$, where $r(X)$ is as in the introduction.

9.3. Theorem. *Let $f : M \rightarrow N$ be a harmonic map. If $\text{rank}(d_x f) > 2\nu_0$ for some $x \in M$, then f lifts to a variation of Hodge structure.*

Proof: As in the proof of Theorem 7.1, we can view f as defined by a flat connection ∇ and a decomposition $D + \theta$, where D is a K connection and the \mathfrak{p} -valued one-form represents df . Let $D = D' + D''$ and $\theta = \theta' + \theta''$ be the decompositions into type. The pluriharmonic equation $D''\theta' = 0$ and its consequences $D''^2 = 0$ and $[\theta', \theta'] = 0$ assert that the pair (D'', θ') constitute a Higgs bundle. (These notions were introduced by Hitchin [14] and extended by Simpson [21]. Here we shall follow the excellent exposition of [8]). Now there is a moduli space of Higgs bundles and a circle action on it, the fixed points of which are variations of Hodge structure:

$$D + \theta' + \theta'' \mapsto D + e^{it}\theta' + e^{-it}\theta''$$

The infinitesimal generator of this action is, relative to the natural symplectic form defined on the space of Higgs bundles, the gradient of the energy ([14], second equation of §7). The latter is defined as $E(\theta) = i\eta(\theta', \theta'')$, where

$$\eta(\phi, \psi) = \int_M B(\phi, \psi) \wedge \omega^{m-1}$$

is the symplectic form. Here B is the Killing form and ω is the Kähler form. Now, by a remark of Corlette, the energy is a plurisubharmonic function on the space of Higgs bundles. The relevant computation is

$$\frac{\partial^2 E}{\partial t \partial \bar{t}} i\eta(\theta' + t\phi', \theta'' + t\phi'') = i\eta(\phi', \phi'')$$

where

$$\nabla_t = \nabla + t\zeta = (D + t\psi) + (\theta + t\phi)$$

is a family of Higgs bundles. Since ϕ is real, $\phi'' = \bar{\phi}'$, and so the Levi form is positive on the space of connections. It is also basic, and so descends to the moduli space of Higgs bundles.

It remains to show that the component of the moduli space in which the Higgs bundle associated to f lies is compact. For then the energy, which is plurisubharmonic, is constant, and so its gradient, which generates the circle action, is zero, hence the circle action is trivial and f is necessarily a fixed point. But our hypothesis on the rank (at some point $x \in M$, hence on the complement of an analytic subvariety) implies that all deformations θ'_t of the given Higgs field are nilpotent, i.e., $\theta'_t(X_p)$ is a nilpotent endomorphism for any holomorphic tangent vector X_p . But, according to Lemma 3.17 of [21] (cf. also §4 of [14], the map which associates to a Higgs field its characteristic polynomial is proper, and so the result follows.

Finally we make some remarks on problems that are more global in nature.

The first concerns the algebras of Type II, where the largest-dimensional abelian subalgebra (consisting of nilpotents) does not correspond to a Hermitian symmetric subspace. The space of leading roots is a maximum commutative system C which contains a subsystem C' of cardinality one less which is the $(1,0)$ -tangent space of a Hermitian symmetric subspace. We first observe that for the groups $SO(2p, 2q + 1)$ these algebras can be realized as tangent spaces of pluriharmonic maps. This follows from the example of a variation of Hodge structure constructed in §7, IV, of [4], which, by projection to the symmetric space produces, by Lemma 8.1, a pluriharmonic map. Comparing the description of C^+ in Table 2 with the correspondence between tangent vectors and root vectors displayed in p. 685 of [4], we see that this pluriharmonic map realizes the abelian space W_C corresponding to the maximum commutative system $C = C^+$. The domain of this map is a Siegel domain of the third kind which fibres over the unit disk in \mathbb{C} , with fibre the Hermitian symmetric space corresponding to the subsystem C' .

It seems unlikely that these domains admit compact, or even quasiprojective quotients. It also seems possible that these examples are the only ones in the sense that any other factors through one of the “standard” ones on the level of universal covers. If both these statements turn out to be true, they would imply that these algebras cannot be realized by a pluriharmonic map of a *compact* complex manifold to a quotient of G/K . It would then follow that the maximum homotopical rank of a map of a compact Kähler manifold to a quotient of G/K would still be the largest dimension of a Hermitian symmetric subspace.

It also seems possible that the maximum-dimensional abelian subalgebra of the remaining groups of Type II (and of the exceptional 9-dimensional commutative system for $SO(6, 6)$, which makes this group, in our terminology, both of Type I and Type II) can also be realized by similar pluriharmonic maps of Siegel domains. If this is the case, then we could also attempt the more global approach suggested above in all these cases.

Finally we would like to point out one interesting problem concerning the group $Sp(n, 1)$. We proved in Theorem (3.5) (c) of [5], and it also follows from our results here (in §7), that first-order rigidity holds in this case. But local rigidity clearly cannot hold, since there are as many local pluriharmonic mappings as there are horizontal mappings into the associated twistor space $Sp(n, 1)/Sp(n) \times U(1)$, and the latter are integral manifolds of a holomorphic contact structure, hence form a space of infinite dimension. The

question of global rigidity (for maps of *compact* Kähler manifolds into quotients of the symmetric space) is open and we have no conjectures on its possible resolution.

Bibliography

- [1] Beauville, A., Le groupe de monodromie d'hypersurfaces et d'intersections completes. Lect. Notes. Math. 1194, 1–18 (1986).
- [2] Borel, A., Compact Clifford-Klein forms of symmetric spaces, *Topology* 2 (1963) 111-122.
- [3] Carlson, J.A., and L. Hernández, Harmonic maps from compact Kähler manifolds to exceptional hyperbolic spaces, *Jour. Geom. Analysis* 1 (1991), 339 - 357.
- [4] Carlson, J., A. Kasparian, and D. Toledo, Variations of Hodge structure of maximal dimension, *Duke Math. J.* 58 (1989), 669–694.
- [5] Carlson, J. and D. Toledo, Harmonic mappings of Kähler manifolds to locally symmetric spaces, *Pub. Math. IHES* 69 (1989), 173–201.
- [6] Corlette, K., Flat G-bundles with canonical metrics, *Jour. Diff. Geom.* 28 (1988), 361-382.
- [7] Corlette, K., Rigid representations of Kahlerian fundamental groups, *Jour. Diff. Geom.* 33 (1991), 239 - 252.
- [8] Corlette, K., Nonabelian Hodge theory, Preprint, Univ. of Chicago, 1990.
- [9] Eells, J., and J.H. Sampson, Harmonic mappings of Riemannian manifolds, *Amer. J. Math.* 86 (1964), 109–160.
- [10] Kostant, B., Lie algebra cohomology and the generalized Borel-Weil Theorem, *Ann. of Math.* 74 (1961) 329-387.
- [11] Griffiths, P.A., Periods of rational integrals, I, II, *Ann. of Math.* 90 (1969), 469–526 and 805–865.
- [12] Griffiths, P.A., Periods of integrals of algebraic manifolds, III, *Pub. Math. I.H.E.S.* 38 (1970), 125–180.
- [13] Helgason, S., *Differential Geometry, Lie Groups, and Symmetric Spaces*, Academic Press, 1978.
- [14] Hitchin, N. J., Self-duality equations on a Riemann surface, *Proc. London Math. Soc.* (3) 55 (1987), 59-126.
- [15] Malcev, A.I., Commutative subalgebras of semisimple Lie algebras, *Izvestia Ak. Nauk USSR (Russian)* 9 (1945), 125–133; English: *Amer. Math. Soc. Translations* No. 40 (1951).
- [16] Margulis, G. A., Discrete groups of motions of manifolds of non-positive curvature (in Russian), *Proc. Int. Cong. Math., Vancouver, 1974*, vol. 2, 35 -44; *AMS Transl.* (2), Vol. 109 (1977), 33 - 45.

- [17] Matsuki, T., The orbits of affine symmetric spaces under the action of minimal parabolic subgroups, *J. Math. Soc. Japan* 31 (1979), 331–357.
- [18] Mostow, G. D., *Strong Rigidity of Locally Symmetric Spaces*, *Annals of Math. Studies* 78, Princeton Univ. Press, 1973.
- [19] Sampson, J.H., Applications of harmonic maps to Kähler geometry, *Contemp. Math.* 49 (1986), 125–133.
- [20] Schur, I., Zur Theorie der vertauschbaren Matrizen, *Jour. Reine und Angew. Math.* 130 (1905), 66-76.
- [21] Simpson, C. T., Higgs bundles and local systems, Preprint, Princeton Univ., 1989.
- [22] Simpson, C. T., Some families of local systems over smooth projective varieties, *Prépubl. Univ. Toulouse III*, 1991.
- [23] Siu, Y.T., Complex analyticity of harmonic maps and strong rigidity of compact Kähler manifolds, *Ann. Math.* 112 (1980), 73–111.
- [24] Siu, Y. T., Strong rigidity of compact quotients of exceptional bounded symmetric domains, *Duke Math. Jour.* 48 (1981), 857 - 871.
- [25] Siu, Y. T., Complex analyticity of harmonic maps, vanishing and Lefschetz theorems, *Jour. Diff. Geom.* 17 (1982), 55 - 138.

Authors' address:

Department of Mathematics
 University of Utah
 Salt Lake City, Utah 84112

Electronic mail:

carlson@math.utah.edu
 toledo@math.utah.edu