

Hitting probabilities for systems of non-linear stochastic heat equations with multiplicative noise

Robert C. Dalang^{1,4}, Davar Khoshnevisan^{2,5}, and Eulalia Nualart³

Abstract

We consider a system of d non-linear stochastic heat equations in spatial dimension 1 driven by d -dimensional space-time white noise. The non-linearities appear both as additive drift terms and as multipliers of the noise. Using techniques of Malliavin calculus, we establish upper and lower bounds on the one-point density of the solution $u(t, x)$, and upper bounds of Gaussian-type on the two-point density of $(u(s, y), u(t, x))$. In particular, this estimate quantifies how this density degenerates as $(s, y) \rightarrow (t, x)$. From these results, we deduce upper and lower bounds on hitting probabilities of the process $\{u(t, x)\}_{t \in \mathbb{R}_+, x \in [0, 1]}$, in terms of respectively Hausdorff measure and Newtonian capacity. These estimates make it possible to show that points are polar when $d \geq 7$ and are not polar when $d \leq 5$. We also show that the Hausdorff dimension of the range of the process is 6 when $d > 6$, and give analogous results for the processes $t \mapsto u(t, x)$ and $x \mapsto u(t, x)$. Finally, we obtain the values of the Hausdorff dimensions of the level sets of these processes.

AMS 2000 subject classifications: Primary: 60H15, 60J45; Secondary: 60H07, 60G60.

Key words and phrases. Hitting probabilities, stochastic heat equation, space-time white noise, Malliavin calculus.

¹Institut de Mathématiques, Ecole Polytechnique Fédérale de Lausanne, Station 8, CH-1015 Lausanne, Switzerland. robert.dalang@epfl.ch

²Department of Mathematics, The University of Utah, 155 S. 1400 E. Salt Lake City, UT 84112-0090, USA. davar@math.utah.edu

³Institut Galilée, Université Paris 13, 93430 Villetaneuse, France. nualart@math.univ-paris13.fr

⁴Supported in part by the Swiss National Foundation for Scientific Research.

⁵Research supported in part by a grant from the US National Science Foundation.

1 Introduction and main results

Consider the following system of non-linear stochastic partial differential equations (*spde*'s)

$$\frac{\partial u_i}{\partial t}(t, x) = \frac{\partial^2 u_i}{\partial x^2}(t, x) + \sum_{j=1}^d \sigma_{i,j}(u(t, x)) \dot{W}^j(t, x) + b_i(u(t, x)), \quad (1.1)$$

for $1 \leq i \leq d$, $t \in [0, T]$, and $x \in [0, 1]$, where $u := (u_1, \dots, u_d)$, with initial conditions $u(0, x) = 0$ for all $x \in [0, 1]$, and Neumann boundary conditions

$$\frac{\partial u_i}{\partial x}(t, 0) = \frac{\partial u_i}{\partial x}(t, 1) = 0, \quad 0 \leq t \leq T. \quad (1.2)$$

Here, $\dot{W} := (\dot{W}^1, \dots, \dot{W}^d)$ is a vector of d independent space-time white noises on $[0, T] \times [0, 1]$. For all $1 \leq i, j \leq d$, $b_i, \sigma_{i,j} : \mathbb{R}^d \rightarrow \mathbb{R}$ are globally Lipschitz functions. We set $b = (b_i)$, $\sigma = (\sigma_{i,j})$. Equation (1.1) is formal: the rigorous formulation of WALSH [W86] will be recalled in Section 2.

The objective of this paper is to develop a potential theory for the \mathbb{R}^d -valued process $u = (u(t, x), t \geq 0, x \in (0, 1))$. In particular, given $A \subset \mathbb{R}^d$, we want to determine whether the process u visits (or hits) A with positive probability.

The only potential-theoretic result that we are aware of for systems of *non-linear* spde's with multiplicative noise (σ non-constant) is DALANG AND NUALART [DN04], who study the case of the reduced hyperbolic spde on \mathbb{R}_+^2 (essentially equivalent to the wave equation in spatial dimension 1):

$$\frac{\partial^2 X_t^i}{\partial t_1 \partial t_2} = \sum_{j=1}^d \sigma_{i,j}(X_t) \frac{\partial^2 W_t^j}{\partial t_1 \partial t_2} + b_i(X_t),$$

where $t = (t_1, t_2) \in \mathbb{R}_+^2$, and $X_t^i = 0$ if $t_1 t_2 = 0$, for all $1 \leq i \leq d$. There, Dalang and Nualart used Malliavin calculus to show that the solution (X_t) of this spde satisfies

$$K^{-1} \text{Cap}_{d-4}(A) \leq \text{P}\{\exists t \in [a, b]^2 : X_t \in A\} \leq K \text{Cap}_{d-4}(A),$$

where Cap_β denotes the capacity with respect to the Newtonian β -kernel $K_\beta(\cdot)$ (see (1.6)). This result, particularly the upper bound, relies heavily on properties of the underlying two-parameter filtration and uses Cairoli's maximal inequality for two-parameter processes.

Hitting probabilities for systems of linear heat equations have been obtained in MUELLER AND TRIBE [MT03]. For systems of non-linear stochastic heat equations with *additive noise*, that is, σ in (1.1) is a constant matrix, so (1.1) becomes

$$\frac{\partial u_i}{\partial t}(t, x) = \frac{\partial^2 u_i}{\partial x^2}(t, x) + \sum_{j=1}^d \sigma_{i,j} \dot{W}^j(t, x) + b_i(u(t, x)), \quad (1.3)$$

estimates on hitting probabilities have been obtained in DALANG, KHOSHNEVISAN AND NUALART [DKN07]. That paper develops some general results that lead to upper and lower bounds on hitting probabilities for continuous two-parameter random fields, and then uses

these, together with a careful analysis of the linear equation ($b \equiv 0$, $\sigma \equiv I_d$, where I_d denotes the $d \times d$ identity matrix) and Girsanov's theorem, to deduce bounds on hitting probabilities for the solution to (1.3).

In this paper, we make use of the general results of [DKN07], but then, in order to handle the solution of (1.1), we use a very different approach. Indeed, the results of [DKN07] require in particular information about the probability density function $p_{t,x}$ of the random vector $u(t, x)$. In the case of multiplicative noise, estimates on $p_{t,x}$ can be obtained via Malliavin calculus.

We refer in particular on the results of BALLY AND PARDOUX [BP98], who used Malliavin calculus in the case $d = 1$ to prove that for any $t > 0$, $k \in \mathbb{N}$ and $0 \leq x_1 < \dots < x_k \leq 1$, the law of $(u(t, x_1), \dots, u(t, x_k))$ is absolutely continuous with respect to Lebesgue measure, with a smooth and strictly positive density on $\{\sigma \neq 0\}^k$, provided σ and b are infinitely differentiable functions which are bounded together with their derivatives of all orders. A Gaussian-type lower bound for this density is established by KOHATSU-HIGA [K03] under a uniform ellipticity condition. MORIEN [M98] showed that the density function is also Hölder-continuous as a function of (t, x) .

In this paper, we shall use techniques of Malliavin calculus to establish the following theorem. Let $p_{t,x}(z)$ denote the probability density function of the \mathbb{R}^d -valued random vector $u(t, x) = (u_1(t, x), \dots, u_d(t, x))$ and for $(s, y) \neq (t, x)$, let $p_{s,y;t,x}(z_1, z_2)$ denote the joint density of the \mathbb{R}^{2d} -valued random vector

$$(u(s, y), u(t, x)) = (u_1(s, y), \dots, u_d(s, y), u_1(t, x), \dots, u_d(t, x)) \quad (1.4)$$

(the existence of $p_{t,x}(\cdot)$ is essentially a consequence of the result of BALLY AND PARDOUX [BP98], see our Corollary 4.3; the existence of $p_{s,y;t,x}(\cdot, \cdot)$ is a consequence of Theorems 3.1 and 6.3).

Consider the following two hypotheses on the coefficients of the system (1.1):

P1 The functions σ_{ij} and b_i are bounded and infinitely differentiable with bounded partial derivatives of all orders, for $1 \leq i, j \leq d$.

P2 The matrix σ is uniformly elliptic, that is, $\|\sigma(x)\xi\|^2 \geq \rho^2 > 0$ for some $\rho > 0$, for all $x \in \mathbb{R}^d$, $\xi \in \mathbb{R}^d$, $\|\xi\| = 1$ ($\|\cdot\|$ denotes the Euclidean norm on \mathbb{R}^d).

Theorem 1.1. *Assume P1 and P2. Fix $T > 0$ and let $I \subset (0, T]$ and $J \subset (0, 1)$ be two compact nonrandom intervals.*

(a) *The density $p_{t,x}(z)$ is uniformly bounded over $z \in \mathbb{R}^d$, $t \in I$ and $x \in J$.*

(b) *There exists $c > 0$ such that for any $t \in I$, $x \in J$ and $z \in \mathbb{R}^d$,*

$$p_{t,x}(z) \geq ct^{-d/4} \exp\left(-\frac{\|z\|^2}{ct^{1/2}}\right).$$

(c) *For all $\eta > 0$, there exists $c > 0$ such that for any $s, t \in I$, $x, y \in J$, $(s, y) \neq (t, x)$ and $z_1, z_2 \in \mathbb{R}^d$,*

$$p_{s,y;t,x}(z_1, z_2) \leq c(|t - s|^{1/2} + |x - y|)^{-(d+\eta)/2} \exp\left(-\frac{\|z_1 - z_2\|^2}{c(|t - s|^{1/2} + |x - y|)}\right). \quad (1.5)$$

(d) *There exists $c > 0$ such that for any $t \in I$, $x, y \in J$, $x \neq y$ and $z_1, z_2 \in \mathbb{R}^d$,*

$$p_{t,y;t,x}(z_1, z_2) \leq c(|x - y|)^{-d/2} \exp\left(-\frac{\|z_1 - z_2\|^2}{c|x - y|}\right).$$

The main technical effort in this paper is to obtain the upper bound in (c). Indeed, it is not difficult to check that for fixed $(s, y; t, x)$, $(z_1, z_2) \mapsto p_{s,y;t,x}(z_1, z_2)$ behaves like a Gaussian density function. However, for $(s, y) = (t, x)$, the \mathbb{R}^{2d} -valued random vector $(u(s, y), u(t, x))$ is concentrated on a d -dimensional subspace in \mathbb{R}^{2d} and therefore does not have a density with respect to Lebesgue measure in \mathbb{R}^{2d} . So the main effort is to estimate how this density blows up as $(s, y) \rightarrow (t, x)$. This is achieved by a detailed analysis of the behavior of the Malliavin matrix of $(u(s, y), u(t, x))$ as a function of $(s, y; t, x)$, using a perturbation argument. The presence of η in statement (c) may be due to the method of proof. When $t = s$, it is possible to set $\eta = 0$ as in Theorem 1.1(d).

This paper is organized as follows. After introducing some notation and stating our main results on hitting probabilities (Theorems 1.2 and 1.6), we assume Theorem 1.1 and use the theorems of [DKN07] to prove these results in Section 2. In Section 3, we recall some basic facts of Malliavin calculus and state and prove two results that are tailored to our needs (Propositions 3.4 and 3.5). In Section 4, we establish the existence, smoothness and uniform boundedness of the one-point density function $p_{t,x}$, proving Theorem 1.1(a). In Section 5, we establish a lower bound on $p_{t,x}$, which proves Theorem 1.1(b). This upper (respectively lower) bound is a fairly direct extension to $d \geq 1$ of a result of BALLY AND PARDOUX [BP98] (respectively KOHATSU-HIGA [K03]) when $d = 1$. In Section 6, we establish Theorem 1.1(c) and (d). The main steps are as follows.

The upper bound on the two-point density function $p_{s,y;t,x}$ involves a bloc-decomposition of the Malliavin matrix of the \mathbb{R}^{2d} -valued random vector $(u(s, y), u(s, y) - u(t, x))$. The entries of this matrix are of different orders of magnitude, depending on which bloc they are in: see Theorem 6.3. Assuming Theorem 6.3, we prove Theorem 1.1(c) and (d) in Section 6.3. The exponential factor in (1.5) is obtained from an exponential martingale inequality, while the factor $(|t - s|^{1/2} + |x - y|)^{-(d+\eta)/2}$ comes from an estimate of the iterated Skorohod integrals that appear in Corollary 3.3 and from the block structure of the Malliavin matrix.

The proof of Theorem 6.3 is presented in Section 6.4: this is the main technical effort in this paper. We need bounds on the inverse of the Malliavin matrix. Bounds on its cofactors are given in Proposition 6.5, while bounds on negative moments of its determinant are given in Proposition 6.6. The determinant is equal to the product of the $2d$ eigenvalues of the Malliavin matrix. It turns out that at least d of these eigenvalues are of order 1 (“large eigenvalues”) and do not contribute to the upper bound in (1.5), and at most d are of the same order as the smallest eigenvalue (“small eigenvalues”), that is, of order $|t - s|^{1/2} + |x - y|$. If we did not distinguish between these two types of eigenvalues, but estimated all of them by the smallest eigenvalue, we would obtain a factor of $(|t - s|^{1/2} + |x - y|)^{-d+\eta/2}$ in (1.5), which would *not* be the correct order. The estimates on the smallest eigenvalue are obtained by refining a technique that appears in [BP98]; indeed, we obtain a precise estimate on the density whereas they only showed existence. The study of the large eigenvalues does not seem to appear elsewhere in the literature.

Coming back to potential theory, let us introduce some notation. For all Borel sets $F \subset \mathbb{R}^d$, we define $\mathcal{P}(F)$ to be the set of all probability measures with compact support contained in F . For all integers $k \geq 1$ and $\mu \in \mathcal{P}(\mathbb{R}^k)$, we let $I_\beta(\mu)$ denote the β -dimensional energy of μ , that is,

$$I_\beta(\mu) := \iint \mathbb{K}_\beta(\|x - y\|) \mu(dx) \mu(dy),$$

where $\|x\|$ denotes the Euclidian norm of $x \in \mathbb{R}^k$,

$$\mathbb{K}_\beta(r) := \begin{cases} r^{-\beta} & \text{if } \beta > 0, \\ \log(N_0/r) & \text{if } \beta = 0, \\ 1 & \text{if } \beta < 0, \end{cases} \quad (1.6)$$

and N_0 is a sufficiently large constant (see DALANG, KHOSHNEVISAN, AND NUALART [DKN07, (1.5)]).

For all $\beta \in \mathbb{R}$, integers $k \geq 1$, and Borel sets $F \subset \mathbb{R}^k$, $\text{Cap}_\beta(F)$ denotes the β -dimensional capacity of F , that is,

$$\text{Cap}_\beta(F) := \left[\inf_{\mu \in \mathcal{P}(F)} I_\beta(\mu) \right]^{-1},$$

where $1/\infty := 0$. Note that if $\beta < 0$, then $\text{Cap}_\beta(\cdot) \equiv 1$.

Given $\beta \geq 0$, the β -dimensional Hausdorff measure of F is defined by

$$\mathcal{H}_\beta(F) = \liminf_{\epsilon \rightarrow 0^+} \left\{ \sum_{i=1}^{\infty} (2r_i)^\beta : F \subseteq \bigcup_{i=1}^{\infty} B(x_i, r_i), \sup_{i \geq 1} r_i \leq \epsilon \right\}, \quad (1.7)$$

where $B(x, r)$ denotes the open (Euclidean) ball of radius $r > 0$ centered at $x \in \mathbb{R}^d$. When $\beta < 0$, we define $\mathcal{H}_\beta(F)$ to be infinite.

Throughout, we consider the following *parabolic metric*: For all $s, t \in [0, T]$ and $x, y \in [0, 1]$,

$$\Delta((t, x); (s, y)) := |t - s|^{1/2} + |x - y|. \quad (1.8)$$

Clearly, this is a metric on \mathbb{R}^2 which generates the usual Euclidean topology on \mathbb{R}^2 . Then we obtain an energy form

$$I_\beta^\Delta(\mu) := \iint \mathbb{K}_\beta(\Delta((t, x); (s, y))) \mu(dt dx) \mu(ds dy),$$

and a corresponding capacity

$$\text{Cap}_\beta^\Delta(F) := \left[\inf_{\mu \in \mathcal{P}(F)} I_\beta^\Delta(\mu) \right]^{-1}.$$

For the Hausdorff measure, we write

$$\mathcal{H}_\beta^\Delta(F) = \liminf_{\epsilon \rightarrow 0^+} \left\{ \sum_{i=1}^{\infty} (2r_i)^\beta : F \subseteq \bigcup_{i=1}^{\infty} B^\Delta((t_i, x_i), r_i), \sup_{i \geq 1} r_i \leq \epsilon \right\},$$

where $B^\Delta((t, x), r)$ denotes the open Δ -ball of radius $r > 0$ centered at $(t, x) \in [0, T] \times [0, 1]$.

Using Theorem 1.1 together with results from DALANG, KHOSHNEVISAN, AND NUALART [DKN07], we shall prove the following result. Let $u(E)$ denote the (random) range of E under the map $(t, x) \mapsto u(t, x)$, where E is some Borel-measurable subset of \mathbb{R}^2 .

Theorem 1.2. *Assume **P1** and **P2**. Fix $T > 0$, $M > 0$, and $\eta > 0$. Let $I \subset (0, T]$ and $J \subset (0, 1)$ be two fixed non-trivial compact intervals.*

- (a) *There exists $c > 0$ depending on M, I, J and η such that for all compact sets $A \subseteq [-M, M]^d$,*

$$c^{-1} \text{Cap}_{d-6+\eta}(A) \leq \mathbb{P}\{u(I \times J) \cap A \neq \emptyset\} \leq c \mathcal{H}_{d-6-\eta}(A).$$

- (b) *For all $t \in (0, T]$, there exists $c_1 > 0$ depending on T, M and J , and $c_2 > 0$ depending on T, M, J and $\eta > 0$ such that for all compact sets $A \subseteq [-M, M]^d$,*

$$c_1 \text{Cap}_{d-2}(A) \leq \mathbb{P}\{u(\{t\} \times J) \cap A \neq \emptyset\} \leq c_2 \mathcal{H}_{d-2-\eta}(A).$$

- (c) *For all $x \in (0, 1)$, there exists $c > 0$ depending on M, I and η such that for all compact sets $A \subseteq [-M, M]^d$,*

$$c^{-1} \text{Cap}_{d-4+\eta}(A) \leq \mathbb{P}\{u(I \times \{x\}) \cap A \neq \emptyset\} \leq c \mathcal{H}_{d-4-\eta}(A).$$

Remark 1.3. (i) Because of the inequalities between capacity and Hausdorff measure, the right-hand sides of Theorem 1.2 can be replaced by $c \text{Cap}_{d-6-\eta}(A)$, $c \text{Cap}_{d-2-\eta}(A)$ and $c \text{Cap}_{d-4-\eta}(A)$ in (a), (b) and (c), respectively (cf. KAHANE [K85, p. 133]).

- (ii) Theorem 1.2 also holds if we consider Dirichlet boundary conditions (i.e. $u_i(t, 0) = u_i(t, 1) = 0$, for $t \in [0, T]$) instead of Neumann boundary conditions.

- (iii) In the upper bounds of Theorem 1.2, the condition in **P1** that σ and b are bounded can be removed, but their derivatives of all orders must exist and be bounded.

As a consequence of Theorem 1.2, we deduce the following result on the polarity of points. Recall that a Borel set $A \subseteq \mathbb{R}^d$ is called *polar* for u if $\mathbb{P}\{u((0, T] \times (0, 1)) \cap A \neq \emptyset\} = 0$; otherwise, A is called *nonpolar*.

Corollary 1.4. *Assume **P1** and **P2**.*

- (a) *Singletons are nonpolar for $(t, x) \mapsto u(t, x)$ when $d \leq 5$, and are polar when $d \geq 7$ (the case $d = 6$ is open).*
- (b) *Fix $t \in (0, T]$. Singletons are nonpolar for $x \mapsto u(t, x)$ when $d = 1$, and are polar when $d \geq 3$ (the case $d = 2$ is open).*
- (c) *Fix $x \in (0, 1)$. Singletons are not polar for $t \mapsto u(t, x)$ when $d \leq 3$ and are polar when $d \geq 5$ (the case $d = 4$ is open).*

Another consequence of Theorem 1.2 is the Hausdorff dimension of the range of the process u .

Corollary 1.5. *Assume **P1** and **P2**.*

- (a) *If $d > 6$, then $\dim_{\mathbb{H}}(u((0, T] \times (0, 1))) = 6$ a.s.*
- (b) *Fix $t \in \mathbb{R}_+$. If $d > 2$, then $\dim_{\mathbb{H}}(u(\{t\} \times (0, 1))) = 2$ a.s.*
- (c) *Fix $x \in (0, 1)$. If $d > 4$, then $\dim_{\mathbb{H}}(u(\mathbb{R}_+ \times \{x\})) = 4$ a.s.*

As in DALANG, KHOSHNEVISAN, AND NUALART [DKN07], it is also possible to use Theorem 1.1 to obtain results concerning level sets of u . Define

$$\begin{aligned}\mathcal{L}(z; u) &:= \{(t, x) \in I \times J : u(t, x) = z\}, \\ \mathcal{T}(z; u) &= \{t \in I : u(t, x) = z \text{ for some } x \in J\}, \\ \mathcal{X}(z; u) &= \{x \in J : u(t, x) = z \text{ for some } t \in I\}, \\ \mathcal{L}_x(z; u) &:= \{t \in I : u(t, x) = z\}, \\ \mathcal{L}^t(z; u) &:= \{x \in J : u(t, x) = z\}.\end{aligned}$$

We note that $\mathcal{L}(z; u)$ is the level set of u at level z , $\mathcal{T}(z; u)$ (resp. $\mathcal{X}(z; u)$) is the projection of $\mathcal{L}(z; u)$ onto I (resp. J), and $\mathcal{L}_x(z; u)$ (resp. $\mathcal{L}^t(z; u)$) is the x -section (resp. t -section) of $\mathcal{L}(z; u)$.

Theorem 1.6. *Assume **P1** and **P2**. Then for all $\eta > 0$ and $R > 0$ there exists a positive and finite constant c such that the following holds for all compact sets $E \subset (0, T] \times (0, 1)$, $F \subset (0, T]$, $G \subset (0, 1)$, and for all $z \in B(0, R)$:*

- (a) $c^{-1} \text{Cap}_{(d+\eta)/2}^{\Delta}(E) \leq \mathbb{P}\{\mathcal{L}(z; u) \cap E \neq \emptyset\} \leq c \mathcal{H}_{(d-\eta)/2}^{\Delta}(E)$;
- (b) $c^{-1} \text{Cap}_{(d-2+\eta)/4}(F) \leq \mathbb{P}\{\mathcal{T}(z; u) \cap F \neq \emptyset\} \leq c \mathcal{H}_{(d-2-\eta)/4}(F)$;
- (c) $c^{-1} \text{Cap}_{(d-4+\eta)/2}(G) \leq \mathbb{P}\{\mathcal{X}(z; u) \cap G \neq \emptyset\} \leq c \mathcal{H}_{(d-4-\eta)/2}(G)$;
- (d) *for all $x \in (0, 1)$, $c^{-1} \text{Cap}_{(d+\eta)/4}(F) \leq \mathbb{P}\{\mathcal{L}_x(z; u) \cap F \neq \emptyset\} \leq c \mathcal{H}_{(d-\eta)/4}(F)$;*
- (e) *for all $t \in (0, T]$, $c^{-1} \text{Cap}_{d/2}(G) \leq \mathbb{P}\{\mathcal{L}^t(z; u) \cap G \neq \emptyset\} \leq c \mathcal{H}_{(d-\eta)/2}(G)$.*

Corollary 1.7. *Assume **P1** and **P2**. Choose and fix $z \in \mathbb{R}^d$.*

- (a) *If $2 < d < 6$, then $\dim_{\mathbb{H}} \mathcal{T}(z; u) = \frac{1}{4}(6 - d)$ a.s. on $\{\mathcal{T}(z; u) \neq \emptyset\}$.*
- (b) *If $4 < d < 6$ (i.e. $d = 5$), then $\dim_{\mathbb{H}} \mathcal{X}(z; u) = \frac{1}{2}(6 - d)$ a.s. on $\{\mathcal{X}(z; u) \neq \emptyset\}$.*
- (c) *If $1 \leq d < 4$, then $\dim_{\mathbb{H}} \mathcal{L}_x(z; u) = \frac{1}{4}(4 - d)$ a.s. on $\{\mathcal{L}_x(z; u) \neq \emptyset\}$.*
- (d) *If $d = 1$, then $\dim_{\mathbb{H}} \mathcal{L}^t(z; u) = \frac{1}{2}(2 - d) = \frac{1}{2}$ a.s. on $\{\mathcal{L}^t(z; u) \neq \emptyset\}$.*

In addition, all four right-most events have positive probability.

Remark 1.8. The results of the two theorems and corollaries above should be compared with those of DALANG, KHOSHNEVISAN AND NUALART [DKN07].

2 Proof of Theorems 1.2, 1.6 and their corollaries (assuming Theorem 1.1)

We first recall that equation (1.1) is formal: a rigorous formulation, following WALSH [W86], is as follows. Let $W^i = (W^i(s, x))_{s \in \mathbb{R}_+, x \in [0, 1]}$, $i = 1, \dots, d$, be independent Brownian sheets defined on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$, and set $W = (W^1, \dots, W^d)$. For $t \geq 0$, let $\mathcal{F}_t = \sigma\{W(s, x), s \in [0, t], x \in [0, 1]\}$. We say that a process $u = \{u(t, x), t \in [0, T], x \in [0, 1]\}$ is adapted to (\mathcal{F}_t) if $u(t, x)$ is \mathcal{F}_t -measurable for each $(t, x) \in [0, T] \times [0, 1]$. We say that u is a solution of (1.1) if u is adapted to (\mathcal{F}_t) and if for $i \in \{1, \dots, d\}$,

$$\begin{aligned} u_i(t, x) = & \int_0^t \int_0^1 G_{t-r}(x, v) \sum_{j=1}^d \sigma_{i,j}(u(r, v)) W^j(dr, dv) \\ & + \int_0^t \int_0^1 G_{t-r}(x, v) b_i(u(r, v)) dr dv, \end{aligned} \quad (2.1)$$

where $G_t(x, y)$ denotes the Green kernel for the heat equation with Neumann boundary conditions (see WALSH [W86, Chap 3]), and the stochastic integral in (2.1) is interpreted as in [W86].

Adapting the results from [W86] to the case $d \geq 1$, one can show that there exists a unique continuous process $u = \{u(t, x), t \in [0, T], x \in [0, 1]\}$ adapted to (\mathcal{F}_t) that is a solution of (1.1). Moreover, it is shown in BALLY, MILLET, AND SANZ-SOLÉ [BMS95] that for any $s, t \in [0, T]$ with $s \leq t$, $x, y \in [0, 1]$, and $p > 1$,

$$\mathbb{E}[|u(t, x) - u(s, y)|^p] \leq C_{T,p}(\Delta((t, x); (s, y)))^{p/2}, \quad (2.2)$$

where Δ is the parabolic metric defined in (1.8). In particular, for any $0 < \alpha < 1/2$, u is a.s. α -Hölder continuous in x and $\alpha/2$ -Hölder continuous in t .

Assuming Theorem 1.1, we now prove Theorems 1.2, 1.6 and their corollaries.

Proof of Theorem 1.2. (a) In order to prove the upper bound we use DALANG, KHOSHNEVISAN, AND NUALART [DKN07, Theorem 3.3]. Indeed, Theorem 1.1(a) and (2.2) imply that the hypotheses (i) and (ii), respectively, of this theorem, are satisfied, and so the conclusion (with $\beta = d - \eta$) is too.

In order to prove the lower bound, we shall use of [DKN07, Theorem 2.1]. This requires checking hypotheses **A1** and **A2** in that paper. Hypothesis **A1** is a lower bound on the one-point density function $p_{t,x}(z)$, which is an immediate consequence of Theorem 1.1(b). Hypothesis **A2** is an upper bound on the two-point density function $p_{s,y;t,x}(z_1, z_2)$, which involves a parameter β ; we take $\beta = d + \eta$. In this case, Hypothesis **A2** is an immediate consequence of Theorem 1.1(c). Therefore, the lower bound in Theorem 1.2(a) follows from [DKN07, Theorem 2.1]. This proves (a).

(b) For the upper bound, we again refer to [DKN07, Theorem 3.3] (see also [DKN07, Theorem 3.1]). For the lower bound, which involves $\text{Cap}_{d-2}(A)$ instead of $\text{Cap}_{d-2+\eta}(A)$, we refer to [DKN07, Remark 2.5] and observe that hypotheses **A1**^t and **A2**^t there are satisfied with $\beta = d$ (by Theorem 1.1(d)). This proves (b).

(c) As in (a), the upper bound follows from [DKN07, Theorem 3.3] with $\beta = d - \eta$ (see also [DKN07, Theorem 3.1(3)]), and the lower bound follows from [DKN07, Theorem 2.1(3)], with $\beta = d + \eta$. Theorem 1.2 is proved. \square

Proof of Corollary 1.4. We first prove (a). Let $z \in \mathbb{R}^d$. If $d \leq 5$, then there is $\eta > 0$ such that $d - 6 + \eta < 0$, and thus $\text{Cap}_{d-6+\eta}(\{z\}) = 1$. Hence, the lower bound of Theorem 1.2 (a) implies that $\{z\}$ is not polar. On the other hand, if $d > 6$, then for small $\eta > 0$, $d - 6 - \eta > 0$. Therefore, $\mathcal{H}_{d-6-\eta}(\{z\}) = 0$ and the upper bound of Theorem 1.2(a) implies that $\{z\}$ is polar. This proves (a). One proves (b) and (c) exactly along the same lines using Theorem 1.2(b) and (c). \square

Proof of Theorem 1.6. For the upper bounds in (a)-(e), we use DALANG, KHOSHNEVISAN, AND NUALART [DKN07, Theorem 3.3] whose assumptions we verified above with $\beta = d - \eta$; these upper bounds then follow immediately from [DKN07, Theorem 3.2].

For the lower bounds in (a)-(d), we use [DKN07, Theorem 2.4] since we have shown above that the assumptions of this theorem, with $\beta = d + \eta$, are satisfied by Theorem 1.1. For the lower bound in (e), we refer to [DKN07, Remark 2.5] and note that by Theorem 1.1(d), Hypothesis **A2**^t there is satisfied with $\beta = d$. This proves Theorem 1.6. \square

Proof of Corollaries 1.5 and 1.7. The final positive-probability assertion in Corollary 1.7 is an immediate consequence of Theorem 1.6 and Taylor's theorem KHOSHNEVISAN [K02, Corollary 2.3.1 p. 523].

Let E be a random set. When it exists, the codimension of E is the real number $\beta \in [0, d]$ such that for all compact sets $A \subset \mathbb{R}^d$,

$$\mathbb{P}\{E \cap A \neq \emptyset\} \begin{cases} > 0 & \text{whenever } \dim_{\text{H}}(A) > \beta, \\ = 0 & \text{whenever } \dim_{\text{H}}(A) < \beta. \end{cases}$$

See KHOSHNEVISAN [K02, Chap.11, Section 4]. When it is well defined, we write the said codimension as $\text{codim}(E)$. Theorems 1.2 and 1.6 imply that for $d \geq 1$: $\text{codim}(u(\mathbb{R}_+ \times (0, 1))) = (d - 6)^+$; $\text{codim}(u(\{t\} \times (0, 1))) = (d - 2)^+$; $\text{codim}(u(\mathbb{R}_+ \times \{x\})) = (d - 4)^+$; $\text{codim}(\mathcal{T}(z)) = (\frac{d-2}{4})^+$; $\text{codim}(\mathcal{X}(z)) = (\frac{d-4}{2})^+$; $\text{codim}(\mathcal{L}_x(z)) = \frac{d}{4}$; and $\text{codim}(\mathcal{L}^t(z)) = \frac{d}{2}$. According to Theorem 4.7.1 of KHOSHNEVISAN [K02, Chapter 11], given a random set E in \mathbb{R}^n whose codimension is strictly between 0 and n ,

$$\dim_{\text{H}} E + \text{codim } E = n \quad \text{a.s. on } \{E \neq \emptyset\}. \quad (2.3)$$

This implies the statements of Corollaries 1.5 and 1.7. \square

3 Elements of Malliavin calculus

In this section, we introduce, following NUALART [N95] (see also SANZ-SOLÉ [S05]), some elements of Malliavin calculus. Let \mathcal{S} denote the class of smooth random variables of the form

$$F = f(W(h_1), \dots, W(h_n)),$$

where $n \geq 1$, $f \in \mathcal{C}_P^\infty(\mathbb{R}^n)$, the set of real-valued functions f such that f and all its partial derivatives have at most polynomial growth, $h_i \in \mathcal{H} := L^2([0, T] \times [0, 1], \mathbb{R}^d)$, and $W(h_i)$ denotes the Wiener integral

$$W(h_i) = \int_0^T \int_0^1 h_i(t, x) \cdot W(dx, dt), \quad 1 \leq i \leq n.$$

Given $F \in \mathcal{S}$, its derivative is defined to be the \mathbb{R}^d -valued stochastic process $DF = (D_{t,x}F = (D_{t,x}^{(1)}F, \dots, D_{t,x}^{(d)}F), (t, x) \in [0, T] \times [0, 1])$ given by

$$D_{t,x}F = \sum_{i=1}^n \frac{\partial f}{\partial x_i}(W(h_1), \dots, W(h_n)) h_i(t, x).$$

More generally, we can define the derivative $D^k F$ of order k of F by setting

$$D_\alpha^k F = \sum_{i_1, \dots, i_k=1}^n \frac{\partial}{\partial x_{i_1}} \cdots \frac{\partial}{\partial x_{i_k}} f(W(h_1), \dots, W(h_n)) h_{i_1}(\alpha_1) \otimes \cdots \otimes h_{i_k}(\alpha_k),$$

where $\alpha = (\alpha_1, \dots, \alpha_k)$, and $\alpha_i = (t_i, x_i)$, $1 \leq i \leq k$.

For $p, k \geq 1$, the space $\mathbb{D}^{k,p}$ is the closure of \mathcal{S} with respect to the seminorm $\|\cdot\|_{k,p}^p$ defined by

$$\|F\|_{k,p}^p = \mathbb{E}[|F|^p] + \sum_{j=1}^k \mathbb{E}[\|D^j F\|_{\mathcal{H}^{\otimes j}}^p],$$

where

$$\|D^j F\|_{\mathcal{H}^{\otimes j}}^2 = \sum_{i_1, \dots, i_j=1}^d \int_0^T dt_1 \int_0^1 dx_1 \cdots \int_0^T dt_j \int_0^1 dx_j \left(D_{(t_1, x_1)}^{(i_1)} \cdots D_{(t_j, x_j)}^{(i_j)} F \right)^2.$$

We set $(\mathbb{D}^\infty)^d = \bigcap_{p \geq 1} \bigcap_{k \geq 1} \mathbb{D}^{k,p}$.

The derivative operator D on $L^2(\Omega)$ has an adjoint, termed the Skorohod integral and denoted by δ , which is an unbounded operator on $L^2(\Omega, \mathcal{H})$. Its domain, denoted by $\text{Dom } \delta$, is the set of elements $u \in L^2(\Omega, \mathcal{H})$ such that there exists a constant c such that $|\mathbb{E}[\langle DF, u \rangle_{\mathcal{H}}]| \leq c \|F\|_{0,2}$, for any $F \in \mathbb{D}^{1,2}$.

If $u \in \text{Dom } \delta$, then $\delta(u)$ is the element of $L^2(\Omega)$ characterized by the following duality relation:

$$\mathbb{E}[F \delta(u)] = \mathbb{E} \left[\sum_{j=1}^d \int_0^T \int_0^1 D_{t,x}^{(j)} F u_j(t, x) dt dx \right], \quad \text{for all } F \in \mathbb{D}^{1,2}.$$

A first application of Malliavin calculus to the study of probability laws is the following global criterion for smoothness of densities.

Theorem 3.1. [N95, Thm.2.1.2 and Cor.2.1.2] or [S05, Thm.5.2] *Let $F = (F^1, \dots, F^d)$ be an \mathbb{R}^d -valued random vector satisfying the following two conditions:*

- (i) $F \in (\mathbb{D}^\infty)^d$;
- (ii) the Malliavin matrix of F defined by $\gamma_F = (\langle DF^i, DF^j \rangle_{\mathcal{H}})_{1 \leq i, j \leq d}$ is invertible a.s. and $(\det \gamma_F)^{-1} \in L^p(\Omega)$ for all $p \geq 1$.

Then the probability law of F has an infinitely differentiable density function.

A random vector F that satisfies conditions (i) and (ii) of Theorem 3.1 is said to be nondegenerate. For a nondegenerate random vector, the following integration by parts formula plays a key role.

Proposition 3.2. [N98, Prop.3.2.1] or [S05, Prop.5.4] *Let $F = (F^1, \dots, F^d) \in (\mathbb{D}^\infty)^d$ be a nondegenerate random vector, let $G \in \mathbb{D}^\infty$ and let $g \in \mathcal{C}_P^\infty(\mathbb{R}^d)$. Fix $k \geq 1$. Then for any multi-index $\alpha = (\alpha_1, \dots, \alpha_k) \in \{1, \dots, d\}^k$, there is an element $H_\alpha(F, G) \in \mathbb{D}^\infty$ such that*

$$\mathbb{E}[(\partial_\alpha g)(F)G] = \mathbb{E}[g(F)H_\alpha(F, G)].$$

In fact, the random variables $H_\alpha(F, G)$ are recursively given by

$$\begin{aligned} H_\alpha(F, G) &= H_{(\alpha_k)}(F, H_{(\alpha_1, \dots, \alpha_{k-1})}(F, G)), \\ H_{(i)}(F, G) &= \sum_{j=1}^d \delta(G(\gamma_F^{-1})_{i,j} DF^j). \end{aligned}$$

Proposition 3.2 with $G = 1$ and $\alpha = (1, \dots, d)$ implies the following expression for the density of a nondegenerate random vector.

Corollary 3.3. [N98, Corollary 3.2.1] *Let $F = (F^1, \dots, F^d) \in (\mathbb{D}^\infty)^d$ be a nondegenerate random vector and let $p_F(z)$ denote the density of F . Then for every subset σ of the set of indices $\{1, \dots, d\}$,*

$$p_F(z) = (-1)^{d-|\sigma|} \mathbb{E}[1_{\{F^i > z^i, i \in \sigma, F^i < z^i, i \notin \sigma\}} H_{(1, \dots, d)}(F, 1)],$$

where $|\sigma|$ is the cardinality of σ , and, in agreement with Proposition 3.2,

$$H_{(1, \dots, d)}(F, 1) = \delta((\gamma_F^{-1} DF)^d \delta((\gamma_F^{-1} DF)^{d-1} \delta(\dots \delta((\gamma_F^{-1} DF)^1) \dots))).$$

The next result gives a criterion for uniform boundedness of the density of a nondegenerate random vector.

Proposition 3.4. *For all $p > 1$ and $\ell \geq 1$, let $c_1 = c_1(p) > 0$ and $c_2 = c_2(\ell, p) \geq 0$ be fixed. Let $F \in (\mathbb{D}^\infty)^d$ be a nondegenerate random vector such that*

- (a) $\mathbb{E}[(\det \gamma_F)^{-p}] \leq c_1$;
- (b) $\mathbb{E}[\|D^l(F^i)\|_{\mathcal{H}^{\otimes \ell}}^p] \leq c_2$, $i = 1, \dots, d$.

Then the density of F is uniformly bounded, and the bound does not depend on F but only on the constants $c_1(p)$ and $c_2(\ell, p)$.

Proof. The proof of this result uses the same arguments as in the proof of DALANG AND NUALART [DN04, Lemma 4.11]. Therefore, we will only give the main steps.

Fix $z \in \mathbb{R}^d$. Thanks to Corollary 3.3 and the Cauchy-Schwarz inequality we find that

$$|p_F(z)| \leq \|H_{(1,\dots,d)}(F, 1)\|_{0,2}.$$

Using the continuity of the Skorohod integral δ (cf. NUALART [N95, Proposition 3.2.1] and NUALART [N98, (1.11) and p.131]) and Hölder's inequality for Malliavin norms (cf. WATANABE [W84, Proposition 1.10, p.50]), we obtain

$$\|H_{(1,\dots,d)}(F, 1)\|_{0,2} \leq c \|H_{(1,\dots,d-1)}(F, 1)\|_{1,4} \sum_{j=1}^d \|(\gamma_F^{-1})_{d,j}\|_{1,8} \|D(F^j)\|_{1,8}. \quad (3.1)$$

In agreement with hypothesis (b), $\|D(F^j)\|_{m,p} \leq c$. In order to bound the second factor in (3.1), note that

$$\|(\gamma_F^{-1})_{i,j}\|_{m,p} = \left\{ \mathbb{E}[|(\gamma_F^{-1})_{i,j}|^p] + \sum_{k=1}^m \mathbb{E}[\|D^k(\gamma_F^{-1})_{i,j}\|_{\mathcal{H}^{\otimes k}}^p] \right\}^{1/p}. \quad (3.2)$$

For the first term in (3.2), we use Cramer's formula to get that

$$|(\gamma_F^{-1})_{i,j}| = |(\det \gamma_F)^{-1} (A_F)_{i,j}|,$$

where A_F denotes the cofactor matrix of γ_F . By means of Cauchy-Schwarz inequality and hypotheses (a) and (b) we find that

$$\begin{aligned} \mathbb{E}[|(\gamma_F^{-1})_{i,j}|^p] &\leq c_{d,p} \{ \mathbb{E}[(\det \gamma_F)^{-2p}] \}^{1/2} \times \{ \mathbb{E}[\|D(F)\|_{\mathcal{H}}^{4p(d-1)}] \}^{1/2} \\ &\leq c_{d,p}, \end{aligned}$$

where none of the constants depend on F . For the second term on the right-hand side of (3.2), we iterate the equality (cf. NUALART [N95, Lemma 2.1.6])

$$D(\gamma_F^{-1})_{i,j} = - \sum_{k,\ell=1}^d (\gamma_F^{-1})_{i,k} D(\gamma_F)_{k,\ell} (\gamma_F^{-1})_{\ell,j}, \quad (3.3)$$

in the same way as in the proof of DALANG AND NUALART [DN04, Lemma 4.11]. Then, appealing again to hypotheses (a) and (b) and iterating the inequality (3.1) to bound the first factor on the right-hand side of (3.2), we obtain the uniform boundedness of $p_F(z)$. \square

We finish this section with a result that will be used later on to bound negative moments of a random variable, as is needed to check hypothesis (a) of Proposition 3.4.

Proposition 3.5. *Suppose $Z \geq 0$ is a random variable for which we can find $\epsilon_0 \in (0, 1)$, processes $\{Y_{i,\epsilon}\}_{\epsilon \in (0,1)}$ ($i = 1, 2$), and constants $c > 0$ and $0 \leq \alpha_2 \leq \alpha_1$ with the property that*

$Z \geq \min(c\epsilon^{\alpha_1} - Y_{1,\epsilon} c\epsilon^{\alpha_2} - Y_{2,\epsilon})$ for all $\epsilon \in (0, \epsilon_0)$. Also suppose that we can find $\beta_i > \alpha_i$ ($i = 1, 2$), not depending on ϵ_0 , such that

$$C(q) := \sup_{0 < \epsilon < 1} \max \left(\frac{\mathbf{E}[|Y_{1,\epsilon}|^q]}{\epsilon^{q\beta_1}}, \frac{\mathbf{E}[|Y_{2,\epsilon}|^q]}{\epsilon^{q\beta_2}} \right) < \infty \quad \text{for all } q \geq 1.$$

Then for all $p \geq 1$, there exists a constant $c' \in (0, \infty)$, not depending on ϵ_0 , such that $\mathbf{E}[|Z|^{-p}] \leq c'\epsilon_0^{-p\alpha_1}$.

Remark 3.6. This lemma is of interest mainly when $\beta_2 \leq \alpha_1$.

Proof. Define $k := (2/c)\epsilon_0^{-\alpha_1}$. Suppose that $y \geq k$, and let $\epsilon := (2/c)^{1/\alpha_1}y^{-1/\alpha_1}$. Then $0 < \epsilon \leq \epsilon_0$, $y^{-1} = (c/2)\epsilon^{\alpha_1}$, and for all $q \geq 1$,

$$\begin{aligned} \mathbf{P}\{Z^{-1} > y\} &= \mathbf{P}\{Z < y^{-1}\} \\ &\leq \mathbf{P}\left\{Y_{1,\epsilon} \geq \frac{c}{2}\epsilon^{\alpha_1}\right\} + \mathbf{P}\left\{Y_{2,\epsilon} \geq c\epsilon^{\alpha_2} - \frac{c}{2}\epsilon^{\alpha_1}\right\} \\ &\leq \frac{C(q)}{c^q} \left(2^q \epsilon^{q(\beta_1 - \alpha_1)} + \epsilon^{q\beta_2} \left[\epsilon^{\alpha_2} - \frac{1}{2}\epsilon^{\alpha_1} \right]^{-q} \right). \end{aligned}$$

The inequality $\epsilon^{\alpha_2} - (1/2)\epsilon^{\alpha_1} \geq (1/2)\epsilon^{\alpha_2}$ implies that

$$\mathbf{P}\{Z^{-1} > y\} \leq \frac{C(q)}{c^q} \left(2^q \epsilon^{q(\beta_1 - \alpha_1)} + 2^q \epsilon^{q(\beta_2 - \alpha_2)} \right) \leq ay^{-qb},$$

where a and b are positive and finite constants that do not depend on y, ϵ_0 or q . We apply this with $q := (p/b) + 1$ to find that for all $p \geq 1$,

$$\begin{aligned} \mathbf{E}[|Z|^{-p}] &= p \int_0^\infty y^{p-1} \mathbf{P}\{Z^{-1} > y\} dy \\ &\leq k^p + ap \int_k^\infty y^{-b-1} dy = k^p + \left(\frac{ap}{b}\right) k^{-b}. \end{aligned}$$

Because $k \geq (2/c)$ and $b > 0$, it follows that $\mathbf{E}[|Z|^{-p}] \leq (1 + c_1(ap/b))k^p$, where $c_1 := (c/2)^{b+p}$. This is the desired result. \square

4 Existence, smoothness and uniform boundedness of the one-point density

Let $u = \{u(t, x), t \in [0, T], x \in [0, 1]\}$ be the solution of equation (2.1). In this section, we prove the existence, smoothness and uniform boundedness of the density of the random vector $u(t, x)$. In particular, this will prove Theorem 1.1(a).

The first result concerns the Malliavin differentiability of u and the equations satisfied by its derivatives. We refer to BALLY AND PARDOUX [BP98, Proposition 4.3, (4.16), (4.17)] for its proof in dimension one. As we work coordinate by coordinate, the following proposition follows in the same way and its proof is therefore omitted.

Proposition 4.1. *Assume P1. Then $u(t, x) \in (\mathbb{D}^\infty)^d$ for any $t \in [0, T]$ and $x \in [0, 1]$. Moreover, its iterated derivative satisfies*

$$\begin{aligned} & D_{r_1, v_1}^{(k_1)} \cdots D_{r_n, v_n}^{(k_n)} (u_i(t, x)) \\ &= \sum_{l=1}^d G_{t-r_l}(x, v_l) \left(D_{r_1, v_1}^{(k_1)} \cdots D_{r_{l-1}, v_{l-1}}^{(k_{l-1})} D_{r_{l+1}, v_{l+1}}^{(k_{l+1})} \cdots D_{r_n, v_n}^{(k_n)} (\sigma_{ik_l}(u(r_l, v_l))) \right) \\ & \quad + \sum_{j=1}^d \int_{r_1 \vee \cdots \vee r_n}^t \int_0^1 G_{t-\theta}(x, \eta) \prod_{l=1}^n D_{r_l, v_l}^{(k_l)} (\sigma_{ij}(u(\theta, \eta))) W^j(d\theta, d\eta) \\ & \quad + \int_{r_1 \vee \cdots \vee r_n}^t \int_0^1 G_{t-\theta}(x, \eta) \prod_{l=1}^n D_{r_l, v_l}^{(k_l)} (b_i(u(\theta, \eta))) d\theta d\eta \end{aligned}$$

if $t \leq r_1 \vee \cdots \vee r_n$ and $D_{r_1, v_1}^{(k_1)} \cdots D_{r_n, v_n}^{(k_n)} (u_i(t, x)) = 0$ otherwise. Finally, for any $p > 1$,

$$\sup_{(t,x) \in [0,T] \times [0,1]} \mathbb{E} \left[\left\| D^n(u_i(t, x)) \right\|_{\mathcal{H}^{\otimes n}}^p \right] < +\infty. \quad (4.1)$$

Note that, in particular, the first-order Malliavin derivative satisfies, for $r < t$,

$$D_{r,v}^{(k)}(u_i(t, x)) = G_{t-r}(x, v) \sigma_{ik}(u(r, v)) + a_i(k, r, v, t, x), \quad (4.2)$$

where

$$\begin{aligned} a_i(k, r, v, t, x) &= \sum_{j=1}^d \int_r^t \int_0^1 G_{t-\theta}(x, \eta) D_{r,v}^{(k)} (\sigma_{ij}(u(\theta, \eta))) W^j(d\theta, d\eta) \\ & \quad + \int_r^t \int_0^1 G_{t-\theta}(x, \eta) D_{r,v}^{(k)} (b_i(u(\theta, \eta))) d\theta d\eta, \end{aligned} \quad (4.3)$$

and $D_{r,v}^{(k)}(u_i(t, x)) = 0$ when $r > t$.

The next result proves property (a) in Proposition 3.4 when F is replaced by $u(t, x)$.

Proposition 4.2. *Assume P1 and P2. Let I and J two compact intervals as in Theorem 1.1. Then, for any $p \geq 1$,*

$$\mathbb{E}[(\det \gamma_{u(t,x)})^{-p}]$$

is uniformly bounded over $(t, x) \in I \times J$.

Proof. This proof follows NUALART [N98, Proof of (3.22)], where it is shown that for fixed (t, x) , $\mathbb{E}[(\det \gamma_{u(t,x)})^{-p}] < +\infty$. Our emphasis here is on the uniform bound over $(t, x) \in I \times J$. Assume that $I = [t_1, t_2]$ and $J = [x_1, x_2]$, where $0 < t_1 < t_2 \leq T$, $0 < x_1 < x_2 < 1$. Let $(t, x) \in I \times J$ be fixed. We write

$$\det \gamma_{u(t,x)} \geq \left(\inf_{\xi \in \mathbb{R}^d: \|\xi\|=1} \xi^T \gamma_{u(t,x)} \xi \right)^d.$$

Let $\xi = (\xi_1, \dots, \xi_d) \in \mathbb{R}^d$ with $\|\xi\| = 1$ and fix $\epsilon \in (0, 1)$. Note the inequality

$$(a + b)^2 \geq \frac{2}{3}a^2 - 2b^2, \quad (4.4)$$

valid for all $a, b \geq 0$. Using (4.2) and the fact that $\gamma_{u(t,x)}$ is a matrix whose entries are inner-products, this implies that

$$\begin{aligned} \xi^T \gamma_{u(t,x)} \xi &= \int_0^t dr \int_0^1 dv \left\| \sum_{i=1}^d D_{r,v}(u_i(t,x)) \xi_i \right\|^2 \\ &\geq \int_{t(1-\epsilon)}^t dr \int_0^1 dv \left\| \sum_{i=1}^d D_{r,v}(u_i(t,x)) \xi_i \right\|^2 \geq I_1 - I_2, \end{aligned}$$

where

$$\begin{aligned} I_1 &= \frac{2}{3} \int_{t(1-\epsilon)}^t dr \int_0^1 dv \sum_{k=1}^d \left(\sum_{i=1}^d G_{t-r}(x,v) \sigma_{ik}(u(r,v)) \xi_i \right)^2, \\ I_2 &= 2 \int_{t(1-\epsilon)}^t dr \int_0^1 dv \sum_{k=1}^d \left(\sum_{i=1}^d a_i(k,r,v,t,x) \xi_i \right)^2, \end{aligned}$$

and $a_i(k,r,v,t,x)$ is defined in (4.3). In accord with hypothesis **P2** and thanks to Lemma 7.2,

$$I_1 \geq c(t\epsilon)^{1/2}, \quad (4.5)$$

where c is uniform over $(t,x) \in I \times J$.

Next we apply the Cauchy-Schwarz inequality to find that, for any $q \geq 1$,

$$\mathbb{E} \left[\sup_{\xi \in \mathbb{R}^d: \|\xi\|=1} |I_2|^q \right] \leq c(\mathbb{E}[|A_1|^q] + \mathbb{E}[|A_2|^q]),$$

where

$$\begin{aligned} A_1 &= \sum_{i,j,k=1}^d \int_{t(1-\epsilon)}^t dr \int_0^1 dv \left(\int_r^t \int_0^1 G_{t-\theta}(x,\eta) D_{r,v}^{(k)}(\sigma_{ij}(u(\theta,\eta))) W^j(d\theta, d\eta) \right)^2, \\ A_2 &= \sum_{i,k=1}^d \int_{t(1-\epsilon)}^t dr \int_0^1 dv \left(\int_r^t \int_0^1 G_{t-\theta}(x,\eta) D_{r,v}^{(k)}(b_i(u(\theta,\eta))) d\theta d\eta \right)^2. \end{aligned}$$

We bound the q -th moment of A_1 and A_2 separately. As regards A_1 , we use Burkholder's inequality for martingales with values in a Hilbert space (Lemma 7.6) to obtain

$$\mathbb{E}[|A_1|^q] \leq c \sum_{k,i=1}^d \mathbb{E} \left[\left\| \int_{t(1-\epsilon)}^t d\theta \int_0^1 d\eta \int_{t(1-\epsilon)}^t dr \int_0^1 dv \Theta^2 \right\|^q \right], \quad (4.6)$$

where

$$\begin{aligned}\Theta &:= 1_{\{\theta>r\}} G_{t-\theta}(x, \eta) \left| D_{r,v}^{(k)}(\sigma_{ij}(u(\theta, \eta))) \right| \\ &\leq c 1_{\{\theta>r\}} G_{t-\theta}(x, \eta) \left| \sum_{l=1}^d D_{r,v}^{(k)}(u_l(\theta, \eta)) \right|,\end{aligned}$$

thanks to hypothesis **P1**. Hence,

$$\mathbb{E}[|A_1|^q] \leq c \sum_{k=1}^d \mathbb{E} \left[\left| \int_{t(1-\epsilon)}^t d\theta \int_0^1 d\eta G_{t-\theta}^2(x, \eta) \int_{t(1-\epsilon)}^{t \wedge \theta} dr \int_0^1 dv \Psi^2 \right|^q \right],$$

where $\Psi := \sum_{l=1}^d D_{r,v}^{(k)}(u_l(\theta, \eta))$. We now apply Hölder's inequality with respect to the measure $G_{t-\theta}^2(x, \eta) d\theta d\eta$ to find that

$$\begin{aligned}\mathbb{E}[|A_1|^q] &\leq C \left| \int_{t(1-\epsilon)}^t d\theta \int_0^1 d\eta G_{t-\theta}^2(x, \eta) \right|^{q-1} \\ &\quad \times \int_{t(1-\epsilon)}^t d\theta \int_0^1 d\eta G_{t-\theta}^2(x, \eta) \sum_{k=1}^d \mathbb{E} \left[\left| \int_{t(1-\epsilon)}^t dr \int_0^1 dv \Psi^2 \right|^q \right].\end{aligned}$$

Lemmas 7.3 and 7.5 assure that

$$\mathbb{E}[|A_1|^q] \leq C_T (t\epsilon)^{\frac{q-1}{2}} (t\epsilon)^{q/2} \int_{t(1-\epsilon)}^t \int_0^1 G_{t-\theta}^2(x, \eta) d\theta d\eta \leq C_T (t\epsilon)^q,$$

where C_T is uniform over $(t, x) \in I \times J$.

We next derive a similar bound for A_2 . By the Cauchy–Schwarz inequality,

$$\mathbb{E}[|A_2|^q] \leq c(t\epsilon)^q \sum_{i,k=1}^d \mathbb{E} \left[\left| \int_{t(1-\epsilon)}^t dr \int_0^1 dv \int_r^t d\theta \int_0^1 d\eta \Phi^2 \right|^q \right],$$

where $\Phi := G_{t-\theta}(x, \eta) |D_{r,v}^{(k)}(b_i(u(\theta, \eta)))|$. From here on, the q -th moment of A_2 is estimated as that of A_1 was; cf. (4.6), and this yields $\mathbb{E}[|A_2|^q] \leq C_T (t\epsilon)^{2q}$.

Thus, we have proved that

$$\mathbb{E} \left[\sup_{\xi \in \mathbb{R}^d: \|\xi\|=1} |I_2|^q \right] \leq C_T (t\epsilon)^q, \quad (4.7)$$

where the constant C_T is clearly uniform over $(t, x) \in I \times J$.

Finally, we apply Proposition 3.5 with $Z := \inf_{\|\xi\|=1} (\xi^T \gamma_{u(t,x)} \xi)$, $Y_{1,\epsilon} = Y_{2,\epsilon} = \sup_{\|\xi\|=1} I_2$, $\epsilon_0 = 1$, $\alpha_1 = \alpha_2 = 1/2$ and $\beta_1 = \beta_2 = 1$, to get

$$\mathbb{E} \left[(\det \gamma_{u(t,x)})^{-p} \right] \leq C_T,$$

where all the constants are clearly uniform over $(t, x) \in I \times J$. This is the desired result. \square

Corollary 4.3. *Assume **P1** and **P2**. Fix $T > 0$ and let I and J be compact intervals as in Theorem 1.1. Then, for any $(t, x) \in (0, T) \times (0, 1)$, $u(t, x)$ is a nondegenerate random vector and its density function is infinitely differentiable and uniformly bounded over $z \in \mathbb{R}^d$ and $(t, x) \in I \times J$.*

Proof of Theorem 1.1(a). This is a consequence of Propositions 4.1 and 4.2 together with Theorem 3.1 and Proposition 3.4. \square

5 The Gaussian-type lower bound on the one-point density

The aim of this section is to prove the lower bound of Gaussian-type for the density of u stated in Theorem 1.1(b). The proof of this result was given in KOHATSU-HIGA [K03, Theorem 10] for dimension 1, therefore we will only sketch the main steps.

Proof of Theorem 1.1(b). We follow [K03] and we show that for each (t, x) , $F = u(t, x)$ is a d -dimensional uniformly elliptic random vector and then we apply [K03, Theorem 5]. Let

$$F_n^i = \int_0^{t_n} \int_0^1 G_{t-r}(x, v) \sum_{j=1}^d \sigma_{ij}(u(r, v)) W^j(dr, dv) + \int_0^{t_n} \int_0^1 G_{t-r}(x, v) b_i(u(r, v)) dr dv,$$

$1 \leq i \leq d$, where $0 = t_0 < t_1 < \dots < t_N = t$ is a sufficiently fine partition of $[0, t]$. Note that $F_n \in \mathcal{F}_{t_n}$. Set $g(s, y) = G_{t-s}(x, y)$. We shall need the following two lemmas.

Lemma 5.1. [K03, Lemma 7] *Assume **P1** and **P2**. Then:*

- (i) $\|F_n^i\|_{k,p} \leq c_{k,p}$, $1 \leq i \leq d$;
- (ii) $\|(\gamma_{F_n}(t_{n-1}))_{ij}^{-1}\|_{p,t_{n-1}} \leq c_p (\Delta_{n-1}(g))^{-1} = c_p (\|g\|_{L^2([t_{n-1}, t_n] \times [0, 1])}^2)^{-1}$,

where $\gamma_{F_n}(t_{n-1})$ denotes the conditional Malliavin matrix of F_n given $\mathcal{F}_{t_{n-1}}$ and $\|\cdot\|_{p,t_{n-1}}$ denotes the conditional L^p -norm.

We define

$$\begin{aligned} u_i^{n-1}(s_1, y_1) &= \int_0^{t_{n-1}} \int_0^1 G_{s_1-s_2}(y_1, y_2) \sum_{j=1}^d \sigma_{ij}(u(s_2, y_2)) W^j(ds_2, dy_2) \\ &\quad + \int_0^{t_{n-1}} \int_0^1 G_{s_1-s_2}(y_1, y_2) b_i(u(s_2, y_2)) ds_2 dy_2, \quad 1 \leq i \leq d. \end{aligned}$$

Note that $u^{n-1} \in \mathcal{F}_{t_{n-1}}$. As in [K03], the following holds.

Lemma 5.2. [K03, Lemma 8] *Under hypothesis **P1**, for $s \in [t_{n-1}, t_n]$,*

$$\|u_i(s, y) - u_i^{n-1}(s, y)\|_{n,p,t_{n-1}} \leq (s - t_{n-1})^{1/8}, \quad 1 \leq i \leq d,$$

where $\|\cdot\|_{n,p,t_{n-1}}$ denotes the conditional Malliavin norm given $\mathcal{F}_{t_{n-1}}$.

The rest of the proof of Theorem 1.1(b) follows along the same lines as in [K03] for $d = 1$. We only sketch the remaining main points where the fact that $d > 1$ is important. In order to obtain the expansion of $F_n^i - F_{n-1}^i$ as in [K03, Lemma 9], we proceed as follows. By the mean value theorem,

$$\begin{aligned} F_n^i - F_{n-1}^i &= \int_{t_{n-1}}^{t_n} \int_0^1 G_{t-r}(x, v) \sum_{j=1}^d \sigma_{ij}(u^{n-1}(r, v)) W^j(dr, dv) \\ &\quad + \int_{t_{n-1}}^{t_n} \int_0^1 G_{t-r}(x, v) b_i(u(r, v)) dr dv \\ &\quad + \int_{t_{n-1}}^{t_n} \int_0^1 G_{t-r}(x, v) \sum_{j,l=1}^d \left(\int_0^1 \partial_l \sigma_{ij}(u(r, v, \lambda)) d\lambda \right) (u_l(r, v) - u_l^{n-1}(r, v)) W^j(dr, dv), \end{aligned}$$

where $u(r, v, \lambda) = (1 - \lambda)u(r, v) + \lambda u^{n-1}(r, v)$. Using the terminology of [K03], the first term is a process of order 1 and the next two terms are residues of order 1 (as in [K03]). In the next step, we write the residues of order 1 as the sum of processes of order 2 and residues of order 2 and 3 as follows:

$$\begin{aligned} &\int_{t_{n-1}}^{t_n} \int_0^1 G_{t-r}(x, v) b_i(u(r, v)) dr dv \\ &= \int_{t_{n-1}}^{t_n} \int_0^1 G_{t-r}(x, v) b_i(u^{n-1}(r, v)) dr dv \\ &\quad + \int_{t_{n-1}}^{t_n} \int_0^1 G_{t-r}(x, v) \sum_{l=1}^d \left(\int_0^1 \partial_l b_i(u(r, v, \lambda)) d\lambda \right) (u_l(r, v) - u_l^{n-1}(r, v)) dr dv \end{aligned}$$

and

$$\begin{aligned} &\int_{t_{n-1}}^{t_n} \int_0^1 G_{t-r}(x, v) \sum_{j,l=1}^d \left(\int_0^1 \partial_l \sigma_{ij}(u(r, v, \lambda)) d\lambda \right) (u_l(r, v) - u_l^{n-1}(r, v)) W^j(dr, dv) \\ &= \int_{t_{n-1}}^{t_n} \int_0^1 G_{t-r}(x, v) \sum_{j,l=1}^d \partial_l \sigma_{ij}(u^{n-1}(r, v)) (u_l(r, v) - u_l^{n-1}(r, v)) W^j(dr, dv) \\ &\quad + \int_{t_{n-1}}^{t_n} \int_0^1 G_{t-r}(x, v) \sum_{j,l,l'=1}^d \left(\int_0^1 \partial_l \partial_{l'} \sigma_{ij}(u(r, v, \lambda)) d\lambda \right) \\ &\quad \quad \times (u_l(r, v) - u_l^{n-1}(r, v)) (u_{l'}(r, v) - u_{l'}^{n-1}(r, v)) W^j(dr, dv). \end{aligned}$$

It is then clear that the remainder of the proof of [K03, Lemma 9] follows for $d > 1$ along the same lines as in [K03], working coordinate by coordinate.

Finally, in order to complete the proof of the proposition, it suffices to verify the hypotheses of [K03, Theorem 5]. Again the proof follows as in the proof of [K03, Theorem 10], working coordinate by coordinate. We will only sketch the proof of his (H2c), where

hypothesis **P2** is used:

$$\begin{aligned} & (\Delta_{n-1}(g))^{-1} \int_{t_{n-1}}^{t_n} \int_0^1 (G_{t-r}(x, v))^2 \|\sigma(u^{n-1}(r, v))\xi\|^2 dr dv \\ & \geq \rho^2 (\Delta_{n-1}(g))^{-1} \int_{t_{n-1}}^{t_n} \int_0^1 (G_{t-r}(x, v))^2 dr dv = \rho^2 > 0, \end{aligned}$$

by the definition of g . This concludes the proof of Theorem 1.1 (b). \square

6 The Gaussian-type upper bound on the two-point density

Let $p_{s,y;t,x}(z_1, z_2)$ denote the joint density of the $2d$ -dimensional random vector

$$(u_1(s, y), \dots, u_d(s, y), u_1(t, x), \dots, u_d(t, x)),$$

for $s, t \in (0, T]$, $x, y \in (0, 1)$, $(s, y) \neq (t, x)$ and $z_1, z_2 \in \mathbb{R}^d$ (the existence of this joint density will be a consequence of Theorem 3.1, Proposition 4.1 and Theorem 6.3).

The next subsections lead to the proofs of Theorem 1.1(c) and (d).

6.1 Bounds on the increments of the Malliavin derivatives

In this subsection, we prove an upper bound for the Sobolev norm of the derivative of the increments of our process u . For this, we will need the following preliminary estimate.

Lemma 6.1. *For any $s, t \in [0, T]$, $s \leq t$, and $x, y \in [0, 1]$,*

$$\int_0^T \int_0^1 (g(r, v))^2 dr dv \leq C_T (|t - s|^{1/2} + |x - y|),$$

where

$$g(r, v) := g_{t,x,s,y}(r, v) = 1_{\{r \leq t\}} G_{t-r}(x, v) - 1_{\{r \leq s\}} G_{s-r}(y, v).$$

Proof. Using BALLY, MILLET, AND SANZ-SOLÉ [BMS95, Lemma B.1] with $\alpha = 2$, we see that

$$\begin{aligned} & \int_0^T \int_0^1 (g(r, v))^2 dr dv \\ & \leq \int_s^t \int_0^1 (G_{t-r}(x, v))^2 dr dv + 2 \int_0^s \int_0^1 (G_{t-r}(x, v) - G_{s-r}(x, v))^2 dr dv \\ & \quad + 2 \int_0^s \int_0^1 (G_{s-r}(x, v) - G_{s-r}(y, v))^2 dr dv \\ & \leq C_T (|t - s|^{1/2} + |x - y|). \end{aligned}$$

\square

Proposition 6.2. *Assuming **P1**, for any $s, t \in [0, T]$, $s \leq t$, $x, y \in [0, 1]$, $p > 1$, $m \geq 1$,*

$$\mathbb{E} \left[\left\| D^m(u_i(t, x) - u_i(s, y)) \right\|_{\mathcal{H}^{\otimes m}}^p \right] \leq C_T (|t - s|^{1/2} + |x - y|)^{p/2}, \quad i = 1, \dots, d.$$

Proof. Let $m = 1$. Consider the function $g(r, v)$ defined in Lemma 6.1. Using the integral equation (4.2) satisfied by the first-order Malliavin derivative, we find that

$$\mathbb{E} \left[\left\| D(u_i(t, x) - u_i(s, y)) \right\|_{\mathcal{H}}^p \right] \leq C (\mathbb{E}[|I_1|^{p/2}] + \mathbb{E}[|I_2|^{p/2}] + \mathbb{E}[|I_3|^{p/2}]),$$

where

$$\begin{aligned} I_1 &= \sum_{k=1}^d \int_0^T dr \int_0^1 dv (g(r, v) \sigma_{ik}(u(r, v)))^2, \\ I_2 &= \sum_{j,k=1}^d \int_0^T dr \int_0^1 dv \left(\int_0^T \int_0^1 g(\theta, \eta) D_{r,v}^{(k)}(\sigma_{ij}(u(\theta, \eta))) W^j(d\theta, d\eta) \right)^2, \\ I_3 &= \sum_{k=1}^d \int_0^T dr \int_0^1 dv \left(\int_0^T \int_0^1 g(\theta, \eta) D_{r,v}^{(k)}(b_i(u(\theta, \eta))) d\theta d\eta \right)^2. \end{aligned}$$

We bound the $p/2$ -moments of I_1, I_2 and I_3 separately.

By hypothesis **P1** and Lemma 6.1, $\mathbb{E}[|I_1|^{p/2}] \leq C_T (|t - s|^{1/2} + |x - y|)^{p/2}$. Using Burkholder's inequality for Hilbert-space-valued martingales (Lemma 7.6) and hypothesis **P1**, we obtain

$$\mathbb{E}[|I_2|^{p/2}] \leq C \sum_{k=1}^d \mathbb{E} \left[\left| \int_0^T d\theta \int_0^1 d\eta (g(\theta, \eta))^2 \int_0^T dr \int_0^1 dv \Theta^2 \right|^{p/2} \right],$$

where $\Theta := \sum_{l=1}^d D_{r,v}^{(k)}(u_l(\theta, \eta))$. From Hölder's inequality with respect to the measure $(g(\theta, \eta))^2 d\theta d\eta$, we see that this is bounded above by

$$\begin{aligned} & C \left(\int_0^T \int_0^1 (g(\theta, \eta))^2 d\theta d\eta \right)^{\frac{p}{2}-1} \\ & \quad \times \sup_{(\theta, \eta) \in [0, T] \times [0, 1]} \sum_{k=1}^d \mathbb{E} \left[\left| \int_0^T d\theta \int_0^1 d\eta (g(\theta, \eta))^2 \int_0^T dr \int_0^1 dv \Theta^2 \right|^{p/2} \right] \\ & \leq C_T (|t - s|^{1/2} + |x - y|)^{p/2}, \end{aligned}$$

thanks to (4.1) and Lemma 6.1.

We next derive a similar bound for I_3 . By the Cauchy-Schwarz inequality,

$$\mathbb{E}[|I_3|^{p/2}] \leq C_T \sum_{k=1}^d \mathbb{E} \left[\left| \int_0^T d\theta \int_0^1 d\eta (g(\theta, \eta))^2 \int_0^T dr \int_0^1 dv \Theta^2 \right|^{p/2} \right].$$

From here on, the $p/2$ -moment of I_3 is estimated as was that of I_2 , and this yields $\mathbb{E}[|I_3|^{p/2}] \leq C_T(|t-s|^{1/2} + |x-y|)^{p/2}$. This proves the desired result for $m = 1$. The case $m > 1$ follows using the stochastic differential equation satisfied by the iterated Malliavin derivatives (Proposition 4.1), Hölder's and Burkholder's inequalities, hypothesis **P1**, (4.1) and Lemma 6.1 in the same way as we did for $m = 1$, to obtain the desired bound. \square

6.2 Study of the Malliavin matrix

For $s, t \in [0, T]$, $s \leq t$, and $x, y \in [0, 1]$ consider the $2d$ -dimensional random vector

$$Z := (u_1(s, y), \dots, u_d(s, y), u_1(t, x) - u_1(s, y), \dots, u_d(t, x) - u_d(s, y)). \quad (6.1)$$

Let γ_Z the Malliavin matrix of Z . Note that $\gamma_Z = ((\gamma_Z)_{m,l})_{m,l=1,\dots,2d}$ is a symmetric $2d \times 2d$ random matrix with four $d \times d$ blocs of the form

$$\gamma_Z = \begin{pmatrix} \gamma_Z^{(1)} & \vdots & \gamma_Z^{(2)} \\ \dots & \vdots & \dots \\ \gamma_Z^{(3)} & \vdots & \gamma_Z^{(4)} \end{pmatrix},$$

where

$$\begin{aligned} \gamma_Z^{(1)} &= (\langle D(u_i(s, y)), D(u_j(s, y)) \rangle_{\mathcal{H}})_{i,j=1,\dots,d}, \\ \gamma_Z^{(2)} &= (\langle D(u_i(s, y)), D(u_j(t, x) - u_j(s, y)) \rangle_{\mathcal{H}})_{i,j=1,\dots,d}, \\ \gamma_Z^{(3)} &= (\langle D(u_i(t, x) - u_i(s, y)), D(u_j(s, y)) \rangle_{\mathcal{H}})_{i,j=1,\dots,d}, \\ \gamma_Z^{(4)} &= (\langle D(u_i(t, x) - u_i(s, y)), D(u_j(t, x) - u_j(s, y)) \rangle_{\mathcal{H}})_{i,j=1,\dots,d}. \end{aligned}$$

We let **(1)** denote the set of indices $\{1, \dots, d\} \times \{1, \dots, d\}$, **(2)** the set $\{1, \dots, d\} \times \{d+1, \dots, 2d\}$, **(3)** the set $\{d+1, \dots, 2d\} \times \{1, \dots, d\}$ and **(4)** the set $\{d+1, \dots, 2d\} \times \{d+1, \dots, 2d\}$.

The following theorem gives an estimate on the Sobolev norm of the entries of the inverse of the matrix γ_Z , which depends on the position of the entry in the matrix.

Theorem 6.3. *Fix $\eta, T > 0$. Assume **P1** and **P2**. Let I and J be two compact intervals as in Theorem 1.1.*

(a) *For any $(s, y) \in I \times J$, $(t, x) \in I \times J$, $s \leq t$, $(s, y) \neq (t, x)$, $k \geq 0$, $p > 1$,*

$$\|(\gamma_Z^{-1})_{m,l}\|_{k,p} \leq \begin{cases} c_{k,p,\eta,T}(|t-s|^{1/2} + |x-y|)^{-d\eta} & \text{if } (m,l) \in \mathbf{(1)}, \\ c_{k,p,\eta,T}(|t-s|^{1/2} + |x-y|)^{-1/2-d\eta} & \text{if } (m,l) \in \mathbf{(2)} \text{ or } \mathbf{(3)}, \\ c_{k,p,\eta,T}(|t-s|^{1/2} + |x-y|)^{-1-d\eta} & \text{if } (m,l) \in \mathbf{(4)}. \end{cases}$$

(b) *For any $s = t \in (0, T]$, $(t, y) \in I \times J$, $(t, x) \in I \times J$, $x \neq y$, $k \geq 0$, $p > 1$,*

$$\|(\gamma_Z^{-1})_{m,l}\|_{k,p} \leq \begin{cases} c_{k,p,T} & \text{if } (m,l) \in \mathbf{(1)}, \\ c_{k,p,T}|x-y|^{-1/2} & \text{if } (m,l) \in \mathbf{(2)} \text{ or } \mathbf{(3)}, \\ c_{k,p,T}|x-y|^{-1} & \text{if } (m,l) \in \mathbf{(4)}. \end{cases}$$

(Note the slight improvements in the exponents in case (b) where $s = t$.)

The proof of this theorem is deferred to Section 6.4. We assume it for the moment and complete the proof of Theorem 1.1(c) and (d).

6.3 Proof of Theorem 1.1(c) and (d)

Fix two compact intervals I and J as in Theorem 1.1. Let $(s, y), (t, x) \in I \times J$, $s \leq t$, $(s, y) \neq (t, x)$, and $z_1, z_2 \in \mathbb{R}^d$. Let Z be as in (6.1) and let p_Z be the density of Z . Then

$$p_{s,y;t,x}(z_1, z_2) = p_Z(z_1, z_1 - z_2).$$

Apply Corollary 3.3 with $\sigma = \{i \in \{1, \dots, d\} : z_1^i - z_2^i \geq 0\}$ and Hölder's inequality to see that

$$p_Z(z_1, z_1 - z_2) \leq \prod_{i=1}^d \left(\mathbb{P} \left\{ |u_i(t, x) - u_i(s, y)| > |z_1^i - z_2^i| \right\} \right)^{\frac{1}{2d}} \times \|H_{(1, \dots, 2d)}(Z, 1)\|_{0,2}.$$

Therefore, in order to prove the desired results (c) and (d) of Theorem 1.1, it suffices to prove that:

$$\prod_{i=1}^d \left(\mathbb{P} \left\{ |u_i(t, x) - u_i(s, y)| > |z_1^i - z_2^i| \right\} \right)^{\frac{1}{2d}} \leq c \exp \left(-\frac{\|z_1 - z_2\|^2}{c_T(|t - s|^{1/2} + |x - y|)} \right), \quad (6.2)$$

$$\|H_{(1, \dots, 2d)}(Z, 1)\|_{0,2} \leq c_T(|t - s|^{1/2} + |x - y|)^{-(d+\eta)/2}, \quad (6.3)$$

and if $s = t$, then

$$\|H_{(1, \dots, 2d)}(Z, 1)\|_{0,2} \leq c_T|x - y|^{-d/2}. \quad (6.4)$$

Proof of (6.2). Let \tilde{u} denote the solution of (2.1) for $b \equiv 0$. Consider the continuous one-parameter martingale $(M_u = (M_u^1, \dots, M_u^d), 0 \leq u \leq t)$ defined by

$$M_u^i = \begin{cases} \int_0^u \int_0^1 (G_{t-r}(x, v) - G_{s-r}(y, v)) \sum_{j=1}^d \sigma_{ij}(\tilde{u}(r, v)) W^j(dr, dv) & \text{if } 0 \leq u \leq s, \\ \int_0^s \int_0^1 (G_{t-r}(x, v) - G_{s-r}(y, v)) \sum_{j=1}^d \sigma_{ij}(\tilde{u}(r, v)) W^j(dr, dv) \\ \quad + \int_s^u \int_0^1 G_{t-r}(x, v) \sum_{j=1}^d \sigma_{ij}(\tilde{u}(r, v)) W^j(dr, dv) & \text{if } s \leq u \leq t, \end{cases}$$

for all $i = 1, \dots, d$, with respect to the filtration $(\mathcal{F}_u, 0 \leq u \leq t)$. Notice that

$$M_0 = 0, \quad M_t = \tilde{u}(t, x) - \tilde{u}(s, y).$$

Moreover, by hypothesis **P1** and Lemma 6.1,

$$\begin{aligned}
\langle M^i \rangle_t &= \int_0^s \int_0^1 (G_{t-r}(x, v) - G_{s-r}(y, v))^2 \sum_{j=1}^d (\sigma_{ij}(\tilde{u}(r, v)))^2 dr dv \\
&\quad + \int_s^t \int_0^1 (G_{t-r}(x, v))^2 \sum_{j=1}^d (\sigma_{ij}(\tilde{u}(r, v)))^2 dr dv \\
&\leq C \int_0^T \int_0^1 (g(r, v))^2 dr dv \\
&\leq C_T (|t - s|^{1/2} + |x - y|).
\end{aligned}$$

By the exponential martingale inequality NUALART [N95, A.5],

$$\mathbb{P} \left\{ |\tilde{u}^i(t, x) - \tilde{u}^i(s, y)| > |z_1^i - z_2^i| \right\} \leq 2 \exp \left(- \frac{|z_1^i - z_2^i|^2}{C_T (|t - s|^{1/2} + |x - y|)} \right). \quad (6.5)$$

We will now treat the case $b \not\equiv 0$ using Girsanov's theorem. Consider the random variable

$$\begin{aligned}
L_t &= \exp \left(- \int_0^t \int_0^1 \sigma^{-1}(u(r, v)) b(u(r, v)) \cdot W(dr, dv) \right. \\
&\quad \left. - \frac{1}{2} \int_0^t \int_0^1 \|\sigma^{-1}(u(r, v)) b(u(r, v))\|^2 dr dv \right).
\end{aligned}$$

The following Girsanov's theorem holds.

Theorem 6.4. [N94, Prop.1.6] $E[L_t] = 1$, and if $\tilde{\mathbb{P}}$ denotes the probability measure on (Ω, \mathcal{F}) defined by

$$\frac{d\tilde{\mathbb{P}}}{d\mathbb{P}}(\omega) = L_t(\omega),$$

then $\tilde{W}(t, x) = W(t, x) + \int_0^t \int_0^x \sigma^{-1}(u(r, v)) b(u(r, v)) dr dv$ is a standard Brownian sheet under $\tilde{\mathbb{P}}$.

Consequently, the law of u under $\tilde{\mathbb{P}}$ coincides with the law of \tilde{u} under \mathbb{P} . Consider now the random variable

$$\begin{aligned}
J_t &= \exp \left(- \int_0^t \int_0^1 \sigma^{-1}(\tilde{u}(r, v)) b(\tilde{u}(r, v)) \cdot W(dr, dv) \right. \\
&\quad \left. + \frac{1}{2} \int_0^t \int_0^1 \|\sigma^{-1}(\tilde{u}(r, v)) b(\tilde{u}(r, v))\|^2 dr dv \right).
\end{aligned}$$

Then, by Theorem 6.4, the Cauchy-Schwarz inequality and (6.5),

$$\begin{aligned}
\mathbb{P}\left\{|u_i(t, x) - u_i(s, y)| > |z_1^i - z_2^i|\right\} &= \mathbb{E}_{\tilde{\mathbb{P}}}\left[1_{\{|u^i(t, x) - u^i(s, y)| > |z_1^i - z_2^i|\}} L_t^{-1}\right] \\
&= \mathbb{E}_{\mathbb{P}}\left[1_{\{|\tilde{u}^i(t, x) - \tilde{u}^i(s, y)| > |z_1^i - z_2^i|\}} J_t^{-1}\right] \\
&\leq \left(\mathbb{P}\left\{|\tilde{u}^i(t, x) - \tilde{u}^i(s, y)| > |z_1^i - z_2^i|\right\}\right)^{1/2} \left(\mathbb{E}_{\mathbb{P}}[J_t^{-2}]\right)^{1/2} \\
&\leq 2 \exp\left(-\frac{|z_1^i - z_2^i|^2}{C_T(|t - s|^{1/2} + |x - y|)}\right) \left(\mathbb{E}_{\mathbb{P}}[J_t^{-2}]\right)^{1/2}.
\end{aligned}$$

Now, hypothesis **P1** and **P2** give

$$\begin{aligned}
\mathbb{E}_{\mathbb{P}}[J_t^{-2}] &\leq \mathbb{E}_{\mathbb{P}}\left[\exp\left(\int_0^t \int_0^1 2\sigma^{-1}(\tilde{u}(r, v)) b(\tilde{u}(r, v)) \cdot W(dr, dv)\right.\right. \\
&\quad \left.\left. - \frac{1}{2} \int_0^t \int_0^1 4\|\sigma^{-1}(\tilde{u}(r, v)) b(\tilde{u}(r, v))\|^2 dr dv\right)\right. \\
&\quad \left.\times \exp\left(\int_0^t \int_0^1 \|\sigma^{-1}(\tilde{u}(r, v)) b(\tilde{u}(r, v))\|^2 dr dv\right)\right] \\
&\leq C,
\end{aligned}$$

since the second exponential is bounded and the first is an exponential martingale.

Therefore, we have proved that

$$\mathbb{P}\left\{|u_i(t, x) - u_i(s, y)| > |z_1^i - z_2^i|\right\} \leq C \exp\left(-\frac{|z_1^i - z_2^i|^2}{C_T(|t - s|^{1/2} + |x - y|)}\right),$$

from which we conclude that

$$\prod_{i=1}^d \left(\mathbb{P}\left\{|u_i(t, x) - u_i(s, y)| > |z_1^i - z_2^i|\right\}\right)^{\frac{1}{2d}} \leq C \exp\left(-\frac{\|z_1 - z_2\|^2}{C_T(|t - s|^{1/2} + |x - y|)}\right).$$

This proves (6.2). □

Proof of (6.3). As in (3.1), using the continuity of the Skorohod integral δ and Hölder's inequality for Malliavin norms, we obtain

$$\begin{aligned}
\|H_{(1, \dots, 2d)}(Z, 1)\|_{0,2} &\leq C \|H_{(1, \dots, 2d-1)}(Z, 1)\|_{1,4} \left(\sum_{j=1}^d \|(\gamma_Z^{-1})_{2d,j}\|_{1,8} \|D(Z^j)\|_{1,8}\right. \\
&\quad \left. + \sum_{j=d+1}^{2d} \|(\gamma_Z^{-1})_{2d,j}\|_{1,8} \|D(Z^j)\|_{1,8}\right).
\end{aligned}$$

Notice that the entries of γ_Z^{-1} that appear in this expression belong to sets **(3)** and **(4)** of indices, as defined before Theorem 6.3. From Theorem 6.3(a) and Propositions 4.1 and 6.2, we find that this is bounded above by

$$C_T \|H_{(1,\dots,2d-1)}(Z, 1)\|_{1,4} \left(\sum_{j=1}^d (|t-s|^{1/2} + |x-y|)^{-\frac{1}{2}-d\eta} + \sum_{j=d+1}^{2d} (|t-s|^{1/2} + |x-y|)^{-1-d\eta+\frac{1}{2}} \right),$$

that is, by

$$C_T \|H_{(1,\dots,2d-1)}(Z, 1)\|_{1,4} (|t-s|^{1/2} + |x-y|)^{-1/2-d\eta}.$$

Iterating this procedure d times (during which we only encounter coefficients $(\gamma_Z^{-1})_{m,l}$ for (m,l) in blocs **(3)** and **(4)**, cf. Theorem 6.3(a)), we get, for some integers $m_0, k_0 > 0$,

$$\|H_{(1,\dots,2d)}(Z, 1)\|_{0,2} \leq C_T \|H_{(1,\dots,d)}(Z, 1)\|_{m_0,k_0} (|t-s|^{1/2} + |x-y|)^{-d/2-d^2\eta}.$$

Again, using the continuity of δ and Hölder's inequality for the Malliavin norms, we obtain

$$\begin{aligned} \|H_{(1,\dots,d)}(Z, 1)\|_{m,k} \leq C \|H_{(1,\dots,d-1)}(Z, 1)\|_{m_1,k_1} & \left(\sum_{j=1}^d \|(\gamma_Z^{-1})_{d,j}\|_{m_2,k_2} \|D(Z^j)\|_{m_3,k_3} \right. \\ & \left. + \sum_{j=d+1}^{2d} \|(\gamma_Z^{-1})_{d,j}\|_{m_4,k_4} \|D(Z^j)\|_{m_5,k_5} \right), \end{aligned}$$

for some integers $m_i, k_i > 0$, $i = 1, \dots, 5$. This time, the entries of γ_Z^{-1} that appear in this expression come from the sets **(1)** and **(2)** of indices. We appeal again to Theorem 6.3(a) and Propositions 4.1 and 6.2 to get

$$\|H_{(1,\dots,d)}(Z, 1)\|_{m,k} \leq C_T \|H_{(1,\dots,d-1)}(Z, 1)\|_{m_1,k_1} (|t-s|^{1/2} + |x-y|)^{-d\eta}.$$

Finally, iterating this procedure d times (during which we encounter coefficients $(\gamma_Z^{-1})_{m,l}$ for (m,l) in blocs **(1)** and **(2)** only, cf. Theorem 6.3(a)), and choosing $\eta' = 4d^2\eta$, we conclude that

$$\|H_{(1,\dots,2d)}(Z, 1)\|_{0,2} \leq C_T (|t-s|^{1/2} + |x-y|)^{-(d+\eta')/2},$$

which proves (6.3) and concludes the proof of Theorem 1.1(c). \square

Proof of (6.4). In order to prove (6.4), we proceed exactly along the same lines as in the proof of (6.3) but we appeal to Theorem 6.3(b). This concludes the proof of Theorem 1.1(d). \square

6.4 Proof of Theorem 6.3

Let Z as in (6.1). Since the inverse of the matrix γ_Z is the inverse of its determinant multiplied by its cofactor matrix, we examine these two factors separately.

Proposition 6.5. Fix $T > 0$ and let I and J be compact intervals as in Theorem 1.1. Assuming **P1**, for any $(s, y), (t, x) \in I \times J$, $(s, y) \neq (t, x)$, $p > 1$,

$$\mathbb{E}[|(A_Z)_{m,l}|^p]^{1/p} \leq \begin{cases} c_{p,T}(|t-s|^{1/2} + |x-y|)^d & \text{if } (m,l) \in \mathbf{(1)}, \\ c_{p,T}(|t-s|^{1/2} + |x-y|)^{d-\frac{1}{2}} & \text{if } (m,l) \in \mathbf{(2)} \text{ or } \mathbf{(3)}, \\ c_{p,T}(|t-s|^{1/2} + |x-y|)^{d-1} & \text{if } (m,l) \in \mathbf{(4)}, \end{cases}$$

where A_Z denotes the cofactor matrix of γ_Z .

Proof. We consider the four different cases.

- If $(m, l) \in \mathbf{(1)}$, we claim that

$$\begin{aligned} |(A_Z)_{m,l}| \leq C \sum_{k=0}^{d-1} \left\{ & \|D(u(s, y))\|_{\mathcal{H}}^{2k} \times \|D(u(t, x) - u(s, y))\|_{\mathcal{H}}^{2(d-1-k)} \right. \\ & \left. \times \|D(u(s, y))\|_{\mathcal{H}}^{2(d-1-k)} \times \|D(u(t, x) - u(s, y))\|_{\mathcal{H}}^{2(k+1)} \right\}. \end{aligned} \quad (6.6)$$

Indeed, let $A_Z^{m,l} = (a_{\bar{m}, \bar{l}}^{m,l})_{\bar{m}, \bar{l}=1, \dots, 2d-1}$ be the $(2d-1) \times (2d-1)$ -matrix obtained by removing from γ_Z its row m and column l . Then

$$(A_Z)_{m,l} = \det((A_Z)^{m,l}) = \sum_{\pi \text{ permutation of } (1, \dots, 2d-1)} a_{1, \pi(1)}^{m,l} \cdots a_{2d-1, \pi(2d-1)}^{m,l}.$$

Each term of this sum contains one entry from each row and column of $A_Z^{m,l}$. If there are k entries taken from bloc **(1)** of γ_Z , these occupy k rows and columns of $A_Z^{m,l}$. Therefore, $d-1-k$ entries must come from the $d-1$ remaining rows of bloc **(2)**, and the same number from the columns of bloc **(3)**. Finally, there remain $k+1$ entries to be taken from bloc **(4)**. Therefore,

$$\begin{aligned} |(A_Z)_{m,l}| \leq C \sum_{k=0}^{d-1} \sum \left\{ & (\text{product of } k \text{ entries from } \mathbf{(1)}) \right. \\ & \times (\text{product of } d-1-k \text{ entries from } \mathbf{(2)}) \\ & \left. \times (\text{product of } d-1-k \text{ entries from } \mathbf{(3)}) \times (\text{product of } k+1 \text{ entries from } \mathbf{(4)}) \right\}. \end{aligned}$$

Adding all the terms and using the particular form of these terms establishes (6.6).

Regrouping the various factors in (6.6), applying the Cauchy-Schwarz inequality and using (4.1) and Proposition 6.2, we obtain

$$\begin{aligned} \mathbb{E} \left[|(A_Z)_{m,l}|^p \right] & \leq C \sum_{k=0}^d \mathbb{E} \left[\|D(u(s, y))\|_{\mathcal{H}}^{2(d-1)p} \times \|D(u(t, x) - u(s, y))\|_{\mathcal{H}}^{2dp} \right] \\ & \leq C_T (|t-s|^{1/2} + |x-y|)^{dp}. \end{aligned}$$

- If $(m, l) \in \mathbf{(2)}$ or $(m, l) \in \mathbf{(3)}$, then using the same arguments as above, we obtain

$$\begin{aligned}
& |(A_Z)_{m,l}| \\
& \leq C \sum_{k=0}^{d-1} \sum \left\{ \|D(u(s, y))\|_{\mathcal{H}}^{2(d-1-k)} \times \|D(u(s, y))\|_{\mathcal{H}}^k \times \|D(u(t, x) - u(s, y))\|_{\mathcal{H}}^k \right. \\
& \quad \left. \times \|D(u(s, y))\|_{\mathcal{H}}^{k+1} \times \|D(u(t, x) - u(s, y))\|_{\mathcal{H}}^{k+1} \times \|D(u(t, x) - u(s, y))\|_{\mathcal{H}}^{2(d-1-k)} \right\} \\
& \leq C \sum_{k=0}^{d-1} \left\{ \|D(u(s, y))\|_{\mathcal{H}}^{2d-1} \times \|D(u(t, x) - u(s, y))\|_{\mathcal{H}}^{2d-1} \right\},
\end{aligned}$$

from which we conclude, using (4.1) and Proposition 6.2, that

$$\mathbb{E} \left[|(A_Z)_{m,l}|^p \right] \leq C_T (|t - s|^{1/2} + |x - y|)^{(d-\frac{1}{2})p}.$$

- If $(i, j) \in \mathbf{(4)}$, we obtain

$$\begin{aligned}
|(A_Z)_{m,l}| & \leq C \sum_{k=0}^{d-1} \sum \left\{ \|D(u(s, y))\|_{\mathcal{H}}^{2(k+1)} \times \|D(u(s, y))\|_{\mathcal{H}}^{2(d-1-k)} \right. \\
& \quad \left. \times \|D(u(t, x) - D(u(s, y)))\|_{\mathcal{H}}^{2(d-1-k)} \times \|D(u(t, x) - D(u(s, y)))\|_{\mathcal{H}}^{2k} \right\} \\
& \leq C \sum_{k=0}^{d-1} \left\{ \|D(u(s, y))\|_{\mathcal{H}}^{2d} \times \|D(u(t, x) - u(s, y))\|_{\mathcal{H}}^{2d-2} \right\},
\end{aligned}$$

from which we conclude that

$$\mathbb{E} \left[|(A_Z)_{m,l}|^p \right] \leq C_T (|t - s|^{1/2} + |x - y|)^{(d-1)p}.$$

This concludes the proof of the proposition. \square

Proposition 6.6. *Fix $\eta, T > 0$. Assume **P1** and **P2**. Let I and J be compact intervals as in Theorem 1.1.*

- (a) *There exists C depending on T and η such that for any $(s, y), (t, x) \in I \times J$, $(s, y) \neq (t, x)$, $p > 1$,*

$$\mathbb{E} \left[(\det \gamma_Z)^{-p} \right]^{1/p} \leq C (|t - s|^{1/2} + |x - y|)^{-d(1+\eta)}. \quad (6.7)$$

- (b) *There exists C only depending on T such that for any $s = t \in I$, $x, y \in J$, $x \neq y$, $p > 1$,*

$$\mathbb{E} \left[(\det \gamma_Z)^{-p} \right]^{1/p} \leq C (|x - y|)^{-d}.$$

Assuming this proposition, we will be able to conclude the proof of Theorem 6.3, after establishing the following estimate on the derivative of the Malliavin matrix.

Proposition 6.7. Fix $T > 0$. Let I and J be compact intervals as in Theorem 1.1. Assuming **P1**, for any $(s, y), (t, x) \in I \times J$, $(s, y) \neq (t, x)$, $p > 1$ and $k \geq 1$,

$$\mathbb{E}[\|D^k(\gamma_Z)_{m,l}\|_{\mathcal{H}^{\otimes k}}^p]^{1/p} \leq \begin{cases} c_{k,p,T} & \text{if } (m,l) \in \mathbf{(1)}, \\ c_{k,p,T}(|t-s|^{1/2} + |x-y|)^{1/2} & \text{if } (m,l) \in \mathbf{(2)} \text{ or } \mathbf{(3)}, \\ c_{k,p,T}(|t-s|^{1/2} + |x-y|) & \text{if } (m,l) \in \mathbf{(4)}. \end{cases}$$

Proof. We consider the four different blocs.

- If $(m, l) \in \mathbf{(4)}$, proceeding as in DALANG AND NUALART [DN04, p.2131] and appealing to Proposition 6.2 twice, we obtain

$$\begin{aligned} & \mathbb{E}[\|D^k(\gamma_Z)_{m,l}\|_{\mathcal{H}^{\otimes k}}^p] \\ &= \mathbb{E}[\|D^k(\int_0^T dr \int_0^1 dv D_{r,v}(u_m(t,x) - u_m(s,y)) \cdot D_{r,v}(u_l(t,x) - u_l(s,y)))\|_{\mathcal{H}^{\otimes k}}^p] \\ &\leq (k+1)^{p-1} \sum_{j=0}^k \binom{k}{j}^p \mathbb{E}[\|\int_0^T dr \int_0^1 dv D^j D_{r,v}(u_m(t,x) - u_m(s,y)) \\ &\quad \cdot D^{k-j} D_{r,v}(u_l(t,x) - u_l(s,y))\|_{\mathcal{H}^{\otimes k}}^p] \\ &\leq \tilde{C}_T (k+1)^{p-1} \sum_{j=0}^k \binom{k}{j}^p \{(\mathbb{E}[\|D^j D(u_m(t,x) - u_m(s,y))\|_{\mathcal{H}^{\otimes(j+1)}}^{2p}])^{1/2} \\ &\quad \times (\mathbb{E}[\|D^{k-j} D(u_l(t,x) - u_l(s,y))\|_{\mathcal{H}^{\otimes(k-j+1)}}^{2p}])^{1/2}\} \\ &\leq C_T (|t-s|^{1/2} + |x-y|)^p. \end{aligned}$$

- If $(m, l) \in \mathbf{(2)}$ or $(m, l) \in \mathbf{(3)}$, proceeding as above and appealing to (4.1) and Proposition 6.2, we get

$$\begin{aligned} & \mathbb{E}[\|D^k(\gamma_Z)_{m,l}\|_{\mathcal{H}^{\otimes k}}^p] \\ &\leq \tilde{C}_T (k+1)^{p-1} \sum_{j=0}^k \binom{k}{j}^p \{(\mathbb{E}[\|D^j D(u_m(t,x) - u_m(s,y))\|_{\mathcal{H}^{\otimes(j+1)}}^{2p}])^{1/2} \\ &\quad \times (\mathbb{E}[\|D^{k-j} D(u_l(s,y))\|_{\mathcal{H}^{\otimes(k-j+1)}}^{2p}])^{1/2}\} \\ &\leq C_T (|t-s|^{1/2} + |x-y|)^{p/2}. \end{aligned}$$

- If $(m, l) \in \mathbf{(1)}$, using (4.1), we obtain

$$\begin{aligned} & \mathbb{E}[\|D^k(\gamma_Z)_{m,l}\|_{\mathcal{H}^{\otimes k}}^p] \\ &\leq \tilde{C}_T (k+1)^{p-1} \sum_{j=0}^k \binom{k}{j}^p \{(\mathbb{E}[\|D^j D(u_m(s,y))\|_{\mathcal{H}^{\otimes(j+1)}}^{2p}])^{1/2} \\ &\quad \times (\mathbb{E}[\|D^{k-j} D(u_l(s,y))\|_{\mathcal{H}^{\otimes(k-j+1)}}^{2p}])^{1/2}\} \\ &\leq C_T. \end{aligned}$$

□

Proof of Theorem 6.3. When $k = 0$, the result follows directly using the fact that the inverse of a matrix is the inverse of its determinant multiplied by its cofactor matrix and the estimates of Propositions 6.5 and 6.6.

For $k \geq 1$, we shall establish the following two properties.

(a) For any $(s, y), (t, x) \in I \times J$, $(s, y) \neq (t, x)$, $s \leq t$, $k \geq 1$ and $p > 1$,

$$\mathbb{E}[\|D^k(\gamma_Z^{-1})_{m,l}\|_{\mathcal{H}^{\otimes k}}^p]^{1/p} \leq \begin{cases} c_{k,p,\eta,T}(|t-s|^{1/2} + |x-y|)^{-d\eta} & \text{if } (m,l) \in \mathbf{(1)}, \\ c_{k,p,\eta,T}(|t-s|^{1/2} + |x-y|)^{-1/2-d\eta} & \text{if } (m,l) \in \mathbf{(2)}, \mathbf{(3)}, \\ c_{k,p,\eta,T}(|t-s|^{1/2} + |x-y|)^{-1-d\eta} & \text{if } (m,l) \in \mathbf{(4)}. \end{cases}$$

(b) For any $s = t \in I$, $x, y \in J$, $x \neq y$, $k \geq 1$ and $p > 1$,

$$\mathbb{E}[\|D^k(\gamma_Z^{-1})_{m,l}\|_{\mathcal{H}^{\otimes k}}^p]^{1/p} \leq \begin{cases} c_{k,p,T} & \text{if } (m,l) \in \mathbf{(1)}, \\ c_{k,p,T}|x-y|^{-1/2} & \text{if } (m,l) \in \mathbf{(2)} \text{ or } \mathbf{(3)}, \\ c_{k,p,T}|x-y|^{-1} & \text{if } (m,l) \in \mathbf{(4)}. \end{cases}$$

Since

$$\|(\gamma_Z^{-1})_{m,l}\|_{k,p} = \left\{ \mathbb{E}[|(\gamma_Z^{-1})_{m,l}|^p] + \sum_{j=1}^k \mathbb{E}[\|D^j(\gamma_Z^{-1})_{m,l}\|_{\mathcal{H}^{\otimes j}}^p] \right\}^{1/p},$$

(a) and (b) prove the theorem.

We now prove (a) and (b). When $k = 1$, we will use (3.3) written as a matrix product:

$$D(\gamma_Z^{-1}) = \gamma_Z^{-1} D(\gamma_Z) \gamma_Z^{-1}. \quad (6.8)$$

Writing (6.8) in bloc product matrix notation with blocs **(1)**, **(2)**, **(3)** and **(4)**, we get that

$$\begin{aligned} D((\gamma_Z^{-1})^{(1)}) &= (\gamma_Z^{-1})^{(1)} D(\gamma_Z^{(1)}) (\gamma_Z^{-1})^{(1)} + (\gamma_Z^{-1})^{(1)} D(\gamma_Z^{(2)}) (\gamma_Z^{-1})^{(3)} \\ &\quad + (\gamma_Z^{-1})^{(2)} D(\gamma_Z^{(3)}) (\gamma_Z^{-1})^{(1)} + (\gamma_Z^{-1})^{(2)} D(\gamma_Z^{(4)}) (\gamma_Z^{-1})^{(3)}, \\ D((\gamma_Z^{-1})^{(2)}) &= (\gamma_Z^{-1})^{(1)} D(\gamma_Z^{(1)}) (\gamma_Z^{-1})^{(2)} + (\gamma_Z^{-1})^{(1)} D(\gamma_Z^{(2)}) (\gamma_Z^{-1})^{(4)} \\ &\quad + (\gamma_Z^{-1})^{(2)} D(\gamma_Z^{(3)}) (\gamma_Z^{-1})^{(2)} + (\gamma_Z^{-1})^{(2)} D(\gamma_Z^{(4)}) (\gamma_Z^{-1})^{(4)}, \\ D((\gamma_Z^{-1})^{(3)}) &= (\gamma_Z^{-1})^{(3)} D(\gamma_Z^{(1)}) (\gamma_Z^{-1})^{(1)} + (\gamma_Z^{-1})^{(3)} D(\gamma_Z^{(2)}) (\gamma_Z^{-1})^{(3)} \\ &\quad + (\gamma_Z^{-1})^{(4)} D(\gamma_Z^{(3)}) (\gamma_Z^{-1})^{(1)} + (\gamma_Z^{-1})^{(4)} D(\gamma_Z^{(4)}) (\gamma_Z^{-1})^{(3)}, \\ D((\gamma_Z^{-1})^{(4)}) &= (\gamma_Z^{-1})^{(3)} D(\gamma_Z^{(1)}) (\gamma_Z^{-1})^{(2)} + (\gamma_Z^{-1})^{(3)} D(\gamma_Z^{(2)}) (\gamma_Z^{-1})^{(4)} \\ &\quad + (\gamma_Z^{-1})^{(4)} D(\gamma_Z^{(3)}) (\gamma_Z^{-1})^{(2)} + (\gamma_Z^{-1})^{(4)} D(\gamma_Z^{(4)}) (\gamma_Z^{-1})^{(4)}. \end{aligned}$$

It now suffices to apply Hölder's inequality to each block and use the estimates of the case $k = 0$ and Proposition 6.7 to obtain the desired result for $k = 1$. For instance, for

$(m, l) \in (\mathbf{1})$,

$$\begin{aligned}
& \mathbb{E} \left[\left\| ((\gamma_Z^{-1})^{(2)}) D(\gamma_Z^{(4)}) (\gamma_Z^{-1})^{(3)} \right\|_{m,l}^p \right]^{1/p} \\
& \leq \sup_{m_1, l_1} \mathbb{E} \left[\left\| ((\gamma_Z^{-1})^{(2)}) \right\|_{m_1, l_1}^{2p} \right]^{1/(2p)} \sup_{m_2, l_2} \mathbb{E} \left[\left\| (D(\gamma_Z^{(4)})) \right\|_{m_2, l_2}^{4p} \right]^{1/(4p)} \\
& \quad \times \sup_{m_3, l_3} \mathbb{E} \left[\left\| ((\gamma_Z^{-1})^{(3)}) \right\|_{m_3, l_3}^{4p} \right]^{1/(4p)} \\
& \leq c (|t - s|^{1/2} + |x - y|)^{-\frac{1}{2} - d\eta + 1 - \frac{1}{2} - d\eta} = c (|t - s|^{1/2} + |x - y|)^{-2d\eta}.
\end{aligned}$$

For $k \geq 1$, in order to calculate $D^{k+1}(\gamma_Z^{(\cdot)})$, we will need to compute $D^k(\gamma_Z^{-1} D(\gamma_Z) \gamma_Z^{-1})$. For bloc numbers $i_1, i_2, i_3 \in \{1, 2, 3, 4\}$ and $k \geq 1$, we have

$$\begin{aligned}
& D^k((\gamma_Z^{-1})^{(i_1)}) D(\gamma_Z^{(i_2)}) (\gamma_Z^{-1})^{(i_3)} \\
& = \sum_{\substack{j_1 + j_2 + j_3 = k \\ j_i \in \{0, \dots, k\}}} \binom{k}{j_1 j_2 j_3} D^{j_1}((\gamma_Z^{-1})^{(i_1)}) D^{j_2}(D(\gamma_Z^{(i_2)})) D^{j_3}((\gamma_Z^{-1})^{(i_3)}).
\end{aligned}$$

Note that by Proposition 6.7, the norms of the derivatives $D^{j_2}(D(\gamma_Z^{(i_2)}))$ of $\gamma_Z^{(i_2)}$ are of the same order for all j_2 . Hence, we appeal again to Hölder's inequality and Proposition 6.7, and use a recursive argument in order to obtain the desired bounds. \square

Proof of Proposition 6.6. The main idea for the proof of Proposition 6.6 is to use a perturbation argument. Indeed, for (t, x) close to (s, y) , the matrix γ_Z is close to

$$\hat{\gamma} = \begin{pmatrix} \gamma_Z^{(1)} & \vdots & 0 \\ \dots & \vdots & \dots \\ 0 & \vdots & 0 \end{pmatrix}.$$

The matrix $\hat{\gamma}$ has d eigenvectors of the form $(\hat{\lambda}^1, \mathbf{0}), \dots, (\hat{\lambda}^d, \mathbf{0})$, where $\hat{\lambda}^1, \dots, \hat{\lambda}^d \in \mathbb{R}^d$ are eigenvectors of $\gamma_Z^{(1)} = \gamma_{u(s,y)}$, and $\mathbf{0} = (0, \dots, 0) \in \mathbb{R}^d$, and d other eigenvectors of the form $(\mathbf{0}, e^i)$ where e^1, \dots, e^d is a basis of \mathbb{R}^d . These last eigenvectors of $\hat{\gamma}$ are associated with the eigenvalue 0, while the former are associated with eigenvalues of order 1, as can be seen in the proof of Proposition 4.2.

We now write

$$\det \gamma_Z = \prod_{i=1}^{2d} (\xi^i)^T \gamma_Z \xi^i, \tag{6.9}$$

where $\xi = \{\xi^1, \dots, \xi^{2d}\}$ is an orthonormal basis of \mathbb{R}^{2d} consisting of eigenvectors of γ_Z . We then expect that for (t, x) close to (s, y) , there will be d eigenvectors close to the subspace

generated by the $(\tilde{\lambda}^i, \mathbf{0})$, which will contribute a factor of order 1 to the product in (6.9), and d other eigenvectors, close to the subspace generated by the $(\mathbf{0}, e^i)$, that will each contribute a factor of order $(|t-s|^{1/2} + |x-y|)^{-1-\eta}$ to the product. Note that if we do not distinguish between these two types of eigenvectors, but simply bound below the product by the smallest eigenvalue to the power $2d$, following the approach used in the proof of Proposition 4.2, then we would obtain $C(|t-s|^{1/2} + |x-y|)^{-2dp}$ in the right-hand side of (6.7), which would not be the correct order.

We now carry out this somewhat involved perturbation argument. Consider the spaces $E_1 = \{(\lambda, \mathbf{0}) : \lambda \in \mathbb{R}^d, \mathbf{0} \in \mathbb{R}^d\}$ and $E_2 = \{(\mathbf{0}, \mu) : \mu \in \mathbb{R}^d, \mathbf{0} \in \mathbb{R}^d\}$. Note that every ξ^i can be written as

$$\xi^i = (\lambda_i, \mu_i) = \alpha_i(\tilde{\lambda}^i, \mathbf{0}) + \sqrt{1 - \alpha_i^2}(\mathbf{0}, \tilde{\mu}^i), \quad (6.10)$$

where $\lambda_i, \mu_i \in \mathbb{R}^d$, $(\tilde{\lambda}^i, \mathbf{0}) \in E_1$, $(\mathbf{0}, \tilde{\mu}^i) \in E_2$, with $\|\tilde{\lambda}^i\| = \|\tilde{\mu}^i\| = 1$ and $0 \leq \alpha_i \leq 1$. Note in particular that $\|\xi^i\|^2 = \|\lambda_i\|^2 + \|\mu_i\|^2 = 1$ (norms of elements of \mathbb{R}^d or \mathbb{R}^{2d} are Euclidean norms).

Lemma 6.8. *Given a sufficiently small $\alpha_0 > 0$, with probability one, there exist at least d of these vectors, say ξ^1, \dots, ξ^d , such that $\alpha_1 \geq \alpha_0, \dots, \alpha_d \geq \alpha_0$.*

Proof. Observe that as ξ is an orthogonal family and for $i \neq j$, the Euclidean inner product of ξ^i and ξ^j is

$$\xi^i \cdot \xi^j = \alpha_i \alpha_j (\tilde{\lambda}^i \cdot \tilde{\lambda}^j) + \sqrt{1 - \alpha_i^2} \sqrt{1 - \alpha_j^2} (\tilde{\mu}^i \cdot \tilde{\mu}^j) = 0.$$

For $\alpha_0 > 0$, let $D = \{i \in \{1, \dots, 2d\} : \alpha_i < \alpha_0\}$. Then, for $i, j \in D$, $i \neq j$, if $\alpha_0 < \frac{1}{2}$, then

$$|\tilde{\mu}^i \cdot \tilde{\mu}^j| = \frac{\alpha_i \alpha_j}{\sqrt{1 - \alpha_i^2} \sqrt{1 - \alpha_j^2}} |\tilde{\lambda}^i \cdot \tilde{\lambda}^j| \leq \frac{\alpha_0^2}{1 - \alpha_0^2} \|\tilde{\lambda}^i\| \|\tilde{\lambda}^j\| \leq \frac{1}{3} \alpha_0^2.$$

Since the diagonal terms of the matrix $(\tilde{\mu}^i \cdot \tilde{\mu}^j)_{i,j \in D}$ are all equal to 1, for α_0 sufficiently small, it follows that $\det((\tilde{\mu}^i \cdot \tilde{\mu}^j)_{i,j \in D}) \neq 0$. Therefore, $\{\tilde{\mu}^i, i \in D\}$ is a linearly independent family, and, as $(\mathbf{0}, \tilde{\mu}^i) \in E_2$, for $i = 1, \dots, 2d$, we conclude that a.s., $\text{card}(D) \leq \dim(E_2) = d$. We can therefore assume that $\{1, \dots, d\} \subset D^c$ and so $\alpha_1 \geq \alpha_0, \dots, \alpha_d \geq \alpha_0$. \square

By Lemma 6.8 and Cauchy-Schwarz inequality one can write

$$\begin{aligned} \mathbb{E}[(\det \gamma_Z)^{-p}]^{1/p} &\leq \left(\mathbb{E} \left[\left(\prod_{i=1}^d (\xi^i)^T \gamma_Z \xi^i \right)^{-2p} \right] \right)^{1/(2p)} \\ &\quad \times \left(\mathbb{E} \left[\left(\inf_{\substack{\xi = (\lambda, \mu) \in \mathbb{R}^{2d} \\ \|\lambda\|^2 + \|\mu\|^2 = 1}} \xi^T \gamma_Z \xi \right)^{-2dp} \right] \right)^{1/(2p)}. \end{aligned}$$

With this, Propositions 6.9 and 6.15 below conclude the proof of Proposition 6.6. \square

6.4.1 Small Eigenvalues

Let I and J two compact intervals as in Theorem 1.1.

Proposition 6.9. *Fix $\eta, T > 0$. Assume **P1** and **P2**.*

- (a) *There exists C depending on η and T such that for all $s, t \in I$, $0 < t - s < 1$, $x, y \in J$, $x \neq y$, and $p > 1$,*

$$\mathbb{E} \left[\left(\inf_{\substack{\xi = (\lambda, \mu) \in \mathbb{R}^{2d}: \\ \|\lambda\|^2 + \|\mu\|^2 = 1}} \xi^\top \gamma_Z \xi \right)^{-2dp} \right] \leq C(|t - s|^{1/2} + |x - y|)^{-2dp(1+\eta)}.$$

- (b) *There exists C depending only on T such that for all $s = t \in I$, $x, y \in J$, $x \neq y$, and $p > 1$,*

$$\mathbb{E} \left[\left(\inf_{\substack{\xi = (\lambda, \mu) \in \mathbb{R}^{2d}: \\ \|\lambda\|^2 + \|\mu\|^2 = 1}} \xi^\top \gamma_Z \xi \right)^{-2dp} \right] \leq C(|x - y|)^{-2dp}.$$

Proof. We begin by proving (a). Since γ_Z is a matrix of inner products, we can write

$$\xi^\top \gamma_Z \xi = \sum_{k=1}^d \int_0^T dr \int_0^1 dv \left(\sum_{i=1}^d \left(\lambda_i D_{r,v}^{(k)}(u(s, y)) + \mu_i (D_{r,v}^{(k)}(u(t, x)) - D_{r,v}^{(k)}(u(s, y))) \right) \right)^2.$$

Therefore, for $\epsilon \in (0, t - s)$,

$$\xi^\top \gamma_Z \xi \geq J_1 + J_2,$$

where

$$J_1 := \sum_{k=1}^d \int_{s-\epsilon}^s dr \int_0^1 dv \left(\sum_{i=1}^d (\lambda_i - \mu_i) [G_{s-r}(y, v) \sigma_{ik}(u(r, v)) + a_i(k, r, v, s, y)] + W \right)^2,$$

$$J_2 := \sum_{k=1}^d \int_{t-\epsilon}^t dr \int_0^1 dv W^2,$$

$a_i(k, r, v, s, y)$ is defined in (4.3) and

$$W := \sum_{i=1}^d [\mu_i G_{t-r}(x, v) \sigma_{ik}(u(r, v)) + \mu_i a_i(k, r, v, t, x)].$$

From here on, the proof is divided into two cases.

Case 1. In the first case, we assume that $|x - y|^2 \leq t - s$. Choose and fix an $\epsilon \in (0, t - s)$. Then we may write

$$\inf_{\|\xi\|=1} \xi^\top \gamma_Z \xi \geq \min \left(\inf_{\|\xi\|=1, \|\mu\| \geq \epsilon^{\eta/2}} J_2, \inf_{\|\xi\|=1, \|\mu\| \leq \epsilon^{\eta/2}} J_1 \right).$$

We are going to prove that

$$\begin{aligned} \inf_{\|\xi\|=1, \|\mu\|\geq\epsilon^{\eta/2}} J_2 &\geq \epsilon^{\frac{1}{2}+\eta} - Y_{1,\epsilon}, \\ \inf_{\|\xi\|=1, \|\mu\|\leq\epsilon^{\eta/2}} J_1 &\geq \epsilon^{1/2} - Y_{2,\epsilon}, \end{aligned} \quad (6.11)$$

where, for all $q \geq 1$,

$$\mathbb{E}[|Y_{1,\epsilon}|^q] \leq c_1(q)\epsilon^q \quad \text{and} \quad \mathbb{E}[|Y_{2,\epsilon}|^q] \leq c_2(q)\epsilon^{q(\frac{1}{2}+\eta)}. \quad (6.12)$$

We assume these, for the time being, and finish the proof of the proposition in Case 1. Then we will return to proving (6.11) and (6.12).

We can combine (6.11) and (6.12) with Proposition 3.5 to find that

$$\begin{aligned} \mathbb{E} \left[\left(\inf_{\|\xi\|=1} \xi^\top \gamma_Z \xi \right)^{-2pd} \right] &\leq c(t-s)^{-2pd(\frac{1}{2}+\eta)} \\ &\leq \tilde{c} \left[(t-s)^{1/2} + |x-y| \right]^{-2pd(1+2\eta)}, \end{aligned}$$

whence follows the proposition in the case that $|x-y|^2 \leq t-s$. Now we complete our proof of Case 1 by deriving (6.11) and (6.12).

Let us begin with the term that involves J_2 . Inequality (4.4) implies that

$$\inf_{\|\xi\|=1, \|\mu\|\geq\epsilon^{\eta/2}} J_2 \geq \hat{Y}_{1,\epsilon} - Y_{1,\epsilon},$$

where

$$\begin{aligned} \hat{Y}_{1,\epsilon} &:= \frac{2}{3} \inf_{\|\mu\|\geq\epsilon^{\eta/2}} \sum_{k=1}^d \int_{t-\epsilon}^t dr \int_0^1 dv \left(\sum_{i=1}^d \mu_i \sigma_{ik}(u(r,v)) \right)^2 G_{t-r}^2(x,v), \\ Y_{1,\epsilon} &:= 2 \sup_{\|\mu\|\geq\epsilon^{\eta/2}} \sum_{k=1}^d \int_{t-\epsilon}^t dr \int_0^1 dv \left(\sum_{i=1}^d \mu_i a_i(k,r,v,t,x) \right)^2. \end{aligned}$$

In agreement with hypothesis **P2**, and thanks to Lemma 7.2,

$$\hat{Y}_{1,\epsilon} \geq c \inf_{\|\mu\|\geq\epsilon^{\eta/2}} \|\mu\|^2 \epsilon^{1/2} \geq c\epsilon^{\frac{1}{2}+\eta}.$$

Next we apply Lemma 6.11 below [with $s := t$] to find that $\mathbb{E}[|Y_{1,\epsilon}|^q] \leq c\epsilon^q$. This proves the bounds in (6.11) and (6.12) that concern J_2 and $Y_{1,\epsilon}$.

In order to derive the second bound in (6.11), we appeal to (4.4) once more to find that

$$\inf_{\|\xi\|=1, \|\mu\|\leq\epsilon^{\eta/2}} J_1 \geq \hat{Y}_{2,\epsilon} - Y_{2,\epsilon},$$

where

$$\hat{Y}_{2,\epsilon} := \frac{2}{3} \inf_{\|\mu\|\leq\epsilon^{\eta/2}} \sum_{k=1}^d \int_{s-\epsilon}^s dr \int_0^1 dv \left(\sum_{i=1}^d (\lambda_i - \mu_i) \sigma_{ik}(u(r,v)) \right)^2 G_{s-r}^2(y,v),$$

and

$$Y_{2,\epsilon} := 2(W_1 + W_2 + W_3),$$

where

$$\begin{aligned} W_1 &:= \sup_{\|\mu\| \leq \epsilon^{\eta/2}} \sum_{k=1}^d \int_{s-\epsilon}^s dr \int_0^1 dv \left(\sum_{i=1}^d \mu_i G_{t-r}(x, v) \sigma_{ik}(u(r, v)) \right)^2, \\ W_2 &:= \sup_{\|\xi\|=1} \sum_{k=1}^d \int_{s-\epsilon}^s dr \int_0^1 dv \left(\sum_{i=1}^d (\lambda_i - \mu_i) a_i(k, r, v, s, y) \right)^2, \\ W_3 &:= \sup_{\|\mu\| \leq \epsilon^{\eta/2}} \sum_{k=1}^d \int_{s-\epsilon}^s dr \int_0^1 dv \left(\sum_{i=1}^d \mu_i a_i(k, r, v, t, x) \right)^2. \end{aligned}$$

Hypothesis **P2** and Lemma 7.2 together yield

$$\hat{Y}_{2,\epsilon} \geq c\epsilon^{1/2}. \quad (6.13)$$

Next, we apply the Cauchy–Schwarz inequality to find that

$$\begin{aligned} \mathbb{E}[|W_1|^q] &\leq \sup_{\|\mu\| \leq \epsilon^{\eta/2}} \|\mu\|^{2q} \times \mathbb{E} \left[\left| \sum_{k=1}^d \int_{s-\epsilon}^s dr \int_0^1 dv \sum_{i=1}^d (\sigma_{ik}(u(r, v)))^2 G_{t-r}^2(x, v) \right|^q \right] \\ &\leq c\epsilon^{q\eta} \left| \sum_{k=1}^d \int_{s-\epsilon}^s dr \int_0^1 dv G_{t-r}^2(x, v) \right|^q, \end{aligned}$$

thanks to hypothesis **P1**. In light of this, Lemma 7.4 implies that $\mathbb{E}[|W_1|^q] \leq c\epsilon^{q(\frac{1}{2}+\eta)}$.

In order to bound the q -th moment of $|W_2|$, we use the Cauchy–Schwarz inequality together with hypothesis **P1**, and write

$$\begin{aligned} \mathbb{E}[|W_2|^q] &\leq \sup_{\|\mu\| \leq \epsilon^{\eta/2}} \|\lambda - \mu\|^{2q} \times \mathbb{E} \left[\left| \sum_{k=1}^d \int_{s-\epsilon}^s dr \int_0^1 dv \sum_{i=1}^d a_i^2(k, r, v, s, y) \right|^q \right] \\ &\leq C\mathbb{E} \left[\left| \sum_{k=1}^d \int_{s-\epsilon}^s dr \int_0^1 dv \sum_{i=1}^d a_i^2(k, r, v, s, y) \right|^q \right]. \end{aligned}$$

We apply Lemma 6.11 below [with $s := t$] to find that $\mathbb{E}[|W_2|^q] \leq c\epsilon^q$.

Similarly, we find using Lemma 6.11 that

$$\begin{aligned} \mathbb{E}[|W_3|^q] &\leq \sup_{\|\mu\| \leq \epsilon^{\eta/2}} \|\mu\|^{2q} \times \mathbb{E} \left[\left| \sum_{k=1}^d \int_{s-\epsilon}^s dr \int_0^1 dv \sum_{i=1}^d a_i^2(k, r, v, t, x) \right|^q \right] \\ &\leq c\epsilon^{q\eta} (t - s + \epsilon)^{q/2} \epsilon^{q/2} \\ &\leq c\epsilon^{q(\frac{1}{2}+\eta)}. \end{aligned}$$

The preceding bounds for W_1 , W_2 , and W_3 prove, in conjunction, that $\mathbb{E}[|Y_{2,\epsilon}|^q] \leq c_2(q)\epsilon^{q(\frac{1}{2}+\eta)}$. This and (6.13) together prove the bounds in (6.11) and (6.12) that concern J_1 and $Y_{2,\epsilon}$, whence follows the result in Case 1.

Case 2. Now we work on the second case where $|x - y|^2 \geq t - s \geq 0$. Let $\epsilon > 0$ be such that $(1 + \alpha)\epsilon^{1/2} < \frac{1}{2}|x - y|$, where $\alpha > 0$ is large but fixed; its specific value will be decided on later. Then

$$\xi^\top \gamma_Z \xi \geq I_1 + I_2 + I_3,$$

where

$$\begin{aligned} I_1 &:= \sum_{k=1}^d \int_{s-\epsilon}^s dr \int_{y-\sqrt{\epsilon}}^{y+\sqrt{\epsilon}} dv (\mathcal{S}_1 + \mathcal{S}_2)^2, \\ I_2 &:= \sum_{k=1}^d \int_{s-\epsilon}^s dr \int_{x-\sqrt{\epsilon}}^{x+\sqrt{\epsilon}} dv (\mathcal{S}_1 + \mathcal{S}_2)^2, \\ I_3 &:= \sum_{k=1}^d \int_{(t-\epsilon) \vee s}^t dr \int_{x-\sqrt{\epsilon}}^{x+\sqrt{\epsilon}} dv \mathcal{S}_2^2, \end{aligned}$$

and

$$\begin{aligned} \mathcal{S}_1 &:= \sum_{i=1}^d (\lambda_i - \mu_i) [G_{s-r}(y, v) \sigma_{i,k}(u(r, v)) + a_i(k, r, v, s, y)], \\ \mathcal{S}_2 &:= \sum_{i=1}^d \mu_i [G_{t-r}(x, v) \sigma_{i,k}(u(r, v)) + a_i(k, r, v, t, x)]. \end{aligned}$$

From here on, Case 2 is divided into two further sub-cases.

Sub-Case A. Suppose, in addition, that $\epsilon \geq t - s$. In this case, we are going to prove that

$$\inf_{\|\xi\|=1} \xi^\top \gamma_Z \xi \geq c\epsilon^{1/2} - Z_{1,\epsilon}, \quad (6.14)$$

where for all $q \geq 1$,

$$\mathbb{E}[|Z_{1,\epsilon}|^q] \leq c(q)\epsilon^{3q/4}. \quad (6.15)$$

Apply (4.4) to find that

$$I_1 \geq \frac{2}{3} \tilde{A}_1 - B_1^{(1)} - B_1^{(2)},$$

where

$$\begin{aligned} \tilde{A}_1 &:= \sum_{k=1}^d \int_{s-\epsilon}^s dr \int_{y-\sqrt{\epsilon}}^{y+\sqrt{\epsilon}} dv \left(\sum_{i=1}^d [(\lambda_i - \mu_i) G_{s-r}(y, v) + \mu_i G_{t-r}(x, v)] \sigma_{i,k}(u(r, v)) \right)^2, \\ B_1^{(1)} &:= 4\|\lambda - \mu\|^2 \sum_{k=1}^d \int_{s-\epsilon}^s dr \int_{y-\sqrt{\epsilon}}^{y+\sqrt{\epsilon}} dv \sum_{i=1}^d a_i^2(k, r, v, s, y), \end{aligned} \quad (6.16)$$

$$B_1^{(2)} := 4\|\mu\|^2 \sum_{k=1}^d \int_{s-\epsilon}^s dr \int_{y-\sqrt{\epsilon}}^{y+\sqrt{\epsilon}} dv \sum_{i=1}^d a_i^2(k, r, v, t, x). \quad (6.17)$$

Using the inequality

$$(a - b)^2 \geq \frac{2}{3}a^2 - 2ab, \quad (6.18)$$

we see that

$$\tilde{A}_1 \geq \frac{2}{3}A_1 - B_1^{(3)},$$

where

$$\begin{aligned} A_1 &:= \sum_{k=1}^d \int_{s-\epsilon}^s dr \int_{y-\sqrt{\epsilon}}^{y+\sqrt{\epsilon}} dv \left(\sum_{i=1}^d (\lambda_i - \mu_i) G_{s-r}(y, v) \sigma_{ik}(u(r, v)) \right)^2, \\ B_1^{(3)} &:= 2 \sum_{k=1}^d \int_{s-\epsilon}^s dr \int_{y-\sqrt{\epsilon}}^{y+\sqrt{\epsilon}} dv \left(\sum_{i=1}^d (\lambda_i - \mu_i) G_{s-r}(y, v) \sigma_{ik}(u(r, v)) \right) \\ &\quad \times \left(\sum_{i=1}^d \mu_i G_{t-r}(x, v) \sigma_{ik}(u(r, v)) \right). \end{aligned}$$

We can combine terms to find that

$$I_1 \geq \frac{4}{9}A_1 - B_1^{(1)} - B_1^{(2)} - B_1^{(3)}.$$

We proceed in like manner for I_2 , but obtain slightly sharper estimates as follows. Owing to (6.18),

$$I_2 \geq \frac{2}{3}A_2 - B_2^{(1)} - B_2^{(2)} - B_2^{(3)},$$

where

$$\begin{aligned} A_2 &:= \sum_{k=1}^d \int_{s-\epsilon}^s dr \int_{x-\sqrt{\epsilon}}^{x+\sqrt{\epsilon}} dv \left(\sum_{i=1}^d \mu_i G_{t-r}(x, v) \sigma_{ik}(u(r, v)) \right)^2, \\ B_2^{(1)} &:= 2 \sum_{k=1}^d \int_{s-\epsilon}^s dr \int_{x-\sqrt{\epsilon}}^{x+\sqrt{\epsilon}} dv \left(\sum_{i=1}^d \mu_i G_{t-r}(x, v) \sigma_{ik}(u(r, v)) \right) \\ &\quad \times \left(\sum_{i=1}^d \mu_i a_i(k, r, v, t, x) \right), \\ B_2^{(2)} &:= 2 \sum_{k=1}^d \int_{s-\epsilon}^s dr \int_{x-\sqrt{\epsilon}}^{x+\sqrt{\epsilon}} dv \left(\sum_{i=1}^d \mu_i G_{t-r}(x, v) \sigma_{ik}(u(r, v)) \right) \\ &\quad \times \left(\sum_{i=1}^d (\lambda_i - \mu_i) a_i(k, r, v, s, y) \right), \\ B_2^{(3)} &:= 2 \sum_{k=1}^d \int_{s-\epsilon}^s dr \int_{x-\sqrt{\epsilon}}^{x+\sqrt{\epsilon}} dv \left(\sum_{i=1}^d \mu_i G_{t-r}(x, v) \sigma_{ik}(u(r, v)) \right) \\ &\quad \times \left(\sum_{i=1}^d (\lambda_i - \mu_i) G_{s-r}(y, v) \sigma_{ik}(u(r, v)) \right). \end{aligned}$$

Finally, we appeal to (4.4) to find that

$$I_3 \geq \frac{2}{3}A_3 - B_3,$$

where

$$\begin{aligned} A_3 &:= \sum_{k=1}^d \int_{(t-\epsilon) \vee s}^t dr \int_{x-\sqrt{\epsilon}}^{x+\sqrt{\epsilon}} dv \left(\sum_{i=1}^d \mu_i G_{t-r}(x, v) \sigma_{ik}(u(r, v)) \right)^2, \\ B_3 &:= 2 \sum_{k=1}^d \int_{(t-\epsilon) \vee s}^t dr \int_{x-\sqrt{\epsilon}}^{x+\sqrt{\epsilon}} dv \left(\sum_{i=1}^d \mu_i a_i(k, r, v, t, x) \right)^2. \end{aligned} \quad (6.19)$$

By hypothesis **P2**,

$$\begin{aligned} A_1 + A_2 + A_3 &\geq \rho^2 \left(\|\lambda - \mu\|^2 \int_{s-\epsilon}^s dr \int_{y-\sqrt{\epsilon}}^{y+\sqrt{\epsilon}} dv G_{s-r}^2(y, v) \right. \\ &\quad + \|\mu\|^2 \int_{s-\epsilon}^s dr \int_{x-\sqrt{\epsilon}}^{x+\sqrt{\epsilon}} dv G_{t-r}^2(x, v) \\ &\quad \left. + \|\mu\|^2 \int_s^t dr \int_{x-\sqrt{\epsilon}}^{x+\sqrt{\epsilon}} dv G_{t-r}^2(x, v) \right). \end{aligned}$$

Note that we have used the defining assumption of Sub-Case A, namely, that $\epsilon \geq t - s$. Next, we group the last two integrals and apply Lemma 7.2 to find that

$$\begin{aligned} A_1 + A_2 + A_3 &\geq c \left(\|\lambda - \mu\|^2 \epsilon^{1/2} + \|\mu\|^2 \int_{t-\epsilon}^t dr \int_{x-\sqrt{\epsilon}}^{x+\sqrt{\epsilon}} dv G_{t-r}^2(x, v) \right) \\ &\geq c (\|\lambda - \mu\|^2 + \|\mu\|^2) \epsilon^{1/2} \\ &\geq c \epsilon^{1/2}. \end{aligned} \quad (6.20)$$

We are aiming for (6.14), and propose to bound the absolute moments of $B_1^{(i)}$, $B_2^{(i)}$, $i = 1, 2, 3$ and B_3 , separately. According to Lemma 6.11 below with $s = t$,

$$\mathbb{E} \left[\sup_{\|\xi\|=1} |B_3|^q \right] \leq c(q) \epsilon^q. \quad (6.21)$$

Next we bound the absolute moments of $B_1^{(i)}$, $i = 1, 2, 3$. Using hypothesis **P1** and Lemma 6.11, with $t = s$, we find that for all $q \geq 1$,

$$\mathbb{E} \left[\sup_{\|\xi\|=1} |B_1^{(1)}|^q \right] \leq c \epsilon^q. \quad (6.22)$$

In the same way, we see that

$$\mathbb{E} \left[\sup_{\|\xi\|=1} |B_1^{(2)}|^q \right] \leq c(t - s + \epsilon)^{q/2} \epsilon^{q/2}. \quad (6.23)$$

We are in the sub-case A where $t - s \leq \epsilon$. Therefrom, we obtain the following:

$$\mathbb{E} \left[\sup_{\|\xi\|=1} \left| B_1^{(2)} \right|^q \right] \leq c\epsilon^q. \quad (6.24)$$

Finally, we turn to bounding the absolute moments of $B_1^{(3)}$. Hypothesis **P1** assures us that

$$\begin{aligned} \left| B_1^{(3)} \right| &\leq c \int_{s-\epsilon}^s dr \int_{y-\sqrt{\epsilon}}^{y+\sqrt{\epsilon}} dv G_{s-r}(y, v) G_{t-r}(x, v) \\ &\leq c \int_{s-\epsilon}^s dr \int_0^1 dv G_{s-r}(y, v) G_{t-r}(x, v) \\ &= c \int_{s-\epsilon}^s dr G_{t+s-2r}(x, y), \end{aligned}$$

thanks to the semi-group property WALSH [W86, (3.6)] (see (6.44) below). This and Lemma 7.1 together prove that

$$\begin{aligned} \left| B_1^{(3)} \right| &\leq c \int_0^\epsilon \frac{du}{\sqrt{t-s+2u}} \exp\left(-\frac{|x-y|^2}{2(t-s+2u)}\right) \\ &\leq c \int_0^\epsilon \frac{du}{\sqrt{t-s+2u}} \exp\left(-\frac{\alpha^2\epsilon}{2(t-s+2u)}\right), \end{aligned}$$

since $|x-y| \geq 2(1+\alpha)\epsilon^{1/2} \geq \alpha\epsilon^{1/2}$. Now we can change variables [$z := 2(t-s+2u)/(\alpha^2\epsilon)$], and use the bounds $0 \leq t-s \leq \epsilon$ to find that

$$\left| B_1^{(3)} \right| \leq c\epsilon^{1/2}\Psi(\alpha), \quad \text{where} \quad \Psi(\alpha) := \alpha \int_0^{6/\alpha^2} z^{-1/2} e^{-1/z} dz. \quad (6.25)$$

Following exactly in the same way, we see that

$$\left| B_2^{(3)} \right| \leq c\epsilon^{1/2}\Psi(\alpha). \quad (6.26)$$

We can combine (6.22), (6.24) as follows:

$$\mathbb{E} \left[\sup_{\|\xi\|=1} \left| B_1^{(1)} + B_1^{(2)} \right|^q \right] \leq c(q)\epsilon^q. \quad (6.27)$$

On the other hand, we will see in Lemmas 6.13 and 6.14 below that

$$\mathbb{E} \left[\sup_{\|\xi\|=1} \left| B_2^{(1)} + B_2^{(2)} \right|^q \right] \leq c(q)\epsilon^{3q/4}. \quad (6.28)$$

Now, by (6.20), (6.21), (6.25), (6.26), (6.27) and (6.28),

$$\begin{aligned} &\inf_{\|\xi\|=1} \xi^\top \gamma_Z \xi \\ &\geq \left(\frac{1}{3}A_1 + A_2 + A_3 \right) - B_1^{(3)} - B_2^{(3)} - \left(B_1^{(1)} + B_1^{(2)} + B_2^{(1)} + B_2^{(2)} + B_3 \right) \\ &\geq c_1\epsilon^{1/2} - c_2\Psi(\alpha)\epsilon^{1/2} - Z_{1,\epsilon}, \end{aligned}$$

where $Z_{1,\epsilon} := B_1^{(1)} + B_1^{(2)} + B_2^{(1)} + B_2^{(2)} + B_3$ satisfies $\mathbb{E}[|Z_{1,\epsilon}|^q] \leq c_1(q)\epsilon^{3q/4}$. Because $\lim_{\nu \rightarrow \infty} \Psi(\nu) = 0$, we can choose and fix α so large that $c_2\Psi(\alpha) \leq c_1/4$ for the c_1 and c_2 of the preceding displayed equation. This yields,

$$\inf_{\|\xi\|=1} \xi^\top \gamma_Z \xi \geq c\epsilon^{1/2} - Z_{1,\epsilon}, \quad (6.29)$$

as in (6.14) and (6.15).

Sub-Case B. In this final (sub-) case we suppose that $\epsilon \leq t - s \leq |x - y|^2$. Choose and fix $0 < \epsilon < t - s$. During the course of our proof of Case 1, we established the following:

$$\inf_{\|\xi\|=1} \xi^\top \gamma_Z \xi \geq \min \left(c\epsilon^{\frac{1}{2}+\eta} - Y_{1,\epsilon}, c\epsilon^{1/2} - Y_{2,\epsilon} \right),$$

where

$$\mathbb{E}[|Y_{1,\epsilon}|^q] \leq c(q)\epsilon^q \quad \text{and} \quad \mathbb{E}[|Y_{2,\epsilon}|^q] \leq c(q)\epsilon^{q(\frac{1}{2}+\eta)}.$$

See (6.11) and (6.12). Consider this in conjunction with (6.14) to find that for all $0 < \epsilon < \frac{1}{4}(1 + \alpha)^{-2}|x - y|^2$,

$$\inf_{\|\xi\|=1} \xi^\top \gamma_Z \xi \geq \min \left(c\epsilon^{\frac{1}{2}+\eta} - Y_{1,\epsilon}, c\epsilon^{1/2} - Y_{2,\epsilon} - Z_{1,\epsilon} \mathbf{1}_{\{t-s < \epsilon\}} \right).$$

Because of this and (6.15), Proposition 3.5 implies that

$$\begin{aligned} \mathbb{E} \left[\left(\inf_{\|\xi\|=1} \xi^\top \gamma_Z \xi \right)^{-2pd} \right] &\leq c|x - y|^{2(-2dp)(\frac{1}{2}+\eta)} \\ &\leq c(|t - s|^{1/2} + |x - y|)^{-2dp(1+2\eta)}. \end{aligned}$$

This concludes the proof of Proposition 6.9(a).

If $t = s$, then Sub-Case B does not arise, and so we get directly from (6.29) and Proposition 3.5 that

$$\mathbb{E} \left[\left(\inf_{\|\xi\|=1} \xi^\top \gamma_Z \xi \right)^{-2pd} \right] \leq c|x - y|^{-2dp}.$$

This proves (b) and concludes the proof of Proposition 6.9. \square

Remark 6.10. If σ and b are constant, then $a_i = 0$, so η can be taken to be 0. This gives the correct upper bound in the Gaussian case, which shows that the method of proof of Proposition 6.9 is rather tight.

We finally prove three results that we have used in the proof of Proposition 6.9.

Lemma 6.11. *Assume P1. For all $T > 0$ and $q \geq 1$, there exists a constant $c = c(q, T) \in (0, \infty)$ such that for every $0 < \epsilon \leq s \leq t \leq T$ and $x \in [0, 1]$,*

$$\mathbb{E} \left[\left(\sum_{k=1}^d \int_{s-\epsilon}^s dr \int_0^1 dv \sum_{i=1}^d a_i^2(k, r, v, t, x) \right)^q \right] \leq c(t - s + \epsilon)^{q/2} \epsilon^{q/2}.$$

Proof. Define

$$A := \sum_{k=1}^d \int_{s-\epsilon}^s dr \int_0^1 dv \sum_{i=1}^d a_i^2(k, r, v, t, x).$$

Use (4.3) to write

$$\mathbb{E}[|A|^q] \leq c(\mathbb{E}[|A_1|^q] + \mathbb{E}[|A_2|^q]),$$

where

$$A_1 := \sum_{i,j,k=1}^d \int_{s-\epsilon}^s dr \int_0^1 dv \left| \int_r^t \int_0^1 G_{t-\theta}(x, \eta) D_{r,v}^{(k)}(\sigma_{ij}(u(\theta, \eta))) W^j(d\theta, d\eta) \right|^2,$$

and

$$A_2 := \sum_{i,k=1}^d \int_{s-\epsilon}^s dr \int_0^1 dv \left| \int_r^t d\theta \int_0^1 d\eta G_{t-\theta}(x, \eta) D_{r,v}^{(k)}(b_i(u(\theta, \eta))) \right|^2.$$

We bound the q -th moment of A_1 and A_2 separately.

As regards A_1 , we apply the Burkholder inequality for Hilbert-space-valued martingales (Lemma 7.6) to find that

$$\mathbb{E}[|A_1|^q] \leq c \sum_{i,j,k=1}^d \mathbb{E} \left[\left| \int_{s-\epsilon}^t d\theta \int_0^1 d\eta \int_{s-\epsilon}^s dr \int_0^1 dv \Theta^2 \right|^q \right], \quad (6.30)$$

where

$$\begin{aligned} \Theta &:= \mathbf{1}_{\{\theta > r\}} G_{t-\theta}(x, \eta) \left| D_{r,v}^{(k)}(\sigma_{ij}(u(\theta, \eta))) \right| \\ &\leq c \mathbf{1}_{\{\theta > r\}} G_{t-\theta}(x, \eta) \left| \sum_{l=1}^d D_{r,v}^{(k)}(u_l(\theta, \eta)) \right|, \end{aligned}$$

thanks to hypothesis **P1**. Thus,

$$\mathbb{E}[|A_1|^q] \leq c \sum_{k=1}^d \mathbb{E} \left[\left| \int_{s-\epsilon}^t d\theta \int_0^1 d\eta G_{t-\theta}^2(x, \eta) \int_{s-\epsilon}^{s \wedge \theta} dr \int_0^1 dv \left(\sum_{l=1}^d D_{r,v}^{(k)}(u_l(\theta, \eta)) \right)^2 \right|^q \right].$$

We apply Hölder's inequality with respect to the measure $G_{t-\theta}^2(x, \eta) d\theta d\eta$ to find that

$$\begin{aligned} \mathbb{E}[|A_1|^q] &\leq c \left(\int_{s-\epsilon}^t d\theta \int_0^1 d\eta G_{t-\theta}^2(x, \eta) \right)^{q-1} \\ &\quad \times \int_{s-\epsilon}^t d\theta \int_0^1 d\eta G_{t-\theta}^2(x, \eta) \sum_{k=1}^d \mathbb{E} \left[\left| \int_{s-\epsilon}^{s \wedge \theta} dr \int_0^1 dv \Upsilon^2 \right|^q \right], \end{aligned} \quad (6.31)$$

where $\Upsilon := \sum_{l=1}^d D_{r,v}^{(k)}(u_l(\theta, \eta))$. Lemma 7.3 assures us that

$$\left(\int_{s-\epsilon}^t d\theta \int_0^1 d\eta G_{t-\theta}^2(x, \eta) \right)^{q-1} \leq c(t-s+\epsilon)^{(q-1)/2}. \quad (6.32)$$

On the other hand, Lemma 7.5 implies that

$$\sum_{k=1}^d \mathbb{E} \left[\left| \int_{s-\epsilon}^{s \wedge \theta} dr \int_0^1 dv \Upsilon^2 \right|^q \right] \leq c\epsilon^{q/2},$$

where $c \in (0, \infty)$ does not depend on $(\theta, \eta, s, t, \epsilon, x)$. Consequently,

$$\begin{aligned} & \int_{s-\epsilon}^t d\theta \int_0^1 d\eta G_{t-\theta}^2(x, \eta) \sum_{k=1}^d \mathbb{E} \left[\left| \int_{s-\epsilon}^{s \wedge \theta} dr \int_0^1 dv \Upsilon^2 \right|^q \right] \\ & \leq c\epsilon^{q/2} \int_{s-\epsilon}^t d\theta \int_0^1 d\eta G_{t-\theta}^2(x, \eta) \\ & \leq c\epsilon^{q/2} (t - s + \epsilon)^{1/2}. \end{aligned} \tag{6.33}$$

Equations (6.31), (6.32), and (6.33) together imply that

$$\mathbb{E} [|A_1|^q] \leq c(t - s + \epsilon)^{q/2} \epsilon^{q/2}. \tag{6.34}$$

This is the desired bound for the q -th moment of A_1 . Next we derive a similar bound for A_2 . This will finish the proof. By the Cauchy–Schwarz inequality

$$\mathbb{E} [|A_2|^q] \leq c(t - s + \epsilon)^q \sum_{i,k=1}^d \mathbb{E} \left[\left| \int_{s-\epsilon}^s dr \int_0^1 dv \int_r^t d\theta \int_0^1 d\eta \Phi^2 \right|^q \right],$$

where $\Phi := G_{t-\theta}(x, \eta) |D_{r,v}^{(k)}(b_i(u(\theta, \eta)))|$. From here on, the q -th moment of A_2 is estimated as that of A_1 was; cf. (6.30), and this yields $\mathbb{E}[|A_2|^q] \leq c(t - s + \epsilon)^{3q/2} \epsilon^{q/2}$. This completes the proof. \square

Remark 6.12. It is possible to prove that $\mathbb{E}[|A_1|]$ is *at least* a constant times $(t-s+\epsilon)^{1/2} \epsilon^{1/2}$. In this sense, the preceding result is not improvable.

Lemma 6.13. *Assume P1. Fix $T > 0$ and $q \geq 1$. Then there exists $c = c(q, T)$ such that for all $x \in [0, 1]$, $0 \leq s \leq t \leq T$, and $\epsilon \in (0, 1)$,*

$$\mathbb{E} \left[\sup_{\mu \in \mathbb{R}^d: \|\mu\| \leq 1} |B_2^{(1)}|^q \right] \leq c\epsilon^{3q/4}.$$

Proof. Define

$$\hat{B}_2^{(1)}(k, i) := \int_{s-\epsilon}^s dr \int_{x-\sqrt{\epsilon}}^{x+\sqrt{\epsilon}} dv G_{t-r}(x, v) |a_i(k, r, v, t, x)|.$$

Then, by the Cauchy–Schwarz inequality,

$$\mathbb{E} \left[\left| \hat{B}_2^{(1)}(k, i) \right|^q \right] \leq J_1 J_2, \tag{6.35}$$

where

$$J_1 := \left(\int_{s-\epsilon}^s dr \int_{x-\sqrt{\epsilon}}^{x+\sqrt{\epsilon}} dv G_{t-r}^2(x, v) \right)^{q/2},$$

$$J_2 := \mathbb{E} \left[\left(\int_{s-\epsilon}^s dr \int_{x-\sqrt{\epsilon}}^{x+\sqrt{\epsilon}} dv a_i^2(k, r, v, t, x) \right)^{q/2} \right].$$

On one hand, according to Lemma 7.4,

$$J_1 \leq c \frac{\epsilon^{q/2}}{(t-s+\epsilon)^{q/4}}. \quad (6.36)$$

On the other hand, Lemma 6.11 assures us that

$$J_2 \leq c'(t-s+\epsilon)^{q/4} \epsilon^{q/4}. \quad (6.37)$$

By combining (6.35), (6.36), and (6.37), we find that $\mathbb{E}[|\hat{B}_2^{(1)}(k, i)|^q] \leq c\epsilon^{3q/4}$ for a constant $c \in (0, \infty)$ that does not depend on ϵ . By hypothesis **P1**,

$$\mathbb{E} \left[\sup_{\mu \in \mathbb{R}^d: \|\mu\| \leq 1} |B_2^{(1)}|^q \right] \leq c \sum_{k=1}^d \mathbb{E} \left[\sum_{i=1}^d |\hat{B}_2^{(1)}(k, i)|^q \right] \leq c\epsilon^{3q/4}, \quad (6.38)$$

as asserted. \square

Lemma 6.14. *Assume **P1**. Fix $T > 0$ and $q \geq 1$. Then there exists $c = c(q, T)$ such that for any $x \in [0, 1]$, $0 \leq s \leq t \leq T$, and $\epsilon \in (0, 1)$,*

$$\mathbb{E} \left[\sup_{\xi = (\lambda, \mu) \in \mathbb{R}^{2d}: \|\xi\| = 1} |B_2^{(2)}|^q \right] \leq c\epsilon^{3q/4}.$$

Proof. Define

$$\hat{B}_2^{(2)}(k, i) := \int_{s-\epsilon}^s dr \int_{x-\sqrt{\epsilon}}^{x+\sqrt{\epsilon}} dv G_{t-r}(x, v) |a_i(k, r, v, s, y)|.$$

Then, by the Cauchy–Schwarz inequality,

$$\mathbb{E} \left[\left| \hat{B}_2^{(2)}(k, i) \right|^q \right] \leq J_1 J_2,$$

where

$$J_1 := \left(\int_{s-\epsilon}^s dr \int_{x-\sqrt{\epsilon}}^{x+\sqrt{\epsilon}} dv G_{t-r}^2(x, v) \right)^{q/2},$$

$$J_2 := \mathbb{E} \left[\left(\int_{s-\epsilon}^s dr \int_{x-\sqrt{\epsilon}}^{x+\sqrt{\epsilon}} dv a_i^2(k, r, v, s, y) \right)^{q/2} \right].$$

According to (6.36),

$$J_1 \leq c \frac{\epsilon^{q/2}}{(t-s+\epsilon)^{q/4}} \leq c' \epsilon^{q/4}.$$

On the other hand, Lemma 6.11, with $t = s$, assures us that $J_2 \leq c'' \epsilon^{q/2}$. It follows that the q -th absolute moment of $\hat{B}_2^{(2)}(k, i)$ is at most $c\epsilon^{3q/4}$. An appeal to the triangle inequality finishes the proof; see (6.38) where a similar argument was worked out in detail. \square

6.4.2 Large Eigenvalues

Proposition 6.15. *Assume **P1** and **P2**. Fix $T > 0$ and $p > 1$. Then there exists $C = C(p, T)$ such that for all $0 \leq s < t \leq T$ with $t - s < \frac{1}{2}$, $x, y \in (0, 1)$, $x \neq y$,*

$$\mathbb{E} \left[\left(\prod_{i=1}^d (\xi^i)^T \gamma_Z \xi^i \right)^{-p} \right] \leq C,$$

where ξ^1, \dots, ξ^d are the vectors from Lemma 6.8.

Proof. Let ξ^1, \dots, ξ^d , written as in (6.10), be such that $\alpha_1 \geq \alpha_0, \dots, \alpha_d \geq \alpha_0$ for some $\alpha_0 > 0$. In order to simplify the exposition, we assume that $0 < \alpha = \alpha_1 = \dots = \alpha_d \leq 1$, since the general case follows along the same lines. Let $0 < \epsilon < s \leq t$. As in the proof of Proposition 6.9, we note first that $\prod_{i=1}^d (\xi^i)^T \gamma_Z \xi^i$ is bounded below by

$$\begin{aligned} & \sum_{k=1}^d \int_{s-\epsilon}^s dr \int_0^1 dv \left(\sum_{i=1}^d \left[\left(\alpha \tilde{\lambda}^i G_{s-r}(y, v) \right. \right. \right. \\ & \quad \left. \left. \left. + \tilde{\mu}^i \sqrt{1-\alpha^2} (G_{t-r}(x, v) - G_{s-r}(y, v)) \right) \sigma_{ik}(u(r, v)) \right. \right. \\ & \quad \left. \left. + \alpha \tilde{\lambda}^i a_i(k, r, v, s, y) \right. \right. \\ & \quad \left. \left. \left. + \tilde{\mu}^i \sqrt{1-\alpha^2} (a_i(k, r, v, t, x) - a_i(k, r, v, s, y)) \right] \right)^2 \\ & + \sum_{k=1}^d \int_{s \vee (t-\epsilon)}^t dr \int_0^1 dv \left(\sum_{i=1}^d \left[\tilde{\mu}^i \sqrt{1-\alpha^2} G_{t-r}(x, v) \sigma_{ik}(u(r, v)) \right. \right. \\ & \quad \left. \left. + \tilde{\mu}^i \sqrt{1-\alpha^2} a_i(k, r, v, t, x) \right] \right)^2. \end{aligned} \tag{6.39}$$

We intend to use Proposition 3.5 with $\epsilon_0 > 0$ fixed, so we seek lower bounds for this expression for $0 < \epsilon < \epsilon_0$.

Case 1. $t - s \leq \epsilon$. Then, by (4.4), the expression in (6.39) is bounded below by

$$\frac{2}{3} (f_1(s, t, \epsilon, \alpha, \tilde{\lambda}, \tilde{\mu}, x, y) + f_2(s, t, \epsilon, \alpha, \tilde{\lambda}, \tilde{\mu}, x, y)) - 2I_\epsilon,$$

where, from hypothesis **P2**,

$$f_1 \geq c\rho^2 \int_{s-\epsilon}^s dr \int_0^1 dv \left\| \alpha \tilde{\lambda} G_{s-r}(y, v) + \sqrt{1-\alpha^2} \tilde{\mu} (G_{t-r}(x, v) - G_{s-r}(y, v)) \right\|^2, \quad (6.40)$$

$$f_2 \geq c\rho^2 \int_{s \vee (t-\epsilon)}^t dr \int_0^1 dv \left\| \tilde{\mu} \sqrt{1-\alpha^2} G_{t-r}(x, v) \right\|^2, \quad (6.41)$$

and $I_\epsilon = I_{1,\epsilon} + I_{2,\epsilon} + I_{3,\epsilon}$, where

$$\begin{aligned} I_{1,\epsilon} &:= \sum_{k=1}^d \int_{s-\epsilon}^s dr \int_0^1 dv \left(\sum_{i=1}^d [\alpha \tilde{\lambda}^i - \tilde{\mu}_i \sqrt{1-\alpha^2}] a_i(k, r, v, s, y) \right)^2, \\ I_{2,\epsilon} &:= \sum_{k=1}^d \int_{s-\epsilon}^s dr \int_0^1 dv \left(\sum_{i=1}^d \tilde{\mu}_i \sqrt{1-\alpha^2} a_i(k, r, v, t, x) \right)^2, \\ I_{3,\epsilon} &:= \sum_{k=1}^d \int_{t-\epsilon}^t dr \int_0^1 dv \left(\sum_{i=1}^d \tilde{\mu}_i \sqrt{1-\alpha^2} a_i(k, r, v, t, x) \right)^2. \end{aligned}$$

There are obvious similarities between the terms $I_{1,\epsilon}$ and $B_1^{(1)}$ in (6.16). Thus, we apply the same method that was used to bound $\mathbb{E}[|B_1^{(1)}|^q]$ to deduce that $\mathbb{E}[|I_{1,\epsilon}|^q] \leq c(q)\epsilon^q$. Since $I_{2,\epsilon}$ is similar to $B_1^{(2)}$ from (6.17) and $t-s \leq \epsilon$, we see using (6.24) that $\mathbb{E}[|I_{2,\epsilon}|^q] \leq c(q)\epsilon^q$. Finally, using the similarity between $I_{3,\epsilon}$ and B_3 in (6.19), we see that $\mathbb{E}[|I_{3,\epsilon}|^q] \leq c(q)\epsilon^q$.

We claim that there exists $\alpha_0 > 0$, $\epsilon_0 > 0$ and $c_0 > 0$ such that

$$f_1 + f_2 \geq c_0 \sqrt{\epsilon} \quad \text{for all } \alpha \in [\alpha_0, 1], \epsilon \in (0, \epsilon_0], s, t \in [1, 2], x, y \in [0, 1]. \quad (6.42)$$

This will imply in particular that for $\epsilon \geq t-s$,

$$\prod_{i=1}^d (\xi^i)^\top \gamma_Z \xi^i \geq c_0 \epsilon^{1/2} - 2I_\epsilon,$$

where $\mathbb{E}[|I_\epsilon|^q] \leq c(q)\epsilon^q$.

In order to prove (6.42), first define

$$p_t(x, y) := (4\pi t)^{-1/2} e^{-(x-y)^2/(4t)}.$$

In addition, let $g_1(s, t, \epsilon, \alpha, \tilde{\lambda}, \tilde{\mu}, x, y)$ and $g_2(s, t, \epsilon, \alpha, \tilde{\lambda}, \tilde{\mu}, x, y)$ be defined by the same expressions as the right-hand sides of (6.40) and (6.41), but with $G_{s-r}(x, v)$ replaced by $p_{s-r}(x-v)$, and \int_0^1 replaced by $\int_{-\infty}^{+\infty}$.

Observe that $g_1 \geq 0$, $g_2 \geq 0$, and if $g_1 = 0$, then for all $v \in \mathbb{R}^d$,

$$\left\| \alpha p_{s-r}(y-v) \tilde{\lambda} + \sqrt{1-\alpha^2} (p_{t-r}(x-v) - p_{s-r}(y-v)) \tilde{\mu} \right\| = 0. \quad (6.43)$$

If, in addition, $\tilde{\lambda} = \tilde{\mu}$, then we get that for all $v \in \mathbb{R}^d$,

$$(\alpha - \sqrt{1-\alpha^2}) p_{s-r}(y-v) + \sqrt{1-\alpha^2} p_{t-r}(x-v) = 0.$$

We take Fourier transforms to deduce from this that for all $\xi \in \mathbb{R}^d$,

$$\left(\alpha - \sqrt{1 - \alpha^2}\right) e^{i\xi y} = -\sqrt{1 - \alpha^2} e^{i\xi x} e^{(s-t)\xi^2}.$$

If $x = y$, then it follows that $s = t$ and $\alpha - \sqrt{1 - \alpha^2} = -\sqrt{1 - \alpha^2}$. Hence, if $\alpha \neq 0$, $x = y$ and $\tilde{\lambda} = \tilde{\mu}$, then $g_1 > 0$. We shall make use of this observation shortly.

Because $\|\tilde{\lambda}\| = \|\tilde{\mu}\| = 1$, f_1 is bounded below by

$$\begin{aligned} & c\rho^2 \int_{s-\epsilon}^s dr \int_0^1 dv \left(\alpha^2 G_{s-r}^2(y, v) + (1 - \alpha^2) (G_{t-r}(x, v) - G_{s-r}(y, v))^2 \right. \\ & \quad \left. + 2\alpha\sqrt{1 - \alpha^2} G_{s-r}(y, v) (G_{t-r}(x, v) - G_{s-r}(y, v)) (\tilde{\lambda} \cdot \tilde{\mu}) \right) \\ & = c\rho^2 \int_{s-\epsilon}^s dr \int_0^1 dv \left(\left(\alpha - \sqrt{1 - \alpha^2} \right)^2 G_{s-r}^2(y, v) + (1 - \alpha^2) G_{t-r}^2(x, v) \right. \\ & \quad \left. + 2 \left(\alpha - \sqrt{1 - \alpha^2} \right) \sqrt{1 - \alpha^2} G_{s-r}(y, v) G_{t-r}(x, v) \right. \\ & \quad \left. + 2\alpha\sqrt{1 - \alpha^2} G_{s-r}(y, v) (G_{t-r}(x, v) - G_{s-r}(y, v)) (\tilde{\lambda} \cdot \tilde{\mu} - 1) \right). \end{aligned}$$

Recall the semigroup property

$$\int_0^1 dv G_{s-r}(y, v) G_{t-r}(x, v) = G_{s+t-2r}(x, y) \quad (6.44)$$

(see WALSH [W86, (3.6)]). We set $h := t - s$ and change variables [$\bar{r} := s - r$] to obtain the following bound:

$$\begin{aligned} f_1 & \geq c\rho^2 \int_0^\epsilon dr \left(\left(\alpha - \sqrt{1 - \alpha^2} \right)^2 G_{2r}(y, y) + (1 - \alpha^2) G_{2h+2r}(x, x) \right. \\ & \quad \left. + 2 \left(\alpha - \sqrt{1 - \alpha^2} \right) \sqrt{1 - \alpha^2} G_{h+2r}(x, y) \right. \\ & \quad \left. + 2\alpha\sqrt{1 - \alpha^2} (G_{h+2r}(x, y) - G_{2r}(y, y)) (\tilde{\lambda} \cdot \tilde{\mu} - 1) \right). \end{aligned}$$

Recall ([W86, p.318]), that

$$G_t(x, y) = p_t(x, y) + H_t(x, y),$$

where $H_t(x, y)$ is a continuous function that is uniformly bounded over $(t, x, y) \in (0, \infty) \times (0, 1) \times (0, 1)$. Therefore, $f_1 \geq c\rho^2 \tilde{g}_1 - c\epsilon$, where

$$\begin{aligned} \tilde{g}_1 & := \tilde{g}_1(h, \epsilon, \alpha, \tilde{\lambda}, \tilde{\mu}, x, y) \\ & = \int_0^\epsilon dr \left(\left(\alpha - \sqrt{1 - \alpha^2} \right)^2 p_{2r}(y, y) + (1 - \alpha^2) p_{2h+2r}(x, x) \right. \\ & \quad \left. + 2 \left(\alpha - \sqrt{1 - \alpha^2} \right) \sqrt{1 - \alpha^2} p_{h+2r}(x, y) \right. \\ & \quad \left. + 2\alpha\sqrt{1 - \alpha^2} (p_{h+2r}(x, y) - p_{2r}(y, y)) (\tilde{\lambda} \cdot \tilde{\mu} - 1) \right). \end{aligned}$$

We can recognize that

$$p_{h+2r}(x, y) - p_{2r}(y, y) = \frac{\exp(-(x-y)^2/(4(h+2r)))}{\sqrt{4\pi(h+2r)}} - \frac{1}{\sqrt{4\pi(2r)}} \leq 0.$$

Also, $\tilde{\lambda} \cdot \tilde{\mu} - 1 \leq 0$. Thus,

$$\tilde{g}_1 \geq \hat{g}_1,$$

where

$$\begin{aligned} \hat{g}_1 &:= \hat{g}_1(h, \epsilon, \alpha, x, y) \\ &= \int_0^\epsilon dr \left((\alpha - \sqrt{1-\alpha^2})^2 p_{2r}(y, y) + (1-\alpha^2) p_{2h+2r}(x, x) \right. \\ &\quad \left. + 2(\alpha - \sqrt{1-\alpha^2}) \sqrt{1-\alpha^2} p_{h+2r}(x, y) \right). \end{aligned}$$

Therefore,

$$\begin{aligned} \hat{g}_1 &= \int_0^\epsilon dr \left((\alpha - \sqrt{1-\alpha^2})^2 \frac{1}{\sqrt{8\pi r}} + (1-\alpha^2) \frac{1}{\sqrt{8\pi(h+r)}} \right. \\ &\quad \left. + 2(\alpha - \sqrt{1-\alpha^2}) \sqrt{1-\alpha^2} p_{h+2r}(x, y) \right). \end{aligned}$$

On the other hand, by (6.44) above,

$$\begin{aligned} f_2 &\geq \int_0^{\epsilon \wedge (t-s)} dr (1-\alpha^2) G_{2r}(y, y) \\ &\geq \tilde{g}_2 := \int_0^{\epsilon \wedge h} dr (1-\alpha^2) p_{2r}(y, y) - C\epsilon \\ &= (1-\alpha^2) \sqrt{\epsilon \wedge h} - C\epsilon. \end{aligned}$$

Finally, we conclude that

$$\begin{aligned} f_1 + f_2 &\geq \hat{g}_1 + \tilde{g}_2 - 2C\epsilon \\ &= (\alpha - \sqrt{1-\alpha^2})^2 \frac{\sqrt{\epsilon}}{\sqrt{2\pi}} + \frac{1-\alpha^2}{\sqrt{2\pi}} (\sqrt{h+\epsilon} - \sqrt{h}) \\ &\quad + 2(\alpha - \sqrt{1-\alpha^2}) \sqrt{1-\alpha^2} \int_0^\epsilon dr p_{h+2r}(x, y) \\ &\quad + \frac{1-\alpha^2}{\sqrt{2\pi}} \sqrt{\epsilon \wedge h} - 2C\epsilon. \end{aligned}$$

Now we consider two different cases.

Case (i). Suppose $\alpha - \sqrt{1-\alpha^2} \geq 0$, that is, $\alpha \geq 2^{-1/2}$. Then

$$\epsilon^{-1/2} (\hat{g}_1 + \tilde{g}_2) \geq \phi_1 \left(\alpha, \frac{h}{\epsilon} \right) - 2C\epsilon^{1/2},$$

where

$$\phi_1(\alpha, z) := \frac{1}{\sqrt{2\pi}} \left((\alpha - \sqrt{1 - \alpha^2})^2 + (1 - \alpha^2) \frac{1}{\sqrt{1+z} + \sqrt{z}} + (1 - \alpha^2) \sqrt{1 \wedge z} \right).$$

Clearly,

$$\begin{aligned} \inf_{\alpha \geq 2^{-1/2}} \inf_{z > 0} \phi_1(\alpha, z) &\geq \inf_{\alpha > 2^{-1/2}} \frac{(\alpha - \sqrt{1 - \alpha^2})^2 + c_0(1 - \alpha^2)}{\sqrt{2\pi}} \\ &> \phi_0 > 0. \end{aligned}$$

Thus,

$$\inf_{\alpha \geq 2^{-1/2}, h \geq 0, 0 < \epsilon \leq \epsilon_0} \epsilon^{-1/2} (\hat{g}_1 + \tilde{g}_2) > 0.$$

Case (ii). Now we consider the case where $\alpha - \sqrt{1 - \alpha^2} < 0$, that is, $\alpha < 2^{-1/2}$. In this case,

$$\epsilon^{-1/2} (\hat{g}_1 + \tilde{g}_2) \geq \psi_1 \left(\alpha, \frac{h}{\epsilon} \right) - 2C\epsilon^{1/2},$$

where

$$\begin{aligned} \psi_1(\alpha, z) &:= \frac{1}{\sqrt{2\pi}} \left((\alpha - \sqrt{1 - \alpha^2})^2 + (1 - \alpha^2) \frac{1}{\sqrt{1+z} + \sqrt{z}} \right. \\ &\quad \left. - 2 \left(\sqrt{1 - \alpha^2} - \alpha \right) \sqrt{1 - \alpha^2} \frac{\sqrt{2}}{\sqrt{2+z} + \sqrt{z}} + (1 - \alpha^2) \sqrt{1 \wedge z} \right). \end{aligned}$$

Note that $\psi_1(\alpha, z) > 0$ if $\alpha \neq 0$. This corresponds to the observation made in the lines following (6.43). Moreover, for $\alpha \geq \alpha_0 > 0$, $\lim_{z \downarrow 0} \psi_1(\alpha, z) \geq (2\pi)^{-1/2} \alpha_0^2$, and

$$\lim_{z \uparrow +\infty} \psi_1(\alpha, z) \geq \inf_{\alpha > 0} \frac{1}{\sqrt{2\pi}} \left((\alpha - \sqrt{1 - \alpha^2})^2 + (1 - \alpha^2) \right) > 0.$$

Therefore,

$$\inf_{\alpha \in [\alpha_0, 2^{-1/2}], z \geq 0} \psi_1(\alpha, z) > 0.$$

This concludes the proof of the claim (6.42).

Case 2. $t - s > \epsilon$. In accord with (6.39), we are interested in

$$\inf_{1 \geq \alpha \geq \alpha_0} \prod_{i=1}^d (\xi^i)^\top \gamma_Z \xi^i := \min(E_{1,\epsilon}, E_{2,\epsilon}),$$

where

$$\begin{aligned} E_{1,\epsilon} &:= \inf_{\alpha_0 \leq \alpha \leq \sqrt{1 - \epsilon^\eta}} \prod_{i=1}^d (\xi^i)^\top \gamma_Z \xi^i, \\ E_{2,\epsilon} &:= \inf_{\sqrt{1 - \epsilon^\eta} \leq \alpha \leq 1} \prod_{i=1}^d (\xi^i)^\top \gamma_Z \xi^i. \end{aligned}$$

Clearly,

$$E_{1,\epsilon} \geq \frac{2}{3}f_2 - 2I_{3,\epsilon}.$$

Since $\alpha \leq \sqrt{1 - \epsilon^\eta}$ is equivalent to $\sqrt{1 - \alpha^2} \geq \epsilon^{\eta/2}$, we use hypothesis **P2** to deduce that

$$f_2 \geq c\rho^2\epsilon^\eta \int_{t-\epsilon}^t dr \int_0^1 dv G_{t-r}^2(x, v) \geq c\rho^2\epsilon^{\frac{1}{2}+\eta}.$$

Therefore,

$$E_{1,\epsilon} \geq c\rho^2\epsilon^{\frac{1}{2}+\eta} - I_{3,\epsilon},$$

and we have seen that $I_{3,\epsilon}$ has the desirable property $\mathbb{E}[|I_{3,\epsilon}|^q] \leq c(q)\epsilon^q$.

In order to estimate $E_{2,\epsilon}$, we observe using (6.39) that

$$E_{2,\epsilon} \geq \frac{2}{3}\tilde{f}_1 - \tilde{J}_{1,\epsilon} - \tilde{J}_{2,\epsilon} - \tilde{J}_{3,\epsilon} - \tilde{J}_{4,\epsilon},$$

where

$$\begin{aligned} \tilde{f}_1 &\geq \alpha^2 \sum_{k=1}^d \int_{s-\epsilon}^s dr \int_0^1 dv \left(\sum_{i=1}^d \tilde{\lambda}^i \sigma_{ik}(u(r, v)) \right)^2 G_{s-r}^2(y, v), \\ \tilde{J}_{1,\epsilon} &= 2(1 - \alpha^2) \sum_{k=1}^d \int_{s-\epsilon}^s dr \int_0^1 dv \left(\sum_{i=1}^d \tilde{\mu}_i \sigma_{ik}(u(r, v)) \right)^2 G_{t-r}^2(x, v), \\ \tilde{J}_{2,\epsilon} &= 2(1 - \alpha^2) \sum_{k=1}^d \int_{s-\epsilon}^s dr \int_0^1 dv \left(\sum_{i=1}^d \tilde{\mu}_i \sigma_{ik}(u(r, v)) \right)^2 G_{s-r}^2(y, v), \\ \tilde{J}_{3,\epsilon} &= 2 \sum_{k=1}^d \int_{s-\epsilon}^s dr \int_0^1 dv \left(\sum_{i=1}^d \left(\alpha \tilde{\lambda}^i - \tilde{\mu}_i \sqrt{1 - \alpha^2} \right) a_i(k, r, v, s, y) \right)^2, \\ \tilde{J}_{4,\epsilon} &= 2(1 - \alpha^2) \sum_{k=1}^d \int_{s-\epsilon}^s dr \int_0^1 dv \left(\sum_{i=1}^d \tilde{\mu}_i a_i(k, r, v, t, x) \right)^2. \end{aligned}$$

Because $\alpha^2 \geq 1 - \epsilon^\eta$ and $\epsilon \leq t - s \leq \frac{1}{2}$, hypothesis **P2** and Lemma 7.2 imply that $\tilde{f}_1 \geq c\epsilon^{1/2}$. On the other hand, since $1 - \alpha^2 \leq \epsilon^\eta$, we can use hypothesis **P1** and Lemma 7.4 to see that

$$\mathbb{E} \left[\left| \tilde{J}_{1,\epsilon} \right|^q \right] \leq c(q)\epsilon^{q\eta}\epsilon^{q/2} = c(q)\epsilon^{(\frac{1}{2}+\eta)q},$$

and similarly, using Lemma 7.3, $\mathbb{E}[|\tilde{J}_{2,\epsilon}|^q] \leq c(q)\epsilon^{(\frac{1}{2}+\eta)q}$. The term $\tilde{J}_{3,\epsilon}$ is equal to $2I_{1,\epsilon}$, so $\mathbb{E}[|\tilde{J}_{3,\epsilon}|^q] \leq c\epsilon^q$, and $\tilde{J}_{4,\epsilon}$ is similar to $B_1^{(2)}$ from (6.17), so we find using (6.23) that

$$\mathbb{E} \left[\left| \tilde{J}_{4,\epsilon} \right|^q \right] \leq c\epsilon^{q\eta}(t - s + \epsilon)^{q/2}\epsilon^{q/2} \leq c\epsilon^{(\frac{1}{2}+\eta)q}.$$

We conclude that when $t - s > \epsilon$, then $E_{2,\epsilon} \geq c\epsilon^{1/2} - \tilde{J}_\epsilon$, where $\mathbb{E}[|\tilde{J}_\epsilon|^q] \leq c(q)\epsilon^{(\frac{1}{2}+\eta)q}$. Therefore, when $t - s > \epsilon$,

$$\inf_{1 \geq \alpha \geq \alpha_0} \prod_{i=1}^d (\xi^i)^\top \gamma_Z \xi^i \geq \min \left(c\rho^2\epsilon^{\frac{1}{2}+\eta} - I_{3,\epsilon}, c\epsilon^{\frac{1}{2}} - \tilde{J}_\epsilon \right).$$

Putting together the results of Case 1 and Case 2, we see that for $0 < \epsilon \leq \frac{1}{2}$,

$$\inf_{\|\xi\|=1, 1 \geq \alpha \geq \alpha_0} \prod_{i=1}^d (\xi^i)^\top \gamma_Z \xi^i \geq \min \left(c\rho^2 \epsilon^{\frac{1}{2} + \eta} - I_{3,\epsilon}, c\epsilon^{\frac{1}{2}} - 2I_\epsilon \mathbf{1}_{\{\epsilon \geq t-s\}} - \tilde{J}_\epsilon \mathbf{1}_{\{\epsilon < t-s\}} \right).$$

We take into account the bounds on moments of $I_{3,\epsilon}$, I_ϵ and \tilde{J}_ϵ , and then use Proposition 3.5 to conclude the proof of Proposition 6.15. \square

7 Appendix

On several occasions, we have appealed to the following technical estimates on the Green kernel of the heat equation.

Lemma 7.1. [BP98, (A.1)] *There exists $C > 0$ such that for any $0 < s < t$ and $x, y \in [0, 1]$, $x \neq y$,*

$$G_{t-s}(x, y) \leq C \frac{1}{\sqrt{2\pi(t-s)}} \exp\left(-\frac{|x-y|^2}{2(t-s)}\right).$$

Lemma 7.2. [BP98, (A.3)] *There exists $C > 0$ such that for any $t \geq \epsilon > 0$ and $x \in [0, 1]$,*

$$\int_{t-\epsilon}^t \int_{x-\sqrt{\epsilon}}^{x+\sqrt{\epsilon}} G_{t-s}^2(x, y) dy ds \geq C\sqrt{\epsilon}.$$

Lemma 7.3. [BP98, (A.5)] *There exists $C > 0$ such that for any $\epsilon > 0$, $q < \frac{3}{2}$, $t \geq \epsilon$ and $x \in [0, 1]$,*

$$\int_{t-\epsilon}^t \int_0^1 G_{t-s}^{2q}(x, y) dy ds \leq C\epsilon^{3/2-q}.$$

Lemma 7.4. *There exists $C > 0$ such that for all $0 < a < b$ and $x \in [0, 1]$,*

$$\int_a^b \int_0^1 G_s^2(x, y) dy ds \leq C \frac{b-a}{\sqrt{b} + \sqrt{a}}.$$

Proof. Using Lemma 7.1 and the change of variables $z = \frac{x-y}{\sqrt{s}}$, we see that

$$\begin{aligned} \int_a^b \int_0^1 G_s^2(x, y) dy ds &\leq C \int_a^b \int_{-\infty}^{\infty} \frac{1}{\sqrt{s}} e^{-z^2} dz ds \\ &= \tilde{C} \int_a^b \frac{1}{\sqrt{s}} ds = 2\tilde{C}(\sqrt{b} - \sqrt{a}), \end{aligned}$$

which concludes the proof. \square

The next result is a straightforward extension to $d \geq 1$ of MORIEN [M98, Lemma 4.2] for $d = 1$.

Lemma 7.5. *Assume P1. For all $q \geq 1$, $T > 0$ there exists $C > 0$ such that for all $T \geq t \geq s \geq \epsilon > 0$ and $0 \leq y \leq 1$,*

$$\sum_{k=1}^d \mathbb{E} \left[\left(\int_{s-\epsilon}^s dr \int_0^1 dv \left| \sum_{i=1}^d D_{r,v}^{(k)}(u_i(t, y)) \right|^2 \right)^q \right] \leq C\epsilon^{q/2}.$$

The next result is Burkholder's inequality for Hilbert-space-valued martingales.

Lemma 7.6. [BP98, eq.(4.18)] *Let $H_{s,t}$ be a predictable $L^2([0, t] \times [0, 1]^m)$ -valued process, $m \geq 1$. Then, for any $p > 1$, there exists $C > 0$ such that*

$$\begin{aligned} & \mathbb{E} \left[\left| \int_{([0,t] \times [0,1])^m} \left(\int_0^t \int_0^1 H_{s,y}(\alpha) W(dy, ds) \right)^2 d\alpha \right|^p \right] \\ & \leq C \mathbb{E} \left[\left| \int_0^t \int_0^1 \left(\int_{([0,t] \times [0,1])^m} H_{s,y}^2(\alpha) d\alpha \right) dy ds \right|^p \right]. \end{aligned}$$

Acknowledgement. The authors thank V. Bally for several stimulating discussions.

References

- [BMS95] BALLY, V., MILLET, A. AND SANZ-SOLÉ, M. (1995), Approximation and support theorem in Hölder norm for parabolic stochastic partial differential equations, *The Annals of Probability*, **23**, 178-222.
- [BP98] BALLY, V. AND PARDOUX, E. (1998), Malliavin calculus for white noise driven parabolic SPDEs, *Potential Analysis*, **9**, 27-64.
- [DN04] DALANG, R.C. AND NUALART, E. (2004), Potential theory for hyperbolic SPDEs, *The Annals of Probability*, **32**, 2099-2148.
- [DKN07] DALANG, R.C., KHOSHNEVISAN, D. AND NUALART, E. (2007), Hitting probabilities for systems of non-linear stochastic heat equations with additive noise, *Submitted for publication. See <http://arxiv.org/pdf/math.PR/0702710>*.
- [K85] KAHANE, J.-P. (1985), *Some random series of functions*, Cambridge University Press.
- [K02] KHOSHNEVISAN, D. (2002), *Multiparameter processes. An introduction to random fields*, Springer-Verlag.
- [K03] KOHATSU-HIGA, A. (2003), Lower bound estimates for densities of uniformly elliptic random variables on Wiener space, *Probab. Theory Related Fields*, **126**, 421-457.
- [M98] MORIEN, P.-L. (1998), The Hölder and the Besov regularity of the density for the solution of a parabolic stochastic partial differential equation, *Bernoulli*, **5**, 275-298.

- [MT03] MUELLER, C. AND TRIBE, R. (2002), Hitting properties of the random string, *Electronic Journal of Probability*, **7**, 1-29.
- [N94] NUALART, D. AND PARDOUX, E. (1994), Markov field properties of solutions of white noise driven quasi-linear parabolic PDEs, *Stochastics and Stochastics Reports*, **48**, 17-44.
- [N95] NUALART, D. (1995), *The Malliavin calculus and related topics*, Springer-Verlag.
- [N98] NUALART, D. (1998), Analysis on Wiener space and anticipating stochastic calculus, *Ecole d'Eté de Probabilités de Saint-Flour XXV, Lect. Notes in Math.* **1690**, Springer-Verlag, 123-227.
- [S05] SANZ-SOLÉ, M. (2005), *Malliavin calculus with applications to stochastic partial differential equations*, EPFL Press.
- [W86] WALSH, J.B. (1986), An Introduction to Stochastic Partial Differential Equations, *Ecole d'Eté de Probabilités de Saint-Flour XIV, Lect. Notes in Math.*, **1180**, Springer-Verlag, 266-437.
- [W84] WATANABE, S. (1984), Lectures on Stochastic Differential Equations and Malliavin Calculus, *Tata Institute of Fundamental Research Lectures on Math. and Physics*, **73**, Springer-Verlag, Berlin.