

Level sets of the stochastic wave equation driven by a symmetric Lévy noise

Davar Khoshnevisan^{1,3} and Eulalia Nualart²

Abstract

We consider the solution $\{u(t, x), t \geq 0, x \in \mathbb{R}\}$ of a system of d linear stochastic wave equations driven by a d dimensional symmetric space-time Lévy noise. We provide a necessary and sufficient condition, on the characteristic exponent of the Lévy noise, which describes exactly when the zero set of u is nonvoid. We also compute the Hausdorff dimension of that zero set, when it is nonempty. These results will follow from more general potential-theoretic theorems on level sets of Lévy sheets.

AMS 2000 subject classifications: Primary: 60G60, 60H15; Secondary: 60J45, 60G17.

Key words and phrases. Level sets, stochastic wave equation, Lévy noise, potential theory.

¹Department of Mathematics, The University of Utah, 155 S. 1400 E. Salt Lake City, UT 84112-0090, USA. davar@math.utah.edu, <http://www.math.utah.edu/~davar>

²Institut Galilée, Université Paris 13, 93430 Villetaneuse, France, nualart@math.univ-paris13.fr, <http://www.math.univ-paris13.fr/~nualart>

³Research supported in part by the National Science Foundation

1 Introduction and main results

Consider the solution $u(t, x) := (u_1(t, x), \dots, u_d(t, x))$ to the following system of d linear stochastic wave equations:

$$\frac{\partial^2 u_i}{\partial t^2}(t, x) = \frac{\partial^2 u_i}{\partial x^2}(t, x) + \dot{L}_i(t, x), \quad t \geq 0, x \in \mathbb{R}, \quad (1.1)$$

with initial condition $u_i(0, x) = \partial_t u_i(0, x) = 0$, where i ranges in $\{1, \dots, d\}$, and $\dot{L} := (\dot{L}_1, \dots, \dot{L}_d)$ is a [totally scattered] d -dimensional Lévy noise on \mathbb{R}^2 with Lévy exponent Ψ .

Stochastic PDEs (SPDEs) that are driven by [non-Gaussian] Lévy noises are beginning to receive some attention in the literature. We mention, for example, the work of MUELLER [M98], who investigates nonlinear heat equations in high space dimensions that are driven by multiplicative space-time stable Lévy noises. See also MUELLER, MYTNIK, AND STAN [MMS06] who consider heat equations with a multiplicative space-dependent stable Lévy noise. And the Cauchy problem, in high space dimensions, for the linear wave equation driven by a space-time Lévy noise is treated in ØKSENDAL, PROSKE, AND SIGNAHL [ØPS06].

We now return to the SPDE (1.1), and define the *light cone* at $(t, x) \in \mathbb{R}_+^2$ to be the set

$$\mathcal{C}(t, x) := \{(s, y) \in \mathbb{R}_+^2 : 0 \leq s \leq t \text{ and } x - (t - s) \leq y \leq x + (t - s)\}. \quad (1.2)$$

Then, $\frac{1}{2}\mathbf{1}_{\mathcal{C}(t, x)}$ is the Green function for the wave operator. We follow the approach developed by WALSH [W86], for space-time white noise, and define the solution to the “formal” equation (1.1) as

$$u(t, x) = \frac{1}{2} \int_{\mathbb{R}_+^2} \mathbf{1}_{\mathcal{C}(t, x)}(s, y) L(ds dy) \quad \text{for all } t \geq 0, x \in \mathbb{R}, \quad (1.3)$$

where L denotes the Lévy sheet that corresponds to $\dot{L} := (\dot{L}_1, \dots, \dot{L}_d)$ [see §2.1], and the stochastic integral is a Wiener-type integral with respect to L [see §2.3].

The aim of this paper is to study the geometry of the zero set $u^{-1}\{0\}$ of the solution to (1.1). In particular, we seek to know when $u^{-1}\{0\}$ is nonvoid. In order to describe when $u^{-1}\{0\} \neq \emptyset$, we first analyze the zero set of a Lévy sheet. This will be done in a more general setting. Then, we devise a “comparison principle” to relate the solution to the SPDE (1.1) to theorems about additive Lévy process developed earlier in [KX02, KX04, KSX06]. This comparison method might appear to be a little round-about, but as it turns out, it is not so easy to work directly with the solution to (1.1).

In order to state the main result of this paper we introduce the following regularity conditions on the characteristic exponent Ψ of the noise \dot{L} ; see (2.1) below for a precise definition of Ψ :

A1. \dot{L} is *symmetric*. That is, $\Psi(\xi)$ is real and nonnegative for all $\xi \in \mathbb{R}^d$.

A2. $\Phi(\lambda) < \infty$ for all $\lambda > 0$, where Φ is the *gauge function* defined by

$$\Phi(\lambda) := \frac{1}{(2\pi)^d} \int_{\mathbb{R}^d} e^{-\lambda \Psi(\xi)} d\xi \quad \text{for all } \lambda > 0. \quad (1.4)$$

A3. For all $a > 0$ there exists a constant $A_a > 0$ such that

$$\Psi(a\xi) \geq A_a \Psi(\xi) \quad \text{for all } \xi \in \mathbb{R}^d. \quad (1.5)$$

Remark 1.1. In order to understand condition **A3** better, we note that Φ is nonincreasing. Therefore, in particular, $\Phi(2\lambda) \leq \Phi(\lambda)$ for all $\lambda > 0$. Condition **A3** implies that a converse holds. Namely,

$$\limsup_{\lambda \downarrow 0} \frac{\Phi(\lambda)}{\Phi(2\lambda)} < \infty. \quad (1.6)$$

Unfortunately, (1.6) by itself does not seem to be enough to imply our main result. \square

Given an analytic Euclidean set A , we let $\dim_{\text{H}} A$ denote its Hausdorff dimension [K02, Appendix C], with the proviso added that the statement “ $\dim_{\text{H}} A < 0$ ” means that “ A is empty.”

We are ready to present the main result of this paper; the remainder of the article is dedicated to proving this fact.

Theorem 1.2. *Under conditions **A1**–**A3**, the following are equivalent:*

- (i) *Almost surely, $u(t, x) = 0$ for some $t > 0$ and $x \in \mathbb{R}$;*
- (ii) *with positive probability, $u(t, x) = 0$ for some $t > 0$ and $x \in \mathbb{R}$;*
- (iii) $\int_0^1 \lambda \Phi(\lambda) d\lambda < \infty$.

In addition, if one of these conditions holds, then $\dim_{\text{H}} u^{-1}\{0\} = 2 - \overline{\text{ind}} \Phi$ a.s., where

$$\overline{\text{ind}} \Phi := \limsup_{\lambda \downarrow 0} \frac{\log \Phi(\lambda)}{\log(1/\lambda)}. \quad (1.7)$$

In order to understand Theorem 1.2 better, let us emphasize the special case where $\Psi(\xi) = \frac{1}{2} \|\xi\|^\alpha$ for all $\xi \in \mathbb{R}^d$, and $\alpha \in (0, 2]$ is a fixed “index of stability.” In this case, \dot{L} is called the *isotropic stable Lévy noise* with index α , and it is easy to see that $\Phi(\lambda) = \text{const} \cdot \lambda^{d/\alpha}$. Hence, Theorem 1.2 yields the following result.

Corollary 1.3. *Consider the solution u to (1.1) where \dot{L} is the isotropic stable Lévy noise with index $\alpha \in (0, 2]$. Then u has zeros if and only if $d < 2\alpha$. Moreover, if $d < 2\alpha$, then $\dim_{\text{H}} u^{-1}\{0\} = 2 - (d/\alpha)$ a.s.*

When $\alpha = 2$, \dot{L} is the standard d -dimensional white noise on \mathbb{R}^2 , and (1.1) simplifies to the more common form of the linear stochastic wave equation with d i.i.d. components. In that case, there is indeed a large literature on this topic. See, for example, DALANG [D99, D01], DALANG AND WALSH [DW92], CABAÑA [C70, C72], CARMONA AND NUALART [CN88a, CN88b], GAVEAU [G89, G95], PYASETSKAYA [P83], and WALSH [W86].

One could also consider the case where $\Psi(\xi) = \xi' \mathbf{Q} \xi$ where $\mathbf{Q} := (Q_{ij})$ is a nonsingular $d \times d$ covariance matrix. This leads to the weakly-interacting system

$$\frac{\partial^2 u_i}{\partial t^2}(t, x) = \frac{\partial^2 u_i}{\partial x^2}(t, x) + \sum_{j=1}^d Q_{ij} \dot{W}_j(t, x), \quad t \geq 0, x \in \mathbb{R}, 1 \leq i \leq d, \quad (1.8)$$

with initial condition $u_i(x, 0) = \partial_t u_i(x, 0) = 0$. Here, $\dot{W} := (\dot{W}_1, \dots, \dot{W}_d)$ denotes a standard d -dimensional white noise on \mathbb{R}^2 . Corollary 1.3 holds in this case with $\alpha = 2$, with the end-result being that the solution has zeros if and only if $d < 4$. This result is related closely to a hyperbolic problem that was solved in DALANG AND NUALART [DN04]. If we apply their characterization to the problem of the existence of zeros, then we obtain the same critical dimension of four as we do here. We add that the theorem of DALANG AND NUALART [DN04] holds in the presence of smooth multiplicative nonlinearities as well.

This paper is organized as follows: In §2 we recall Lévy sheets and their associated Lévy processes, and present a brief introduction to stochastic integration with respect to a general Lévy noise. Sections 3, 4 and 5 are devoted to the study of the polar sets of the zero set of a Lévy sheet and of the random field defined as a sum of a Lévy sheet and an additive Lévy process, in a general setting. We finally prove Theorem 1.2 in Section 6.

In the rest of this section we give some notation that will be used throughout. The k -dimensional Lebesgue measure on \mathbb{R}^k is denoted by λ_k . For all $x \in \mathbb{R}^k$, $\|x\| := (x_1^2 + \dots + x_k^2)^{1/2}$ denotes the ℓ^2 -norm of x , and $|x| := |x_1| + \dots + |x_k|$ its ℓ^1 -norm.

The underlying parameter space is \mathbb{R}^N , or $\mathbb{R}_+^N = [0, \infty)^N$. A typical parameter $t \in \mathbb{R}^N$ is written as $t = (t_1, \dots, t_N)$. There is a natural partial order \preceq on \mathbb{R}^N . Namely, $s \preceq t$ if and only if $s_i \leq t_i$ for all $i = 1, \dots, N$. When it is the case that $s \preceq t$, we define the interval $[s, t] = \prod_{i=1}^N [s_i, t_i]$. Finally, $s \wedge t$ denotes the N -vector whose i th coordinate is the minimum of s_i and t_i .

We denote by \hat{f} the normalized Fourier transform of $f \in L^1(\mathbb{R}^d)$ given by

$$\hat{f}(\xi) = \int_{\mathbb{R}^d} e^{ix \cdot \xi} f(x) dx \quad \text{for all } \xi \in \mathbb{R}^d. \quad (1.9)$$

2 Lévy sheets and their associated processes

In this section we study the structure of the distribution function of the noise \dot{L} , which is called a Lévy sheet. In fact, we will proceed by studying such sheets under greater generality that is needed for (1.1). This is primarily because we view Lévy sheets as fundamental objects themselves.

2.1 Lévy sheets

Let us begin by introducing some standard notation on N -parameter, d -dimensional Lévy sheets. For more detailed information, particularly in the two-parameter case, see §2 of DALANG AND WALSH [DW92].

Let \dot{L} be a totally scattered symmetric d -dimensional random measure on \mathbb{R}^N that is infinitely divisible in the following sense:

1. If A and B are disjoint Borel subsets of \mathbb{R}^N then $\dot{L}(A)$ and $\dot{L}(B)$ are independent;
2. For every Borel set $A \subset \mathbb{R}^N$ of finite Lebesgue measure, the law of $\dot{L}(A)$ is described by the following:

$$\mathbb{E}[\exp\{i\xi \cdot \dot{L}(A)\}] = \exp(-\lambda_N(A)\Psi(\xi)) \quad \text{for all } \xi \in \mathbb{R}^d, \quad (2.1)$$

where Ψ is a real (and hence non-negative) negative-definite function in the sense of SCHOENBERG [S37]. See also the following Remark.

Remark 2.1. We recall Schoenberg's theorem: Ψ is *negative definite* if and only if $\Psi(0) \geq 0$ and $\xi \mapsto \exp(-t\Psi(\xi))$ is positive definite in the sense of Herzog and Bochner [BF75, Theorem 7.8, p. 41]. Equivalently, Ψ is negative definite if it satisfies the Lévy–Khintchine formula [B96, S99]. In the present case, Ψ is also real-valued, and therefore, $\Psi(\xi) \geq 0$ for all $\xi \in \mathbb{R}^d$. \square

The *Lévy sheet* L with Lévy exponent Ψ is the N -parameter d -dimensional random field $\{L(t); t \in \mathbb{R}_+^N\}$ defined as

$$L(t) := \dot{L}([0, t]) \quad \text{for all } t := (t_1, \dots, t_N) \in \mathbb{R}_+^N. \quad (2.2)$$

Next we mention a few commonly-used families of Lévy noises and their corresponding Lévy sheets.

Example 2.2 (White and Stable Noises). Choose and fix a constant $\chi \in (0, \infty)$. When $\Psi(\xi) = \chi\|\xi\|^2$ for all $\xi \in \mathbb{R}^d$, \dot{L} is called the d -dimensional *white noise* on \mathbb{R}^N , and L is called the N -parameter, d -dimensional *Brownian sheet*. The noise \dot{L} is the [usual] *standard white noise* when $\chi = 1/2$, and the random field L is then called the [usual] *standard Brownian sheet*.

More generally, if $\Psi(\xi) = \chi\|\xi\|^\alpha$ for $\xi \in \mathbb{R}^d$, then by the Lévy–Khintchine formula $\alpha \in (0, 2]$, and \dot{L} is called the d -dimensional *isotropic stable noise* on \mathbb{R}^N with index α . In this case, L is called the N -parameter *isotropic stable sheet* in \mathbb{R}^d with index α ; see EHM [E81].

There are other interesting stable noises that are symmetric. For instance, one could consider $\Psi(\xi) = \frac{1}{2} \sum_{j=1}^d |\xi_j|^\alpha$. This gives rise to the symmetric noise each of whose d components are i.i.d. one-dimensional stable noises on \mathbb{R}^N with common index α . The resulting random field is a stable sheet with i.i.d. coordinates, each with the same stability index α . \square

Example 2.3 (Noises with Stable Components). Let $\alpha_1, \dots, \alpha_d \in (0, 2]$ be fixed, and consider $\Psi(\xi) = \chi \sum_{j=1}^d |\xi_j|^{\alpha_j}$, where $\chi \in (0, \infty)$ is fixed. The resulting random noise \dot{L} has *stable components* with *index* $(\alpha_1, \dots, \alpha_d)$, and the corresponding random field L is the *stable sheet with stable components*. These were introduced in the one-parameter case by PRUITT [PT69], and investigated further by HENDRICKS [H70, H72, H73, H74]. \square

Let us mention also the following ready consequence of Theorem 1.2.

Corollary 2.4. Consider the solution u to (1.1) where \dot{L} is Lévy noise with stable components with index $(\alpha_1, \dots, \alpha_d) \in (0, 2]^d$. Then, u has zeros if and only if $\sum_{j=1}^d (1/\alpha_j) < 2$. If and when $\sum_{j=1}^d (1/\alpha_j) < 2$, then $\dim_{\mathbb{H}} u^{-1}\{0\} = 2 - \sum_{j=1}^d (1/\alpha_j)$ a.s.

2.2 The associated processes

Because Ψ is a Lévy exponent there exists a d -dimensional Lévy process $X := \{X(t); t \geq 0\}$ whose law is described uniquely by $E[e^{i\xi \cdot X(t)}] = e^{-t\Psi(\xi)}$ for all $t \geq 0$ and $\xi \in \mathbb{R}^d$. We may refer to X as the *Lévy process associated to \dot{L}* .

By the inversion theorem, the transition densities of X are given by

$$f(t; x) = \frac{1}{(2\pi)^d} \int_{\mathbb{R}^d} \cos(x \cdot \xi) e^{-t\Psi(\xi)} d\xi \quad \text{for all } t \geq 0, x \in \mathbb{R}^d. \quad (2.3)$$

In particular, $\Phi(\lambda)$ is none other than $f(\lambda; 0)$.

Now let X_1, \dots, X_N be N i.i.d. copies of X . Then *additive Lévy process associated to \dot{L}* is defined as the d -dimensional N -parameter random field $\mathfrak{X} := \{\mathfrak{X}(t); t \in \mathbb{R}_+^N\}$, where

$$\mathfrak{X}(t) := X_1(t_1) + \dots + X_N(t_N) \quad \text{for all } t \in \mathbb{R}_+^N. \quad (2.4)$$

The density function of $\mathfrak{X}(t)$ at $0 \in \mathbb{R}^d$ is $\Phi(\sum_{i=1}^N t_i)$. This should be contrasted with the fact that the density function of $L(t)$ at $0 \in \mathbb{R}^d$ is $\Phi(\prod_{i=1}^N t_i)$.

2.3 Stochastic integrals

In this section we proceed to the construct Wiener-type stochastic integrals of the type $\dot{L}(\varphi) := \int_{\mathbb{R}^N} \varphi(t) \dot{L}(dt)$, where $\varphi : \mathbb{R}^N \rightarrow \mathbb{R}$ is nonrandom, measurable, bounded, and compactly supported. There are in place integration theories that construct $\dot{L}(\varphi)$ abstractly; see, for example the Bartle-type integrals of DUNFORD AND SCHWARTZ [DS58]. But in order to mention some of the probabilistic properties of these integrals, we opt for a more direct approach that is closer to the original method of Wiener. As our method uses standard ideas to produce such stochastic integrals, we will only sketch the main steps.

First suppose φ is a simple function. That is, $\varphi(t) = \sum_{j=1}^m c_j \mathbf{1}_{A_j}$, where A_1, \dots, A_m are disjoint Borel sets in \mathbb{R}^N , and c_1, \dots, c_m are real constants. For such φ we define $\dot{L}(\varphi) := \sum_{j=1}^m c_j \dot{L}(A_j)$. Because $\Psi(0) = 0$,

$$E[\exp\{i\xi \cdot \dot{L}(\varphi)\}] = \exp\left(-\sum_{j=1}^m \lambda^N(A_j) \Psi(c_j \xi)\right) = \exp\left(-\int_{\mathbb{R}^N} \Psi(\varphi(t) \xi) dt\right). \quad (2.5)$$

Next we consider a bounded, compactly-supported measurable function $\varphi : \mathbb{R}^N \rightarrow \mathbb{R}$. Standard measure-theoretic arguments reveal that we can find simple functions $\varphi_1, \varphi_2, \dots : \mathbb{R}^N \rightarrow \mathbb{R}_+$ and a compact set $K \subset \mathbb{R}^N$ such that: (i) $\lim_{n \rightarrow \infty} \varphi_n = \varphi$ pointwise; (ii) $R := \sup_{n \geq 1} \sup_{t \in \mathbb{R}^N} |\varphi_n(t)| < \infty$; and (iii) $\varphi_n(t) = 0$, for all $t \notin K$ and all $n \geq 1$.

By linearity, and since Ψ is real and nonnegative,

$$\begin{aligned} 1 &\geq \liminf_{n, m \rightarrow \infty} E\left[\exp\left(i\xi \cdot \left\{\dot{L}(\varphi_n) - \dot{L}(\varphi_m)\right\}\right)\right] \\ &\geq \exp\left(-\limsup_{n, m \rightarrow \infty} \int_K \Psi(\{\varphi_n(t) - \varphi_m(t)\} \xi) dt\right). \end{aligned} \quad (2.6)$$

Let $c := R \max_{1 \leq j \leq d} |\xi_j|$, and note that Ψ is uniformly continuous on $[-c, c]^d$. Therefore, $\lim_{n, m \rightarrow \infty} \mathbb{E}[\exp(i\xi \cdot \{\dot{L}(\varphi_n) - \dot{L}(\varphi_m)\})] = 1$. Consequently, $\{\dot{L}(\varphi_n)\}_{n \geq 1}$ is a Cauchy sequence in $L^0(\mathbb{P})$, and hence $\dot{L}(\varphi_n)$ converges in probability. The limit is denoted by $\dot{L}(\varphi)$ and has the following properties. As they can be proved directly from the preceding construction, we list them below without proof.

Proposition 2.5. *Let \mathcal{B} denote the algebra of all nonrandom, bounded, compactly supported measurable functions from \mathbb{R}^N to \mathbb{R} . Then there is an “iso-Lévy” process $\{\dot{L}(\varphi)\}_{\varphi \in \mathcal{B}}$ with the following properties:*

- (i) $\mathbb{E}[\exp\{i\xi \cdot \dot{L}(\varphi)\}] = \exp(-\int_{\mathbb{R}^N} \Psi(\varphi(t)\xi) dt)$ for all $\xi \in \mathbb{R}^d$ and $\varphi \in \mathcal{B}$.
- (ii) If $\varphi_1, \varphi_2 \in \mathcal{B}$, then $\dot{L}(\varphi_1 + \varphi_2) = \dot{L}(\varphi_1) + \dot{L}(\varphi_2)$ a.s.
- (iii) If $a \in \mathbb{R}$ and $\varphi \in \mathcal{B}$, then $\dot{L}(a\varphi) = a\dot{L}(\varphi)$ a.s.

That is, \dot{L} defines a *random linear functional* in the sense of MINLOS [M58, M59]. Alternatively, one might write $\int \varphi dL$ in place of $\dot{L}(\varphi)$.

3 The image of a Lévy sheet

Theorem 1.2 is a consequence of the more general results of this section. Before we describe them we need to introduce some notation.

Throughout the rest of the paper $\{L(t); t \in \mathbb{R}_+^N\}$ denotes a d -dimensional N -parameter Lévy sheet whose Lévy exponent Ψ satisfies the symmetry condition **A1**, and whose gauge function Φ satisfies the regularity conditions **A2** and **A3**. Occasionally we need the corresponding Lévy noise, which we denote by \dot{L} . Note that the Lévy sheet that arises from the SPDE (1.1) is one such random field with $N = 2$.

We assume that the underlying sample space Ω is the collection of all cadlag functions $\omega : \mathbb{R}_+^N \mapsto \mathbb{R}^d$. Thus, $\omega \in \Omega$ if and only if:

1. For any net of elements $\{t_\alpha\}_{\alpha \in \mathbb{R}}$ in \mathbb{R}_+^N such that $t_\alpha \preceq t_\beta$ whenever $\alpha \leq \beta$, and t_α converges to some $t \in \mathbb{R}_+^N$ as $\alpha \rightarrow \infty$, then $\lim_{\alpha \rightarrow \infty} \omega(t_\alpha) = \omega(t)$; and
2. for any net of elements $\{t_\alpha\}_{\alpha \in \mathbb{R}}$ in \mathbb{R}_+^N such that $t_\alpha \preceq t_\beta$ whenever $\alpha \geq \beta$, then $\lim_{\alpha \rightarrow \infty} \omega(t_\alpha)$ exists.

We say that the Lévy process L is in *canonical form* if

$$L(t)(\omega) = \omega(t) \quad \text{for all } \omega \in \Omega \text{ and } t \in \mathbb{R}_+^N. \quad (3.1)$$

Throughout, we will assume that our Lévy process L is in canonical form under a fixed probability measure \mathbb{P} . This assumption is made tacitly, and does not incur any loss in generality.

Define \mathbb{P}_x to be the law of the process $x + L$ for every $x \in \mathbb{R}^d$. That is, for every Borel subset A of Ω ,

$$\mathbb{P}_x\{\omega \in \Omega : \omega \in A\} = \mathbb{P}\{\omega \in \Omega : x + \omega \in A\}. \quad (3.2)$$

Let E_x denote the corresponding expectation operator. We will be primarily interested in the sigma-finite (but infinite) measure,

$$P_{\lambda_d}(\bullet) := \int_{\mathbb{R}^d} P_x(\bullet) dx. \quad (3.3)$$

We will write E_{λ_d} for the corresponding “expectation operator.” That is,

$$E_{\lambda_d}(Z) := \int_{\Omega} Z(\omega) P_{\lambda_d}(d\omega) = \int_{\mathbb{R}^d} E_x(Z) dx \quad \text{for all } Z \in L^1(P_{\lambda_d}). \quad (3.4)$$

Let $\mathcal{P}(G)$ denote the collection of all probability measures on any Euclidean set G , and define for all $\mu \in \mathcal{P}(\mathbb{R}_+^N)$,

$$I(\mu) := \iint_{\{s, t \in \mathbb{R}_+^N: s \neq t\}} \Phi(\|s - t\|) \mu(dt) \mu(ds). \quad (3.5)$$

For all compact sets $G \subset \mathbb{R}_+^N$, we consider also the *capacity* of G ,

$$\text{Cap}(G) := \left[\inf_{\mu \in \mathcal{P}(G)} I(\mu) \right]^{-1}, \quad (3.6)$$

where $\inf \emptyset := \infty$, and $1/\infty := 0$. Let $L(G) := \cup_{t \in G} \{L(t)\}$ denote the range of G under the random map $t \mapsto L(t)$. The following is the main result of this section.

Theorem 3.1. *Let G be a nonrandom compact subset of $(0, \infty)^N$. Then there exists a positive and finite constant $c = c(G)$ such that*

$$c^{-1} \text{Cap}(G) \leq P_{\lambda_d} \{0 \in L(G)\} = E [\lambda_d(L(G))] \leq c \text{Cap}(G). \quad (3.7)$$

Moreover, c depends on G only through $\inf_{x \in G} |x|$ and $\sup_{x \in G} |x|$.

As a consequence of this theorem and Theorem 5.1 of KHOSHNEVISAN AND XIAO [KX03], we have the following equivalence theorem between the Lévy sheets of this paper and their associated additive Lévy processes.

Corollary 3.2. *Let $\{\mathfrak{X}(t); t \in \mathbb{R}_+^N\}$ denote the additive Lévy process associated to L . Then for all nonrandom compact sets G in \mathbb{R}_+^N there exists a positive and finite constant $c = c(G)$ such that*

$$c^{-1} E [\lambda_d(\mathfrak{X}(G))] \leq E [\lambda_d(L(G))] \leq c E [\lambda_d(\mathfrak{X}(G))]. \quad (3.8)$$

Moreover, c depends only on G through $\inf_{x \in G} |x|$ and $\sup_{x \in G} |x|$.

Example 3.3. Suppose L is an isotropic α -stable Lévy sheet. Then, we can either calculate directly, or apply the preceding together with Corollary 5.4 of KHOSHNEVISAN AND XIAO [KX03], to find that

$$E [\lambda_d(L(G))] > 0 \iff \text{Cap}(G) > 0 \iff \mathcal{C}_{d/\alpha}(G) > 0, \quad (3.9)$$

where \mathcal{C}_β denotes the standard β -dimensional Riesz capacity. See Appendix D of KHOSHNEVISAN [K02] for information on these capacities. \square

Example 3.4. Suppose L is a Lévy sheet with stable components of index $(\alpha_1, \dots, \alpha_d) \in [0, 2]^d$. Then,

$$\mathbb{E} [\lambda_d(L(G))] > 0 \iff \text{Cap}(G) > 0 \iff \mathcal{C}_{\sum_{j=1}^d (1/\alpha_j)}(G) > 0, \quad (3.10)$$

where \mathcal{C}_β is as in the previous example. \square

Our proof of Theorem 3.1 proceeds by first establishing an elementary lemma.

Lemma 3.5. For all $f \in L^1(\mathbb{R}^d)$ and $t \in \mathbb{R}_+^N$,

$$\mathbb{E}_{\lambda_d} [f(L(t))] = \int_{\mathbb{R}^d} f(x) dx. \quad (3.11)$$

Proof. Because f is measurable and integrable, Fubini's theorem applies, and we have

$$\mathbb{E}_{\lambda_d} [f(L(t))] = \mathbb{E} \left[\int_{\mathbb{R}^d} f(x + L(t)) dx \right], \quad (3.12)$$

and this is manifestly equal to $\int_{\mathbb{R}^d} f(x) dx$. \square

The next lemma is nearly as simple, but particularly useful to us.

Lemma 3.6. If f and g are two probability densities on \mathbb{R}^d such that $\hat{f}, \hat{g} \in L^1(\mathbb{R}^d)$, then

$$\mathbb{E}_{\lambda_d} [f(L(t))g(L(s))] = \frac{1}{(2\pi)^d} \int_{\mathbb{R}^d} e^{-\ell(s,t)\Psi(\xi)} \hat{g}(\xi) \overline{\hat{f}(\xi)} d\xi, \quad (3.13)$$

where ℓ is the symmetric function:

$$\ell(s, t) := \lambda_N([0, s] \triangle [0, t]) \quad \text{for all } s, t \in \mathbb{R}_+^N. \quad (3.14)$$

Proof. We write

$$\begin{aligned} \mathbb{E}_{\lambda_d} [f(L(t))g(L(s))] &= \mathbb{E} \left[\int_{\mathbb{R}^d} f(x + L(t)) g(x + L(s)) dx \right] \\ &= \int_{\mathbb{R}^d} f(y) \mathbb{E} [g(y + L(s) - L(t))] dy \\ &= \int_{\mathbb{R}^d} f(y) \int_{\mathbb{R}^d} g(y - z) \mu_{s,t}(dz) dy, \end{aligned} \quad (3.15)$$

where $\mu_{s,t}$ denotes the distribution of $L(t) - L(s)$. By the inversion theorem of Fourier analysis,

$$\begin{aligned} \mathbb{E}_{\lambda_d} [f(L(s))g(L(t))] &= \frac{1}{(2\pi)^d} \int_{\mathbb{R}^d} f(y) \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} e^{-i(y-z)\cdot\xi} \hat{g}(\xi) d\xi \mu_{s,t}(dz) dy \\ &= \frac{1}{(2\pi)^d} \int_{\mathbb{R}^d} \hat{\mu}_{s,t}(\xi) \hat{g}(\xi) \overline{\hat{f}(\xi)} d\xi. \end{aligned} \quad (3.16)$$

In order to compute the Fourier transform of $\mu_{s,t}$ we first note that

$$\begin{aligned} L(t) - L(s) &= \dot{L}([0, t]) - \dot{L}([0, s]) \\ &= \dot{L}([0, t] \setminus [0, s]) - \dot{L}([0, s] \setminus [0, t]). \end{aligned} \quad (3.17)$$

The last two random variables are independent from one another. Moreover, by symmetry, $-\dot{L}([0, s] \setminus [0, t])$ has the same distribution as $\dot{L}([0, s] \setminus [0, t])$. Therefore, $L(t) - L(s)$ has the same distribution as $\dot{L}([0, s] \triangle [0, t])$. From this and (2.1) it follows that $\hat{\mu}_{s,t}(\xi) = \exp(-\ell(s, t)\Psi(\xi))$. We plug this in to (3.16), in order to conclude the proof. \square

Next we recall a well-known estimate for the function ℓ that was defined in (3.14); see [K02, Lemma 1.3.1, p. 460] for a proof.

Lemma 3.7. *If $u \preceq v$ are in $(0, \infty)^N$, then there exist positive and finite constants a and b such that*

$$a\|s - t\| \leq \ell(s, t) \leq b\|s - t\| \quad \text{for all } s, t \in [u, v]. \quad (3.18)$$

Lemma 3.8. *If $G \subset \mathbb{R}_+^N$ is compact and nonrandom, then*

$$\mathbb{P}_{\lambda_d} \left\{ 0 \in \overline{L(G)} \right\} \geq \text{const} \cdot \text{Cap}(G). \quad (3.19)$$

Moreover, the constant depends on G only through $\inf_{x \in G} |x|$ and $\sup_{x \in G} |x|$.

Proof. Without loss of generality, we assume that $\text{Cap}(G) > 0$; otherwise there is nothing to prove. In that case, there exists $\mu \in \mathcal{P}(G)$ such that $I(\mu) < \infty$. Because $\Phi(0)$ is infinite, for any such probability measure μ , it holds that

$$I(\mu) = \iint_{\mathbb{R}_+^N \times \mathbb{R}_+^N} \Phi(\|s - t\|) \mu(ds) \mu(dt). \quad (3.20)$$

For all $\mu \in \mathcal{P}(G)$ and all f probability density function on \mathbb{R}^d , define the random variable

$$J(f; \mu) := \int_{\mathbb{R}_+^N} f(L(t)) \mu(dt). \quad (3.21)$$

By Lemma 3.5, $\mathbb{E}_{\lambda_d}[J(f)] = 1$. On the other hand, by Lemma 3.6,

$$\begin{aligned} \mathbb{E}_{\lambda_d}(|J(f; \mu)|^2) &= \frac{1}{(2\pi)^d} \iint_{\mathbb{R}_+^N \times \mathbb{R}_+^N} \int_{\mathbb{R}^d} e^{-\ell(s,t)\Psi(\xi)} |\hat{f}(\xi)|^2 d\xi \mu(ds) \mu(dt) \\ &\leq \frac{1}{(2\pi)^d} \iint_{\mathbb{R}_+^N \times \mathbb{R}_+^N} \int_{\mathbb{R}^d} e^{-\ell(s,t)\Psi(\xi)} d\xi \mu(ds) \mu(dt). \end{aligned} \quad (3.22)$$

Now we apply Lemma 3.7 to see that there exists a positive and finite constant a — depending on G only through $\inf_{x \in G} |x|$ and $\sup_{x \in G} |x|$ — such that

$$\mathbb{E}_{\lambda_d}(|J(f; \mu)|^2) \leq \frac{1}{(2\pi)^d} \iint_{\mathbb{R}_+^N \times \mathbb{R}_+^N} \Phi(a\|s - t\|) \mu(ds) \mu(dt) \leq A_a I(\mu). \quad (3.23)$$

The second bound follows from **A3** and (3.20).

Now, the Paley–Zygmund inequality [K02, Lemma 1.4.1, Chap. 3] can be applied to the σ -finite measure P_{λ_d} , and this implies that

$$P_{\lambda_d}\{J(f; \mu) > 0\} \geq \frac{|(E_{\lambda_d}[J(f; \mu)])|^2}{E_{\lambda_d}(|J(f; \mu)|^2)} \geq \frac{1}{A_a I(\mu)}, \quad (3.24)$$

where $1/0 := \infty$ is allowed.

Let f be a probability density function on $B(0; 1)$, where $B(x; r)$ denotes the open (Euclidean) ball of radius $r > 0$ centered at $x \in \mathbb{R}^d$. For all $\epsilon > 0$ define $f_\epsilon(x) := \epsilon^{-d} f(x/\epsilon)$, as x varies over \mathbb{R}^d . Note that f_ϵ is a probability density on $B(0; \epsilon)$. Furthermore, if ω is such that $J(f_\epsilon; \mu)(\omega) > 0$, then there exists $t \in G$ such that $L(t) \in B(0; \epsilon)$. Hence, it follows that for all $\mu \in \mathcal{P}(G)$ such that $I(\mu) < \infty$,

$$P_{\lambda_d}\{L(G) \cap B(0; \epsilon) \neq \emptyset\} \geq \frac{1}{A_a I(\mu)}. \quad (3.25)$$

From here, we can deduce the lemma by first letting ϵ tend to zero, and then optimizing over all $\mu \in \mathcal{P}(G)$. \square

For all $t \in \mathbb{R}_+^N$, we denote by $\mathcal{F}^0(t)$ the sigma-algebra generated by $\{L(s); s \preceq t\}$. Let $\mathcal{F}^1(t)$ denote the completion of $\mathcal{F}^0(t)$ with respect to P_x for all $x \in \mathbb{R}^d$. Finally, define

$$\mathcal{F}(t) := \bigcap_{u \succeq t} \mathcal{F}^1(u). \quad (3.26)$$

In the language of the general theory of random fields, $\mathcal{F} := \{\mathcal{F}(t); t \in \mathbb{R}_+^N\}$ is the *augmented* N -parameter filtration of L .

The following is an analogue of the “Markov property,” in the present setting.

Proposition 3.9. *If $s \preceq t$ are in \mathbb{R}_+^N , then for all measurable functions $f : \mathbb{R}^d \mapsto \mathbb{R}_+$,*

$$E_{\lambda_d}[f(L(t)) \mid \mathcal{F}(s)] = (T_{s,t}^L f)(L(s)), \quad (3.27)$$

where

$$(T_{s,t}^L f)(y) := E[f(L(t) - L(s) + y)] \quad \text{for all } y \in \mathbb{R}_+^N. \quad (3.28)$$

Proof. Consider measurable functions $f, g, h_1, \dots, h_m : \mathbb{R}^d \mapsto \mathbb{R}_+$ and times $t, s, s^1, \dots, s^m \in \mathbb{R}_+^N$ such that $t \succeq s \succeq s^j$, for all $j = 1, \dots, m$. Because $\dot{L}(A)$ and $\dot{L}(B)$ are independent when $A \cap B = \emptyset$,

$$\begin{aligned} & E_{\lambda_d} \left[f(L(t)) \cdot g(L(s)) \cdot \prod_{j=1}^m h_j(L(s^j)) \right] \\ &= \int_{\mathbb{R}^d} E \left[f(L(t) + x) \cdot g(L(s) + x) \cdot \prod_{j=1}^m h_j(L(s^j) + x) \right] dx \\ &= \int_{\mathbb{R}^d} E[f(L(t) - L(s) + y)] E \left[\prod_{j=1}^m h_j(L(s^j) - L(s) + y) \right] g(y) dy. \end{aligned} \quad (3.29)$$

Set $f = h_1 = \dots = h_m \equiv 1$ to see that $P_{\lambda_d}\{L(s) \in \bullet\} = \lambda_d(\bullet)$, and the desired result follows. \square

The next result is Cairoli's maximal inequality. This inequality is proved as a consequence of the *commuting* property of the N -parameter filtration \mathcal{F} .

Proposition 3.10. *For all $Y \in L^2(P_{\lambda_d})$, there exists a version of $t \mapsto E_{\lambda_d}[Y | \mathcal{F}(t)]$ that is cadlag in every variable, uniformly in the remaining $N - 1$ variables, and satisfies*

$$E_{\lambda_d} \left(\sup_{t \in \mathbb{R}_+^N} |E_{\lambda_d}[Y | \mathcal{F}(t)]|^2 \right) \leq 4^N E_{\lambda_d}[Y^2]. \quad (3.30)$$

Proof. When L is the Brownian sheet and the infinite measure P_{λ_d} is replaced by the probability measure P , this result follows from Cairoli's maximal inequality and the Cairoli-Walsh commutation theorem. See Chapter 7 of KHOSHNEVISAN [K02, §2].

In order to prove the present form of the proposition, we need to replace the Gaussian Brownian sheet with the more general random field L , and more significantly, P with P_{λ_d} . Fortunately, none of this requires us to introduce too many new ideas. Therefore, we merely outline the requisite changes in the existing theory to accomodate the present formulation of this proposition.

We appeal to a close analogy with the usual N -parameter theory of processes, and say that \mathcal{F} is a *commuting N parameter filtration* with respect to P_{λ_d} if for all $s, t \in \mathbb{R}_+^N$ and every $\mathcal{F}(s)$ -measurable random variable $Y \in L^2(P_{\lambda_d})$,

$$E_{\lambda_d}[Y | \mathcal{F}(t)] = E_{\lambda_d}[Y | \mathcal{F}(s \wedge t)], \quad (3.31)$$

off of a P_{λ_d} -null set. In the case that $N = 2$ and P is replaced by P_{λ_d} , (3.31) reduces essentially to hypothesis (F4) of CAIROLI AND WALSH [CW75]. The general case is studied in Chapter 7 of KHOSHNEVISAN [K02, §2.1], among other places.

It is known that (3.31) implies the ‘‘Cairoli maximal inequality’’ announced in the statement of the proposition. In the case that P_{λ_d} is replaced everywhere by P , this is covered by Theorem 2.3.2 of [K02, p. 235]. It is not hard to adapt the proof so that the result continues to hold for P_{λ_d} . Thus, it suffices to derive (3.31).

Next, we prove (3.31), but only in the case that $N = 2$. When $N > 2$, the same ideas continue to work, but the notation is more cumbersome to write [and read!].

Define \mathcal{D}_n to be the collection of all left-open right-closed dyadic cubes of side length 2^{-n} in \mathbb{R}^2 . That is,

$$Q \in \mathcal{D}_n \quad \text{iff} \quad Q = \left(\frac{j}{2^n}, \frac{j+1}{2^n} \right] \times \left(\frac{k}{2^n}, \frac{k+1}{2^n} \right] \quad \text{for some } j, k \in \mathbb{Z}. \quad (3.32)$$

For all $s \in (0, \infty)^2$ and $n \geq 0$, let $\mathcal{F}_n(s)$ denote the sigma-algebra generated by all random variables of the form $\dot{L}(A)$, as A ranges over all elements of \mathcal{D}_n that are contained in $(0, s_1] \times (0, s_2]$. For every fixed integer $n \geq 1$, \mathcal{F}_n defines a two-parameter filtration. We might note that if A and B are disjoint elements of \mathcal{D}_n , then $\dot{L}(A)$ and $\dot{L}(B)$ are i.i.d. copies

of one another. Therefore, $\{\mathcal{F}_n(k/2^n, (k+1)/2^n)\}_{k=0}^\infty$ is in fact a two-parameter filtration that is defined by a two-parameter random walk [K02, Chap. 4]. Consequently, Proposition 1.2.1 of KHOSHNEVISAN [K02, p. 107] implies that for all $\mathcal{F}(s)$ -measurable random variables $Y \in L^2(P)$, the following holds with P -probability one:

$$E[Y | \mathcal{F}_n(t)] = E[Y | \mathcal{F}_n(s \wedge t)]. \quad (3.33)$$

And this is valid for all $s, t \in (0, \infty)^2$ that respectively have the form $s = (k/2^n, (k+1)/2^n)$ and $t = (j/2^n, (j+1)/2^n)$ for integers $j, k \geq 0$. Let us call such s and t *dyadic vectors of depth 2^{-n}* .

It is possible to combine the inclusion-exclusion formula [K02, p. 107] with the proof of Proposition 1.2.1 of [K02, p. 107], and derive from the preceding that for all dyadic vectors s and t of depth 2^{-n} , and all $\mathcal{F}(s)$ -measurable random variables $Y \in L^2(P_{\lambda_d})$, the following holds outside a P_{λ_d} -null set:

$$E_{\lambda_d}[Y | \mathcal{F}_n(t)] = E_{\lambda_d}[Y | \mathcal{F}_n(s \wedge t)]. \quad (3.34)$$

The nested property of dyadic cubes is inherited by the \mathcal{F}_n 's. That is, $\mathcal{F}_n(t) \subseteq \mathcal{F}_{n+1}(t)$ and $\mathcal{F}_n(s \wedge t) \subseteq \mathcal{F}_{n+1}(s \wedge t)$. Therefore, we can apply Doob's one-parameter martingale convergence theorem [which is easily adapted to work for P_{λ_d} as well] to deduce that for all dyadic vectors s and t [of any common fixed depth], the following holds off of a P_{λ_d} -null set:

$$E_{\lambda_d}\left[Y \left| \bigvee_{n=0}^{\infty} \mathcal{F}_n(t) \right.\right] = E_{\lambda_d}\left[Y \left| \bigvee_{n=0}^{\infty} \mathcal{F}_n(s \wedge t) \right.\right]. \quad (3.35)$$

We augment the sigma-algebras $\bigvee_{n=0}^{\infty} \mathcal{F}_n(s)$ and $\bigvee_{n=0}^{\infty} \mathcal{F}_n(s \wedge t)$ in the usual way to conclude from this that (3.31) holds for all fixed dyadic vectors s and t . [See the construction of \mathcal{F} via \mathcal{F}^0 , for instance.] Standard arguments involving right continuity imply that the same fact is valid for all $s, t \in (0, \infty)^2$; in the case that P_{λ_d} is replaced everywhere by P , these arguments are summarized in Corollary 2.3.1 of [K02, p. 236]. The case of P_{λ_d} is proved by very much the same arguments. This development implies (3.31), whence the proposition. \square

Lemma 3.11. *If G is a nonrandom compact set in \mathbb{R}_+^N , then*

$$P_{\lambda_d}\{0 \in L(G)\} \leq \text{const} \cdot \text{Cap}(G). \quad (3.36)$$

Moreover, the constant depends on G only through the quantities $\sup_{x \in G} |x|$ and $\inf_{x \in G} |x|$.

Proof. Without loss of generality we may assume that

$$P_{\lambda_d}\{0 \in L(G)\} > 0. \quad (3.37)$$

Otherwise there is nothing to prove.

Let f be a probability density on \mathbb{R}^d , and consider an arbitrary $\mu \in \mathcal{P}(G)$. In accord with Proposition 3.9, for all $s \in G$,

$$E_{\lambda_d}[J(f; \mu) | \mathcal{F}(s)] \geq \int_{t \succeq s} (T_{s,t}^L f)(L(s)) \mu(dt) \quad P_{\lambda_d}\text{-a.s.}, \quad (3.38)$$

where $J(f; \mu)$ is defined in (3.21). The two sides of (3.38) are right-continuous in s and have left limits, both coordinatewise. See DALANG AND WALSH [DW92], and also EHM [E81]. Therefore, for all probability densities f on \mathbb{R}^d such that $\hat{f} \in L^1(\mathbb{R}^d)$, and for all $\mu \in \mathcal{P}(G)$, the following statement holds:

$$\text{There exists one null set off which (3.38) holds for all } s \in G. \quad (3.39)$$

Path regularity of L implies the existence of a random variable σ with values in $G \cup \{\rho\}$ —where ρ is an abstract “cemetery” point not in G —which has the following properties:

1. $\sigma = \rho$ if and only if $L(s) \neq 0$ for every $s \in G$;
2. $L(\sigma) = 0$ on $\{\sigma \neq \rho\}$.

For all integers $k \geq 1$ and all Borel sets $E \subset \mathbb{R}_+^N$, define

$$\mu_{\epsilon,k}(E) := \frac{\mathbb{P}_{\lambda_d} \{\sigma \in E, \sigma \neq \rho, |L(0)| \leq k\}}{\mathbb{P}_{\lambda_d} \{\sigma \neq \rho, |L(0)| \leq k\}}. \quad (3.40)$$

Thanks to (3.37) and the monotone convergence theorem, $\lim_{k \rightarrow \infty} \mathbb{P}_{\lambda_d} \{\sigma \neq \rho, |L(0)| \leq k\} = \mathbb{P}_{\lambda_d} \{0 \in L(G)\} > 0$. Therefore, there exists $k_\epsilon > 0$ such that $\mu_{\epsilon,k}$ is well defined for all $k \geq k_\epsilon$. In fact, $\mu_{\epsilon,k}$ is a probability measure on G , provided that $k \geq k_\epsilon$.

Define for all $\epsilon > 0$, $k \geq k_\epsilon$, and all probability densities f on \mathbb{R}^d ,

$$Q_{\epsilon,k}(f) := \sup_{u \in G} \mathbb{E}_{\lambda_d} [J(f; \mu_{\epsilon,k}) | \mathcal{F}(u)]. \quad (3.41)$$

Then, thanks to (3.38) and (3.39),

$$Q_{\epsilon,k}(f) \geq \mathbf{1}_{\{\sigma \neq \rho, |L(0)| \leq k\}} \cdot \int_{t \geq \sigma} (T_{\sigma,t}^L f)(0) \mu_{\epsilon,k}(dt) \quad \mathbb{P}_{\lambda_d}\text{-a.s.}, \quad (3.42)$$

provided that $\hat{f} \in L^1(\mathbb{R}^d)$. We can square both sides and integrate $[\mathbb{P}_{\lambda_d}]$ to find that

$$\begin{aligned} & \mathbb{E}_{\lambda_d} (|Q_{\epsilon,k}(f)|^2) \\ & \geq \mathbb{P}_{\lambda_d} \{\sigma \neq \rho, |L(0)| \leq k\} \cdot \int_{\mathbb{R}_+^N} \left(\int_{t \geq s} (T_{s,t}^L f)(0) \mu_{\epsilon,k}(dt) \right)^2 \mu_{\epsilon,k}(ds). \end{aligned} \quad (3.43)$$

Since $\mu_{\epsilon,k}$ is a probability measure on G [$\epsilon > 0$ and $k \geq k_\epsilon$], the Cauchy–Schwarz inequality implies that

$$\mathbb{E}_{\lambda_d} (|Q_{\epsilon,k}(f)|^2) \geq \Lambda^2 \mathbb{P}_{\lambda_d} \{\sigma \neq \rho, |L(0)| \leq k\}, \quad (3.44)$$

where

$$\Lambda := \int_{\mathbb{R}_+^N} \int_{t \geq s} (T_{s,t}^L f)(0) \mu_{\epsilon,k}(dt) \mu_{\epsilon,k}(ds). \quad (3.45)$$

Suppose also that $\hat{f}(\xi) \geq 0$ for all $\xi \in \mathbb{R}^d$. Because $f, \hat{f} \in L^1(\mathbb{R}^d)$, can apply the inversion theorem to f —in the same manner as in the proof of Lemma 3.6—to find that for all $s, t \in G$ with $s \preceq t$, and for all $\epsilon > 0$,

$$\begin{aligned} (T_{s,t}^L f)(0) &= \frac{1}{(2\pi)^d} \int_{\mathbb{R}^d} e^{-\ell(s,t)\Psi(\xi)} \hat{f}(\xi) d\xi \\ &\geq \frac{1}{(2\pi)^d} \int_{\mathbb{R}^d} e^{-b\|s-t\|\Psi(\xi)} \hat{f}(\xi) d\xi. \end{aligned} \quad (3.46)$$

Here, b is a positive and finite constant that depends only on the distance between G and the axes of \mathbb{R}_+^N , as well as $\sup_{x \in G} |x|$; see Lemma 3.7. Consequently,

$$\Lambda \geq \frac{1}{2^N (2\pi)^d} \iint_{\mathbb{R}_+^N \times \mathbb{R}_+^N} \int_{\mathbb{R}^d} e^{-b\|s-t\|\Psi(\xi)} \hat{f}(\xi) d\xi \mu_{\epsilon,k}(ds) \mu_{\epsilon,k}(dt). \quad (3.47)$$

According to **A3**, $b\Psi(\xi) \leq \Psi(\xi/A_{1/b})$. Therefore, the preceding implies that

$$\Lambda \geq \frac{A_{1/b}^d}{2^N (2\pi)^d} \iint_{\mathbb{R}_+^N \times \mathbb{R}_+^N} \int_{\mathbb{R}^d} e^{-\|s-t\|\Psi(\xi)} \hat{f}(\xi A_{1/b}) d\xi \mu_{\epsilon,k}(ds) \mu_{\epsilon,k}(dt). \quad (3.48)$$

Now, thanks to Proposition 3.10 and (3.22),

$$\begin{aligned} \mathbb{E}_{\lambda_d} \left(|Q_{\epsilon,k}(f)|^2 \right) &\leq 4^N \mathbb{E}_{\lambda_d} \left(|J(f; \mu_{\epsilon,k})|^2 \right) \\ &\leq \frac{4^N}{(2\pi)^d} \iint_{\mathbb{R}_+^N \times \mathbb{R}_+^N} \int_{\mathbb{R}^d} e^{-\ell(s,t)\Psi(\xi)} |\hat{f}(\xi)|^2 d\xi \mu_{\epsilon,k}(ds) \mu_{\epsilon,k}(dt). \end{aligned} \quad (3.49)$$

Therefore, Lemma 3.7 implies that there exists a positive and finite constant a that depends only on the distance between G and the axes of \mathbb{R}_+^N , as well as $\sup_{x \in G} |x|$, such that

$$\begin{aligned} \mathbb{E}_{\lambda_d} \left(|Q_{\epsilon,k}(f)|^2 \right) &\leq \frac{4^N}{(2\pi)^d} \iint_{\mathbb{R}_+^N \times \mathbb{R}_+^N} \int_{\mathbb{R}^d} e^{-a\|s-t\|\Psi(\xi)} |\hat{f}(\xi)|^2 d\xi \mu_{\epsilon,k}(ds) \mu_{\epsilon,k}(dt) \\ &\leq \frac{4^N}{(2\pi)^d A_a^d} \iint_{\mathbb{R}_+^N \times \mathbb{R}_+^N} \int_{\mathbb{R}^d} e^{-\|s-t\|\Psi(\xi)} |\hat{f}(\xi/A_a)|^2 d\xi \mu_{\epsilon,k}(ds) \mu_{\epsilon,k}(dt). \end{aligned} \quad (3.50)$$

See **A3** for the last inequality. This, (3.44), and (3.48) together imply that

$$\begin{aligned} c \iint_{\mathbb{R}_+^N \times \mathbb{R}_+^N} \int_{\mathbb{R}^d} e^{-\|s-t\|\Psi(\xi)} |\hat{f}(\xi/A_a)|^2 d\xi \mu_{\epsilon,k}(ds) \mu_{\epsilon,k}(dt) \\ \geq \left(\iint_{\mathbb{R}_+^N \times \mathbb{R}_+^N} \int_{\mathbb{R}^d} e^{-\|s-t\|\Psi(\xi)} \hat{f}(\xi A_{1/b}) d\xi \mu_{\epsilon,k}(ds) \mu_{\epsilon,k}(dt) \right)^2 \\ \times \mathbb{P}_{\lambda_d} \{ \sigma \neq \rho, |L(0)| \leq k \}, \end{aligned} \quad (3.51)$$

where $c := 16^N (2\pi)^d / (A_a A_{1/b}^2)^d$.

The preceding is valid for every probability density f on \mathbb{R}^d whose Fourier transform is integrable and nonnegative. Now we make a particular choice of f . Namely, we replace f

by f_η , where $f_\eta(x) := \eta^{-d}g(x/\eta)$, for $\eta > 0$, and g is the density function of an isotropic stable random variable of index $\alpha \in (0, 2]$ satisfying

$$\frac{1}{\alpha} \geq \log_2 (A_a A_{1/b}). \quad (3.52)$$

Note that $\hat{g}(\xi) = \exp(-\|\xi\|^\alpha)$ for all $\xi \in \mathbb{R}^d$. Hence, as $\hat{f}_\eta(\xi) = \exp(-\eta^\alpha \|\xi\|^\alpha)$ for all $\xi \in \mathbb{R}^d$, we have

$$\begin{aligned} |\hat{f}_\eta(\xi/A_a)|^2 &= \exp\left(-\frac{2\eta^\alpha \|\xi\|^\alpha}{A_a^\alpha}\right) \\ &\leq \exp\left(-\eta^\alpha \|\xi\|^\alpha A_{1/b}^\alpha\right) \\ &= \hat{f}_\eta(\xi A_{1/b}). \end{aligned} \quad (3.53)$$

Thus, we find that

$$\begin{aligned} &\mathbb{P}_{\lambda_d} \{\sigma \neq \rho, |L(0)| \leq k\} \\ &\leq c \left[\iint_{\|s-t\| \geq \theta} \int_{\mathbb{R}^d} e^{-\|s-t\|\Psi(\xi)} \exp\left(-\frac{2\eta^\alpha \|\xi\|^\alpha}{A_a^\alpha}\right) d\xi \mu_{\epsilon,k}(ds) \mu_{\epsilon,k}(dt) \right]^{-1}. \end{aligned} \quad (3.54)$$

where $\theta > 0$ is an arbitrary parameter. Replace k by k_ϵ and let $\eta \downarrow 0$ to deduce that for all $\epsilon > 0$,

$$\mathbb{P}_{\lambda_d} \{0 \in L(G), |L(0)| \leq k_\epsilon\} \leq \tilde{c} \left[\iint_{\|s-t\| \geq \theta} \Phi(\|s-t\|) \mu_{\epsilon,k_\epsilon}(ds) \mu_{\epsilon,k_\epsilon}(dt) \right]^{-1}. \quad (3.55)$$

We may observe that $\{\mu_{\epsilon,k_\epsilon}\}_{\epsilon>0}$ is a collection of probability measures, all of which are supported on the same compact set G . Therefore, by Prohorov's theorem we can extract a subsequence of ϵ 's along which $\mu_{\epsilon,k_\epsilon}$ converges weakly to some $\mu_* \in \mathcal{P}(G)$. Because Φ is uniformly continuous on $[\theta, \infty)$, it follows that

$$\liminf_{\epsilon \rightarrow 0^+} \mathbb{P}_{\lambda_d} \{0 \in L(G), |L(0)| \leq k_\epsilon\} \leq \tilde{c} \left[\iint_{\|s-t\| \geq \theta} \Phi(\|s-t\|) \mu_*(ds) \mu_*(dt) \right]^{-1}. \quad (3.56)$$

Because we can always ensure that $\lim_{\epsilon \rightarrow 0^+} k_\epsilon = \infty$, the monotone convergence theorem insures that the left-hand side is precisely $\mathbb{P}_{\lambda_d} \{0 \in L(G)\}$. Finally, let $\theta \downarrow 0$ and appeal to the monotone convergence theorem once again to find that $\mathbb{P}_{\lambda_d} \{0 \in L(G)\} \leq \tilde{c}/I(\mu_*)$, and this is at most a constant \tilde{c} times the capacity $\text{Cap}(G)$ of G . This concludes the proof. \square

We conclude this section by proving Theorem 3.1.

Proof of Theorem 3.1. First note that

$$\begin{aligned} \mathbb{P}_{\lambda_d} \{0 \in L(G)\} &= \int_{\mathbb{R}^d} \mathbb{P} \{-x \in L(G)\} dx \\ &= \mathbb{E} [\lambda_d(L(G))]. \end{aligned} \quad (3.57)$$

This proves the identity in Theorem 3.1. Moreover, Lemma 3.11 proves that the preceding is at most $c\text{Cap}(G)$, whence follows the upper bound of the theorem.

Similarly,

$$P_{\lambda_d} \{0 \in \overline{L(G)}\} = E \left[\lambda_d \left(\overline{L(G)} \right) \right]. \quad (3.58)$$

But the set-difference between $L(G)$ and its euclidean closure is a denumerable union of sets each of which is at most $(d-1)$ -dimensional. See DALANG AND WALSH [DW92, Proposition 2.1] for the case $N = 2$; the general case is proved similarly. It follows that

$$E \left[\lambda_d \left(\overline{L(G)} \setminus L(G) \right) \right] = 0. \quad (3.59)$$

Thus,

$$\begin{aligned} P_{\lambda_d} \{0 \in L(G)\} &= E \left[\lambda_d(L(G)) \right] \\ &= E \left[\lambda_d \left(\overline{L(G)} \right) \right] \\ &= P_{\lambda_d} \{0 \in \overline{L(G)}\}. \end{aligned} \quad (3.60)$$

Apply Lemma 3.8 to finish the proof. \square

4 A perturbed random field

Let \mathfrak{X} be a d -dimensional N -parameter additive Lévy process with Lévy exponent $\nu\Psi$, where $\nu \in (0, \infty)$ is fixed. We assume also that \mathfrak{X} is independent of L . Define

$$F(t) := L(t) + \mathfrak{X}(t) \quad \text{for all } t \in \mathbb{R}_+^N. \quad (4.1)$$

The following proves that the contents of Theorem 3.1 remain unchanged if L is replaced by the perturbed random field F .

Theorem 4.1. *Let G denote a nonrandom compact subset of $(0, \infty)^N$. Then, there exists a positive and finite constant $c = c(G, \nu)$ such that*

$$c^{-1} \text{Cap}(G) \leq P_{\lambda_d} \{0 \in F(G)\} = E \left[\lambda_d(F(G)) \right] \leq c \text{Cap}(G). \quad (4.2)$$

Moreover, c depends on G only through $\inf_{x \in G} |x|$ and $\sup_{x \in G} |x|$.

Proof. Write P_z for the law of $z + F$, and $P_{\lambda_d} := \int_{\mathbb{R}^d} P_z dz$, as before. Also, E_{λ_d} denotes the corresponding expectation operator. As in Lemma 3.5, if $f \in L^1(\mathbb{R}^d)$ and $t \in \mathbb{R}_+^N$, then

$$E_{\lambda_d} [f(F(t))] = \int_{\mathbb{R}^d} f(x) dx. \quad (4.3)$$

Also, the proof of Lemma 3.6 implies that if f and g are two probability densities on \mathbb{R}^d such that $\hat{f}, \hat{g} \in L^1(\mathbb{R}^d)$, then for all $s, t \in \mathbb{R}_+^N$,

$$E_{\lambda_d} [f(F(t))g(F(s))] = \frac{1}{(2\pi)^d} \int_{\mathbb{R}^d} e^{-(\ell(s, t) + c \sum_{j=1}^N |t_j - s_j|) \Psi(\xi)} \hat{g}(\xi) \overline{\hat{f}(\xi)} d\xi. \quad (4.4)$$

Finally, as in Proposition 3.9 we have

$$\mathbb{E}_{\lambda_d} [f(F(t)) \mid \mathcal{F}(s)] = (T_{s,t}^F f)(F(s)), \quad (4.5)$$

valid for all $s, t \in \mathbb{R}_+^N$, and all measurable functions $f : \mathbb{R}^d \rightarrow \mathbb{R}_+$. Here,

$$(T_{s,t}^F f)(y) := \mathbb{E} [f(F(t) + y)] \quad \text{for all } s, t \in \mathbb{R}_+^N, y \in \mathbb{R}^d. \quad (4.6)$$

As the filtration of \mathfrak{X} is also commuting [KX03, Lemma 4.2 and its proof], the remainder of the proof of Theorem 3.1 goes through in the present case without any drastic changes. \square

5 Polar sets for the level sets

The aim of this section is to compute the Hausdorff dimension of the zero set $L^{-1}\{0\}$ of an N -parameter, d -dimensional Lévy sheet L , provided that Conditions **A1**, **A2**, and **A3** are assumed throughout.

Recall that a nonrandom set G is *polar* for a random set S if $\mathbb{P}\{S \cap G \neq \emptyset\} = 0$. We begin by describing all sets that are polar for the level sets of the random field L and its perturbation F .

Theorem 5.1. *Let G be a nonrandom compact subset of $(0, \infty)^N$. Then, the following are equivalent:*

- (i) $\mathbb{P}\{L^{-1}\{0\} \cap G \neq \emptyset\} > 0$;
- (ii) $\mathbb{P}\{F^{-1}\{0\} \cap G \neq \emptyset\} > 0$;
- (iii) $\text{Cap}(G) > 0$.

We can apply the preceding together with Corollary 2.13 of KHOSHNEVISAN AND XIAO [KX03] in order to deduce the following equivalence theorem between the Lévy sheets of this paper and their associated additive Lévy processes.

Corollary 5.2. *Let $\{\mathfrak{X}(t)\}_{t \in \mathbb{R}_+^N}$ denote the additive Lévy process associated to L . Then for all nonrandom compact sets G in \mathbb{R}_+^N there exists a positive and finite constant c such that*

$$c^{-1} \mathbb{P}\{\mathfrak{X}^{-1}\{0\} \cap G \neq \emptyset\} \leq \mathbb{P}\{L^{-1}\{0\} \cap G \neq \emptyset\} \leq c \mathbb{P}\{\mathfrak{X}^{-1}\{0\} \cap G \neq \emptyset\}. \quad (5.1)$$

Moreover, c depends only on G through $\inf_{x \in G} |x|$ and $\sup_{x \in G} |x|$.

Proof of Theorem 5.1. There exists $\alpha > 0$ such that $G \subseteq (\alpha, 1/\alpha)^N$. Therefore, if $t \in G$ then \mathbb{P}_x -a.s. for all $x \in \mathbb{R}^d$,

$$L(t) = L(a) + \dot{L}([0, t] \setminus [0, a]), \quad (5.2)$$

where $a := (\alpha, \dots, \alpha) \in (0, \infty)^N$, and the two terms on the right-hand side describe independent random fields. We can decompose the right-most term as follows:

$$\dot{L}([0, t] \setminus [0, a]) = \sum_{j=1}^N L^j(\pi^j(t - a)) + \tilde{L}(t - a), \quad (5.3)$$

where:

- (1) $L^1, \dots, L^N, \tilde{L}$ are totally independent d -dimensional random fields;
- (2) each L^j is an $(N-1)$ -parameter Lévy sheet with Lévy exponent Ψ ;
- (3) \tilde{L} is an N -parameter Lévy sheet with Lévy exponent Ψ ; and
- (4) π^j maps $t \in \mathbb{R}_+^N$ to $\pi^j t \in \mathbb{R}_+^{N-1}$, which is the same vector as t , but with t_j removed.

This identity is valid \mathbb{P}_x -a.s. for all $x \in \mathbb{R}^d$. But because we can choose path-regular versions of L^1, \dots, L^N and \tilde{L} , it follows that \mathbb{P}_x -a.s. for all $x \in \mathbb{R}^d$,

$$L(t) = L(a) + \sum_{j=1}^N L^j(\pi^j(t-a)) + \tilde{L}(t-a) \quad \text{for all } t \in G. \quad (5.4)$$

Moreover, the $N+2$ processes on the right-hand side (viewed as random fields indexed by t) are totally independent under \mathbb{P}_x for all $x \in \mathbb{R}^d$. Note that

$$\begin{aligned} \mathbb{P}\{L^{-1}\{0\} \cap G \neq \emptyset\} &= \mathbb{P}\{\exists t \in G : L(t) = 0\} \\ &= \int_{\mathbb{R}^d} \mathbb{P}\left\{\exists t \in G : \sum_{j=1}^N L^j(\pi^j(t-a)) + \tilde{L}(t-a) = -x\right\} \mathbb{P}\{L(a) \in dx\} \\ &= \int_{\mathbb{R}^d} \mathbb{P}\left\{\exists t \in G : \sum_{j=1}^N L^j(\pi^j(t-a)) + L(t-a) = -x\right\} \mathbb{P}\{L(a) \in dx\}. \end{aligned} \quad (5.5)$$

Thanks to (1.4) and the inversion theorem,

$$\frac{\mathbb{P}\{L(a) \in dx\}}{dx} = \frac{1}{(2\pi)^d} \int_{\mathbb{R}^d} \exp(-ix \cdot \xi - \alpha^N \Psi(\xi)) d\xi, \quad (5.6)$$

and this bounded above, uniformly for all $x \in \mathbb{R}^d$, by the positive and finite constant $c := c(\alpha) := (2\pi)^{-d} \int_{\mathbb{R}^d} \exp(-\alpha^N \Psi(\xi)) d\xi$. Consequently,

$$\begin{aligned} \mathbb{P}\{L^{-1}\{0\} \cap G \neq \emptyset\} &\leq c \int_{\mathbb{R}^d} \mathbb{P}\left\{\exists t \in G : \sum_{j=1}^N L^j(\pi^j(t-a)) + L(t-a) = -x\right\} dx \\ &= c \mathbb{E}[\lambda_d(W(G \ominus \{a\}))], \end{aligned} \quad (5.7)$$

where $W := \sum_{j=1}^N (L^j \circ \pi^j) + L$, and $G \ominus \{a\} := \{t-a; t \in G\}$. Because $G \ominus \{a\}$ is a nonrandom compact subset of $(0, \infty)^N$, one can prove—in exactly the same manner that we proved Theorem 4.1—that $\mathbb{E}[\lambda_d(W(G \ominus \{a\}))] \leq \text{const} \cdot \text{Cap}(G \ominus \{a\})$, where the constant depends only on G and α . We omit the details. Because $\mathbf{I}(\mu)$ is convolution-based, $G \mapsto \text{Cap}(G)$ is translation invariant, whence we have $\text{Cap}(G \ominus \{a\}) = \text{Cap}(G)$. This proves that (1) implies (3).

Next we prove that (3) implies (1). As above let $\alpha > 0$ such that $G \subset (\alpha, 1/\alpha)^N$. For any $\epsilon > 0$ and $\mu \in \mathcal{P}(\mathbb{R}_+^N)$, we define a radom measure on \mathbb{R}_+^N by

$$\mathcal{J}_\epsilon(B) = \frac{1}{(2\epsilon)^d} \int_B \mathbf{1}_{B(0,\epsilon)}(L(t)) \mu(dt) \quad \text{for all } B \subset \mathbb{R}_+^N \text{ Borel set.} \quad (5.8)$$

We shall need the following two lemmas.

Lemma 5.3. *For every $\mu \in \mathcal{P}(G)$,*

$$\liminf_{\epsilon \rightarrow 0^+} \mathbb{E}[\mathcal{J}_\epsilon(G)] > 0. \quad (5.9)$$

Proof. By Fatou's lemma,

$$\begin{aligned} \liminf_{\epsilon \rightarrow 0^+} \mathbb{E}[\mathcal{J}_\epsilon(G)] &= \liminf_{\epsilon \rightarrow 0^+} \int_G \frac{\mathbb{P}\{|L(t)| \leq \epsilon\}}{(2\epsilon)^d} \mu(dt) \\ &\geq \int_G p(t) \mu(dt), \end{aligned} \quad (5.10)$$

where $p(t)$ denotes the density function of $L(t)$ at zero. By the inversion theorem, $p(t) = \Phi(\prod_{j=1}^N t_j)$. Because μ is supported in $[\alpha, 1/\alpha]^N$,

$$\inf_{t \in G} p(t) \geq \frac{1}{(2\pi)^d} \int_{\mathbb{R}^d} \exp(-\Psi(\xi)\alpha^{-N}) d\xi := c_1 > 0, \quad (5.11)$$

where $c_1 = c_1(\alpha)$. This implies the desired result. \square

Lemma 5.4. *Let $K : \mathbb{R}_+^N \times \mathbb{R}_+^N \mapsto \mathbb{R}_+$ be a measurable function. Then there exists a finite positive constant c , depending only on G , such that for every $\mu \in \mathcal{P}(G)$ and all $\epsilon > 0$,*

$$\mathbb{E} \left[\iint_{G \times G} K(s, t) \mathcal{J}_\epsilon(ds) \mathcal{J}_\epsilon(dt) \right] \leq c \iint_{G \times G} K(s, t) \Phi(\|s - t\|) \mu(dt) \mu(ds). \quad (5.12)$$

In particular,

$$\mathbb{E} \left(|\mathcal{J}_\epsilon(G)|^2 \right) \leq c \mathbb{I}(\mu). \quad (5.13)$$

Proof. By Fubini's theorem,

$$\begin{aligned} \mathbb{E} \left[\iint_{G \times G} K(s, t) \mathcal{J}_\epsilon(ds) \mathcal{J}_\epsilon(dt) \right] \\ = \frac{1}{(2\epsilon)^{2d}} \iint_{G \times G} K(s, t) \mathbb{P}\{|L(s)| \leq \epsilon, |L(t)| \leq \epsilon\} \mu(dt) \mu(ds). \end{aligned} \quad (5.14)$$

We define

$$\begin{aligned} N_1(s, t) &:= L(s) - L(s \wedge t), \\ N_2(s, t) &:= L(t) - L(s \wedge t). \end{aligned} \quad (5.15)$$

Clearly,

$$\begin{aligned}
\mathbb{P}\{|L(s)| \leq \epsilon, |L(t)| \leq \epsilon\} &= \mathbb{P}\{|L(s \wedge t) + N_1(s, t)| \leq \epsilon, |L(s \wedge t) + N_2(s, t)| \leq \epsilon\} \\
&\leq \mathbb{P}\{|L(s \wedge t) + N_1(s, t)| \leq \epsilon, |N_2(s, t) - N_1(s, t)| \leq 2\epsilon\} \quad (5.16) \\
&= \mathbb{P}\{|L(s \wedge t) + N_1(s, t)| \leq \epsilon, |L(t) - L(s)| \leq 2\epsilon\}.
\end{aligned}$$

Note that $L(s \wedge t)$, $N_1(s, t)$, and $N_2(s, t)$ are mutually independent; this follows immediately from considering the representation of L via \dot{L} . Thus,

$$\mathbb{P}\{|L(s)| \leq \epsilon, |L(t)| \leq \epsilon\} \leq \sup_{x \in \mathbb{R}^d} \mathbb{P}_x\{|L(s \wedge t)| \leq \epsilon\} \times \mathbb{P}\{|L(t) - L(s)| \leq 2\epsilon\}. \quad (5.17)$$

Because of symmetry, $L(s \wedge t)$ is a 16^d -weakly unimodal random vector [KX03, Corollary 2.2]. That is,

$$\sup_{x \in \mathbb{R}^d} \mathbb{P}_x\{|L(s \wedge t)| \leq \epsilon\} \leq 16^d \mathbb{P}\{|L(s \wedge t)| \leq \epsilon\}. \quad (5.18)$$

But if $s, t \in G$, then $s \wedge t$ is also in $[\alpha, 1/\alpha]^N$. Hence, by (5.6) the density of $L(s \wedge t)$ is bounded, uniformly in s and t in G . Consequently, we can find a positive and finite constant $c_2 = c_2(\alpha)$ such that

$$\sup_{x \in \mathbb{R}^d} \mathbb{P}_x\{|L(s \wedge t)| \leq \epsilon\} \leq c_2(2\epsilon)^d \quad \text{for all } \epsilon > 0, s, t \in G. \quad (5.19)$$

Consequently,

$$\mathbb{E} \left[\iint_{G \times G} K(s, t) \mathcal{J}_\epsilon(ds) \mathcal{J}_\epsilon(dt) \right] \leq c_3 \iint_{G \times G} K(s, t) p(s, t) \mu(dt) \mu(ds), \quad (5.20)$$

where $p(s, t)$ is the density of $L(t) - L(s)$ at zero. By the inversion theorem,

$$\begin{aligned}
p(s, t) &= \frac{1}{(2\pi)^d} \int_{\mathbb{R}^d} e^{-\ell(s, t)\Psi(\xi)} d\xi \\
&\leq \frac{1}{(2\pi)^d} \int_{\mathbb{R}^d} e^{-a\|s-t\|\Psi(\xi)} d\xi,
\end{aligned} \quad (5.21)$$

where $a \in (0, \infty)$ depends only on G ; see Lemma 3.7. Thanks to **A3** we can write

$$\begin{aligned}
p(s, t) &\leq \frac{1}{(2\pi)^d} \int_{\mathbb{R}^d} e^{-\|s-t\|\Psi(A_a\xi)} d\xi \\
&= \frac{1}{A_a^d} \Phi(\|t - s\|),
\end{aligned} \quad (5.22)$$

and this concludes the proof. \square

Let us now continue the proof that (3) implies (1). Note that (3) implies the existence of $\mu \in \mathcal{P}(G)$ such that $I(\mu) < \infty$. Then there exists a continuous function $\rho : \mathbb{R}^N \mapsto [1, \infty)$ such that $\lim_{s \mapsto s_0} \rho(s) = \infty$ for every $s_0 \in \mathbb{R}^N$ with at least one coordinate equals 0 and such that

$$\iint_{G \times G} \rho(s - t) \Phi(\|s - t\|) \mu(dt) \mu(ds) < \infty. \quad (5.23)$$

See KHOSHNEVISAN AND XIAO [KX02, p. 73] for a construction of ρ .

Consider now the sequence of random measures $\{\mathcal{J}_\epsilon\}_{\epsilon>0}$. It follows from Lemmas 5.3, 5.4, and a second-moment argument, that we can extract a subsequence $\{\mathcal{J}_{\epsilon_n}\}_{n\in\mathbb{N}}$ —converging weakly to a random measure \mathcal{J} —such that

$$\mathbb{P}\{\mathcal{J}(G) > 0\} \geq \frac{(\inf_{0<\epsilon<1} \mathbb{E}[\mathcal{J}_\epsilon(G)])^2}{\sup_{\epsilon>0} \mathbb{E}(|\mathcal{J}_\epsilon(G)|^2)} > 0. \quad (5.24)$$

See Khoshnevisan and Xiao [KX02, proof of Lemma 3.6], and KHOSHNEVISAN, XIAO, AND SHIEH [KSX06, p. 26]). Moreover,

$$\iint_{G\times G} \rho(s-t) \mathcal{J}(dt) \mathcal{J}(ds) \leq c \iint_{G\times G} \rho(s-t) \Phi(\|s-t\|) \mu(dt) \mu(ds). \quad (5.25)$$

This and (5.23) together imply that a.s.,

$$\mathcal{J}\{s \in G : s_j = a \text{ for some } j\} = 0 \quad \text{for all } a \in \mathbb{R}_+. \quad (5.26)$$

In order to deduce (1), it suffices to prove that the random measure \mathcal{J} is supported on $L^{-1}\{0\} \cup G$. For this purpose, it suffices to prove that for every $\delta > 0$, $\mathcal{J}(D(\delta)) = 0$ a.s., where $D(\delta) := \{s \in G : |L(s)| > \delta\}$. Because of (5.26), the proof of this fact follows exactly the arguments of KHOSHNEVISAN AND XIAO [KX02, p. 76].

The equivalence of (2) and (3) is proved in exactly the same way, and is therefore omitted. \square

Next we derive our first dimension result.

Theorem 5.5. *Let G be a nonrandom compact subset of \mathbb{R}_+^N . Then, almost surely on $\{L^{-1}\{0\} \cap G \neq \emptyset\}$,*

$$\dim_{\text{H}}(L^{-1}\{0\} \cap G) = \sup \left\{ q \in (0, N) : I^{(q)}(\mu) < \infty \text{ for some } \mu \in \mathcal{P}(G) \right\}, \quad (5.27)$$

where

$$I^{(q)}(\mu) := \iint_{\{s,t \in \mathbb{R}_+^N : s \neq t\}} \frac{\Phi(\|t-s\|)}{\|t-s\|^q} \mu(ds) \mu(dt). \quad (5.28)$$

Corollary 5.6. *Almost surely on $\{L^{-1}\{0\} \neq \emptyset\}$, $\dim_{\text{H}} L^{-1}\{0\} = N - \overline{\text{ind}} \Phi$.*

Example 5.7. If L is an isotropic stable Lévy sheet of index $\alpha \in (0, 2]$, then we can combine Corollary 5.6 with Example 3.5 of Khoshnevisan, Xiao, and Shieh [KSX06] to find that a.s. on $\{L^{-1}\{0\} \cap G \neq \emptyset\}$,

$$\dim_{\text{H}}(L^{-1}\{0\} \cap G) = \dim_{\text{H}} G - (d/\alpha). \quad (5.29)$$

Also, $\dim_{\text{H}} L^{-1}\{0\} = N - (d/\alpha)$ a.s. on $\{L^{-1}\{0\} \neq \emptyset\}$. This last fact was first proved in [E81]. \square

It is not easy to produce a direct proof of Theorem 5.5. Therefore, instead we use a “comparison principle” to deduce the theorem from an analogous [known] result on the associated additive Lévy process.

Proof of Theorem 5.5. Let \mathfrak{X} denote the additive Lévy process associated to L , and choose and fix an integer $M \geq 1$ and a real number $\alpha \in (0, 2]$. Consider M independent isotropic stable Lévy processes of index α , S^1, \dots, S^M , all taking values in \mathbb{R}^N . We assume that $\{S^1, \dots, S^M, L, \mathfrak{X}\}$ are independent. Consider the M -parameter, N -valued additive stable Lévy process,

$$\mathfrak{S}(u) := S^1(u_1) + \dots + S^M(u_M), \quad \text{for all } u \in \mathbb{R}_+^M. \quad (5.30)$$

Suppose $N > \alpha M$. Then, by Propositions 4.7 and 4.8 of KHOSHNEVISAN, SHIEH, AND XIAO [KSX06] we deduce that

$$\mathbb{P} \{ \mathfrak{X}^{-1}\{0\} \cap G \cap \mathfrak{S}(\mathbb{R}_+^M) = \emptyset \} > 0 \iff \inf_{\mu \in \mathcal{P}(G)} \mathbf{I}^{(N-\alpha M)}(\mu) < \infty. \quad (5.31)$$

A standard approximation argument can be used in conjunction with Theorem 5.1 to deduce that for all nonrandom analytic sets $F \subset \mathbb{R}_+^N$,

$$\mathbb{P} \{ \mathfrak{X}^{-1}\{0\} \cap F = \emptyset \} > 0 \iff \mathbb{P} \{ L^{-1}\{0\} \cap F = \emptyset \} > 0. \quad (5.32)$$

Thanks to the independence properties of \mathfrak{S} we can apply the preceding with $F := G \cap \mathfrak{S}(\mathbb{R}_+^M)$ [first condition on \mathfrak{S} and then take expectations]. Consequently, (5.31) implies that

$$\mathbb{P} \{ L^{-1}\{0\} \cap G \cap \mathfrak{S}(\mathbb{R}_+^M) = \emptyset \} > 0 \iff \inf_{\mu \in \mathcal{P}(G)} \mathbf{I}^{(N-\alpha M)}(\mu) < \infty. \quad (5.33)$$

Now the rest proof of the proof of Theorem 5.5 follows exactly as the proof of Theorem 3.2 of KHOSHNEVISAN AND XIAO [KX03] and is therefore omitted. \square

We use a similar comparison principle to deduce Corollary 5.6.

Proof of Corollary 5.6. Let \mathfrak{X} denote the additive Lévy process associated to L , and assume that L and \mathfrak{X} are independent. Thanks to Theorem 5.5, and Theorem 3.2 of KHOSHNEVISAN AND XIAO [KSX06], the following holds with probability one.

$$\dim_{\mathbb{H}} (L^{-1}\{0\} \cap G) \mathbf{1}_{\{L^{-1}\{0\} \cap G \neq \emptyset\}} = \dim_{\mathbb{H}} (\mathfrak{X}^{-1}\{0\} \cap G) \mathbf{1}_{\{\mathfrak{X}^{-1}\{0\} \cap G \neq \emptyset\}}. \quad (5.34)$$

Now consider a nonrandom *upright cube* $G \subset (0, \infty)^N$; that is a closed set G of the form $G = \prod_{i=1}^N [a_i, b_i]$. Then, the proof of Theorem 1.1 of [KSX06] states that $\dim_{\mathbb{H}} (\mathfrak{X}^{-1}\{0\} \cap G) = N - \overline{\text{ind}} \Phi$ almost surely on $\{\mathfrak{X}^{-1}\{0\} \cap G \neq \emptyset\}$.

Hence, $\dim_{\mathbb{H}} (L^{-1}\{0\} \cap G) = N - \overline{\text{ind}} \Phi$ almost surely on $\{L^{-1}\{0\} \cap G \neq \emptyset\}$. Because we need to consider only upright cubes that have rational coordinates, a limiting argument finishes the proof. \square

6 Proof of Theorem 1.2

We now return to the problem in the Introduction, and study the SPDE (1.1).

Let $x_0 \in \mathbb{R}$ be fixed, and let $\mathcal{S}(x_0)$ be the collection of all $(t, x) \in \mathbb{R}_+ \times \mathbb{R}$ such that $t \geq |x - x_0|$. Elementary geometric considerations lead us to the following equivalent formulation:

$$\mathcal{S}(x_0) = \{(t, x) \in \mathbb{R}_+ \times \mathbb{R} : (x_0, 0) \in \mathcal{C}(t, x)\}. \quad (6.1)$$

We will need the following well-known result. See the Introduction in DALANG AND WALSH [DW92] for the case $x_0 = 0$, for example.

Lemma 6.1. *Let $x_0 \in \mathbb{R}$ be fixed. Then, we can write*

$$u(t, x) = \frac{1}{2} \tilde{F}(t - x, t + x) \quad \text{for all } (t, x) \in \mathcal{S}(x_0), \quad (6.2)$$

where \tilde{F} is a copy of the perturbed random field F of §4 with $N = 2$ and $\nu = \frac{1}{2}$.

Our proof of Theorem 1.2 requires two more simple lemmas.

Lemma 6.2. *Let $x_0 \in \mathbb{R}$ be fixed and let G be a nonrandom compact subset of $(0, \infty)^2 \cap \mathcal{S}(x_0)$. Then,*

$$\mathbb{P} \{u^{-1}\{0\} \cap G \neq \emptyset\} > 0 \iff \text{Cap}(G) > 0. \quad (6.3)$$

Proof. Define the rotation map $\rho : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ by $\rho(t, x) := (t - x, t + x)$. Its inverse is described by $\rho^{-1}(u, v) = \frac{1}{2}(v + u, v - u)$. According to Lemma 6.1,

$$u = \frac{1}{2}(\tilde{F} \circ \rho) \quad \text{on } \mathcal{S}(x_0). \quad (6.4)$$

Now the process $\frac{1}{2}F$ is the perturbed version of $\frac{1}{2}L$, and the latter is a Lévy sheet with Lévy exponent $\xi \mapsto \Psi(\xi/2)$. Thanks to Theorem 5.1, the restriction of $F^{-1}\{0\}$ to $\mathcal{S}(x_0)$ has the same polar sets as the restriction of $L^{-1}\{0\}$ on $\mathcal{S}(x_0)$. Consequently, if G is any nonrandom compact subset of $\mathcal{S}(x_0) \cap (0, \infty)^2$, then

$$\mathbb{P} \{u^{-1}\{0\} \cap G \neq \emptyset\} > 0 \iff \text{Cap}(\rho^{-1}G) > 0. \quad (6.5)$$

Note that if $\mu \in \mathcal{P}(G)$ then $\mu \circ \rho \in \mathcal{P}(\rho^{-1}G)$ and $\mathbf{I}(\mu) = \mathbf{I}(\mu \circ \rho)$. Consequently, $\text{Cap}(\rho^{-1}G) = \text{Cap}(G)$ and the lemma follows. \square

Lemma 6.3. *Let G be a nonrandom compact subset of $(0, \infty)^2$. Then,*

$$\mathbb{P} \{u^{-1}\{0\} \cap G \neq \emptyset\} > 0 \iff \text{Cap}(G) > 0. \quad (6.6)$$

Proof. If $P\{u^{-1}\{0\} \cap G \neq \emptyset\}$ is positive then there must exist $x_0 \in \mathbb{R}$ such that $P\{u^{-1}\{0\} \cap G \cap \mathcal{S}(x_0) \neq \emptyset\}$ is positive. Then Lemma 6.2 implies that $\text{Cap}(G \cap \mathcal{S}(x_0)) > 0$. Because $\text{Cap}(G) \geq \text{Cap}(G \cap \mathcal{S}(x_0))$, this proves the “ \Rightarrow ” portion of the lemma.

The converse is proved similarly. Indeed, it can be shown that Cap is a Choquet capacity, whence $\text{Cap}(G) = \sup_{x_0 \in \mathbb{R}} \text{Cap}(G \cap \mathcal{S}(x_0))$, thanks to Choquet’s capacitability theorem. \square

Finally we prove Theorem 1.2 via a comparison argument.

Proof of Theorem 1.2. Let \mathfrak{X} be the 2-parameter d -dimensional additive Lévy process that is associated to L . Lemma 6.3 used in conjunction with Theorem 5.1 and Corollary 5.2 implies that the zero sets of u and \mathfrak{X} have the same polar sets. Consequently, for all integers $n \geq 1$,

$$\begin{aligned} P\{u^{-1}\{0\} \cap [-n, n] \times [0, n] \neq \emptyset\} &> 0 \\ \iff P\{\mathfrak{X}^{-1}\{0\} \cap [-n, n] \times [0, n] \neq \emptyset\} &> 0. \end{aligned} \quad (6.7)$$

Let n tend to infinity to find that

$$P\{u^{-1}\{0\} \neq \emptyset\} > 0 \iff P\{\mathfrak{X}^{-1}\{0\} \neq \emptyset\} > 0. \quad (6.8)$$

According to Proposition 3.1 of [KX02] this last condition holds if and only if

$$\int_{[0,1]^2} \Phi(\|z\|) dz < \infty. \quad (6.9)$$

Integration in polar coordinates proves “(2) \Leftrightarrow (3)” in Theorem 1.2. Since “(1) \Rightarrow (2)” holds tautologically, in order to prove that (1) and (3) are equivalent, it suffices to prove that if $u^{-1}\{0\}$ is nonempty with positive probability, then it is nonempty a.s.

With this in mind, let us assume that $P\{u^{-1}\{0\} \neq \emptyset\} > 0$. Then there exists an integer $n \geq 1$ large enough such that

$$P\{u^{-1}\{0\} \cap \mathcal{C}(n, 0) \neq \emptyset\} > 0. \quad (6.10)$$

Thanks to (1.3), $u(t, x) = \frac{1}{2} \dot{L}(\mathcal{C}(t, x))$. We have observed in Proposition 2.5 that \dot{L} has the following two properties: (i) For all $a \in \mathbb{R}^2$ and $A \subset \mathbb{R}^2$ Borel-measurable, $\dot{L}(a + A)$ has the same law as $\dot{L}(A)$; and (ii) $\dot{L}(A)$ and $\dot{L}(B)$ are independent whenever A and B are disjoint measurable subsets of \mathbb{R}^2 . It follows from this that

$$\{u^{-1}\{0\} \cap \mathcal{C}(n, 2kn)\}_{k=0}^{\infty} \quad \text{are i.i.d. random sets.} \quad (6.11)$$

Used in conjunction with (6.10) and Borel–Cantelli’s lemma for independent events, this implies that

$$P\left\{u^{-1}\{0\} \cap \bigcup_{k=0}^{\infty} \mathcal{C}(n, 2kn) \neq \emptyset\right\} = 1. \quad (6.12)$$

Because $\bigcup_{k=0}^{\infty} \mathcal{C}(n, 2kn) \subset \mathbb{R}_+ \times \mathbb{R}$ this proves that (2) \Rightarrow (1).

In order to complete the proof of Theorem 1.2 we assume that $\int_0^1 \lambda \Phi(\lambda) d\lambda < \infty$, and proceed to compute the Hausdorff dimension of $u^{-1}\{0\}$. A comparison argument, similar to the one employed at the beginning of this proof, shows that if G is a nonrandom compact subset of $(0, \infty)^2$, then a.s.,

$$\dim_{\text{H}}(u^{-1}\{0\} \cap G) \mathbf{1}_{\{u^{-1}\{0\} \cap G \neq \emptyset\}} = \dim_{\text{H}}(\mathfrak{x}^{-1}\{0\} \cap G) \mathbf{1}_{\{\mathfrak{x}^{-1}\{0\} \cap G \neq \emptyset\}}. \quad (6.13)$$

Let $G = [0, n] \times [-n, n]$ and then let $n \rightarrow \infty$ to find that a.s.,

$$\dim_{\text{H}}(u^{-1}\{0\}) \mathbf{1}_{\{u^{-1}\{0\} \neq \emptyset\}} = \dim_{\text{H}}(\mathfrak{x}^{-1}\{0\}) \mathbf{1}_{\{\mathfrak{x}^{-1}\{0\} \neq \emptyset\}}. \quad (6.14)$$

Because we have assumed that $\int_0^1 \lambda \Phi(\lambda) d\lambda < \infty$, the already-proven equivalence of (1)–(3) shows that $\mathbf{1}_{\{u^{-1}\{0\} \neq \emptyset\}} = 1$ a.s. Therefore, Theorem 1.1 of KHOSHNEVISAN, SHIEH, AND XIAO [KSX06, Eq. (1.11)] implies the stated formula for the Hausdorff dimension of $u^{-1}\{0\}$. \square

References

- [BF75] BERG, C., FORST G. (1975), *Potential Theory on Locally Compact Abelian Groups*, Springer-Verag, New York.
- [B96] BERTOIN, J. (1996), *Lévy Processes*, Cambridge University Press, Cambridge.
- [C70] CABAÑA, E. M. (1970), The vibrating string forced by white noise, *Z. Wahrsch. und Verw. Gebiete* **15**, 111-130.
- [C72] CABAÑA, E. M. (1972), On barrier problems for the vibrating string, *Z. Wahrsch. und Verw. Gebiete* **22**, 13-24.
- [CW75] CAIROLI, R. AND WALSH, J. B. (1975), Stochastic integrals in the plane, *Acta Math.* **134**, 111-183.
- [CN88a] CARMONA, R., NUALART, D. (1988), Random nonlinear wave equations: propagation of singularities, *Ann. Probab.* **2**, 730–751.
- [CN88b] CARMONA, R., NUALART, D. (1988), Random nonlinear wave equations: smoothness of the solutions, *Probab. Theory Related Fields* **4**, 469–508.
- [D99] DALANG, R. C. (1999), Extending the martingale measure stochastic integral with applications to spatially homogeneous s.p.d.e.’s, *Electron. J. Probab.* **4**, .
- [D01] DALANG, R. C. (2001), Corrections to: “Extending the martingale measure stochastic integral with applications to spatially homogeneous s.p.d.e.’s” [Electron J. Probab. **4** (1999),], *Electron. J. Probab.* **6**, .
- [DN04] DALANG, R. C. AND NUALART, E. (2004), Potential theory for hyperbolic SPDEs, *Ann. Probab.* **32**, 2099–2148.

- [DW92] DALANG, R. C. AND WALSH, J. B. (1992), The sharp Markov property of Lévy sheets, *Ann. Probab.* **20**, 591–626.
- [DS58] DUNFORD, N. AND SCHWARTZ, J. T. (1958), *Linear Operators, Part I: General Theory*, Wiley, New York.
- [E81] EHM, W. (1981), Sample function properties of multiparameter stable processes, *Z. Wahrsch. Verw. Gebiete* **56**, 195–228.
- [G95] GAVEAU, B. (1995), The Cauchy problem for the stochastic wave equation, *Bull. Sci. Math.* **119**, 381–407.
- [G89] GAVEAU, B. (1989), Le problème de Cauchy pour une équation d’ondes stochastique, *C. R. Acad. Sci. Paris Sér. I Math.* **308**, 21–24.
- [H70] HENDRICKS, W. J. (1970), Lower envelopes near zero and infinity for processes with stable components, *Z. Wahrsch. und Verw. Gebiete* **16**, 261–278.
- [H72] HENDRICKS, W. J. (1972), Hausdorff dimension in a process with stable components - an interesting counterexample, *Ann. Math. Statist.* **43**, 690–694.
- [H73] HENDRICKS, W. J. (1973), A dimension theorem for sample functions of processes with stable components, *Ann. Probability* **1**, 849–853.
- [H74] HENDRICKS, W. J. (1974), Multiple points for a process in R^2 with stable components, *Z. Wahrsch. und Verw. Gebiete* **28**, 113–128.
- [K02] KHOSHNEVISAN, D. (2002), *Multiparameter Processes*, Springer-Verlag, New-York.
- [KX02] KHOSHNEVISAN, D., XIAO, Y. (2002), Level sets of additive Lévy processes, *Ann. Probab.* **30**, 62–100.
- [KX03] KHOSHNEVISAN, D., XIAO, Y. (2003), Weak unimodality of finite measures, and an application to potential theory of additive Lévy processes, *Proc. Amer. Math. Soc.* **131**, 2611–2616.
- [KXZ03] KHOSHNEVISAN, D., XIAO, Y., AND ZHONG, Y. (2003), Measuring the range of an additive Lévy process, *Ann. Probab.* **31**, 1097–1141.
- [KX04] KHOSHNEVISAN, D. AND XIAO, Y. (2004), Additive Lévy processes: capacity and Hausdorff dimension, *Progress in Probability* **57**, 151–170.
- [KSX06] KHOSHNEVISAN, D., SHIEH, N.-R., AND XIAO, Y. (2006), Hausdorff dimension of the contours of symmetric additive Lévy processes, *Probab. Theory Related Fields* (to appear).
- [M58] MINLOS, R. A. (1958), Continuation of a generalized random process to a completely additive measure, *Dokl. Akad. Nauk SSSR (N.S.)* **119**, 439–442.

- [M59] MINLOS, R. A. (1959), Generalized random processes and their extension in measure, *Trudy Moskov. Mat. Obsc.* **8**, 497–518.
- [M98] MUELLER, C. (1998), The heat equation with Lévy noise, *Stoch. Proc. and Appl.* **74**, 67–82.
- [MMS06] MUELLER, C., MYTNIK, L., AND STAN, A. (2006), The heat equation with time-independent multiplicative stable Lévy noise, *Stoch. Proc. and Appl.* **116**, 70–100.
- [ØPS06] OKSENDAL, B. PROSKE, F., AND SIGNAHL, M. (2006), The Cauchy problem for the wave equation with Lévy noise initial data, *Infinite dimensional analysis quantum probability and related topics* **9**, 249–270.
- [PT69] PRUITT, W. E., TAYLOR, S. J. (1969), Sample path properties of processes with stable components, *Z. Wahrsch. und Verw. Gebiete* **12**, 267–289.
- [P83] PYASETSKAYA, T. E. (1983), A semilinear stochastic hyperbolic equation with two-parameter white noise, *Theory of Random Processes*, **11**, Naukova Dumka, Kiev, 82–89.
- [S99] SATO, K. (1999), *Lévy Processes and Infinitely Divisible Distributions*, Cambridge University Press, Cambridge.
- [S37] SCHOENBERG, I. J. (1937), On certain metric spaces arising from Euclidean spaces by a change of metric and their imbedding in Hilbert space, *Ann. of Math.* (2) **38**, 787–793.
- [W86] WALSH, J. B. (1986), An Introduction to Stochastic Partial Differential Equations, *Ecole d'Été de Probabilités de Saint-Flour XIV, Lect. Notes in Math.* **1180**, Springer-Verlag, 266–437.