

1. Combinatorics

Recall the two basic principles of counting [combinatorics]:

First principle: m distinct garden forks plus n distinct fish forks equals $m + n$ distinct forks.

Second principle: m distinct knives and n distinct forks equals mn distinct ways of taking a knife and a fork.

2. Unordered Selection

Example 6.1. 6 dice are rolled. What is the probability that they all show different faces?

$$\Omega = ?$$

$$|\Omega| = 6^6.$$

If A is the event in question, then $|A| = 6 \times 5 \times 4 \times 3 \times 2 \times 1$.

Definition 6.2. If k is an integer ≥ 1 , then we define “ k factorial” as the following integer:

$$k! = k \cdot (k - 1) \cdot (k - 2) \cdots 2 \cdot 1.$$

For consistency of future formulas, we define also

$$0! = 1.$$

Example 6.3. Five rolls of a fair die. What is $P(A)$, where A is the event that all five show different faces? Note that $|A|$ is equal to 6 [which face is left out] times 6^5 . Thus,

$$P(A) = \frac{6 \cdot 5!}{6^5} = \frac{6!}{6^5}.$$

3. Ordered Selection

Example 6.4. Two-card poker.

$$P(\text{doubles}) = \frac{13 \times \binom{4 \times 3}{2}}{\binom{52 \times 51}{2}}.$$

Theorem 6.5. n objects are divided into r types. n_1 are of type 1; n_2 of type 2; \dots ; n_r are of type r . Thus, $n = n_1 + \dots + n_r$. Objects of the same type are indistinguishable. The number of permutations is

$$\binom{n}{n_1, \dots, n_r} = \frac{n!}{n_1! \cdots n_r!}.$$

Proof. Let N denote the number of permutations; we seek to find N . For every permutation in N there are $n_1! \cdots n_r!$ permutations wherein all n objects are treated differently. Therefore, $n_1! \cdots n_r! N = n!$. Solve to finish. \square

Example 6.6. n people; choose r of them to form a “team.” The number of different teams is then

$$\frac{n!}{r!(n-r)!}.$$

You have to choose r of type 1 (“put this one in the team”), and $n - r$ of type 2 (“leave this one out of the team”).

Definition 6.7. We write the preceding count statistic as “ n choose r ,” and write it as

$$\binom{n}{r} = \frac{n!}{r!(n-r)!} = \frac{n!}{(n-r)!r!} = \binom{n}{n-r}.$$

Example 6.8. Roll 4 dice; let A denote the event that all faces are different. Then,

$$|A| = \binom{6}{4} 4! = \frac{6!}{2!} = \frac{6!}{2}.$$

The 6-choose-4 is there because that is how many ways we can choose the different faces. Thus,

$$P(A) = \frac{6!}{2 \times 4^6}.$$

Example 6.9. There are

$$\binom{52}{5} = 2,598,960$$

different standard poker hands possible.