

MATH 5620 NUMERICAL ANALYSIS II
HOMEWORK 2, DUE FRI FEB 10 2012

Problem 1 B&F 5.1.4 a,b.

Notes:

- Theorem 5.3 is a sufficient condition for a function satisfying a Lipschitz condition. It is not a necessary condition as there are functions that are not differentiable but that are Lipschitz continuous.
- Theorem 5.6 gives a sufficient condition for the well-posedness of an IVP. This condition is not necessary, so Theorem 5.6 does not rule out well-posedness if the Lipschitz condition is not satisfied.

Problem 2 (K&C 8.2.2) Consider the initial value problem:

$$\begin{cases} y' = \sqrt{y} \\ y(0) = 0. \end{cases}$$

- (a) Verify that the function $y(t) = t^2/4$ is a solution to this IVP.
- (b) Apply Euler's method to this IVP **by hand** (i.e. with $h > 0$ and $t_i = ih$ find y_i for $i = 0, 1, \dots$).
- (c) Explain why the numerical solution differs from the solution $t^2/4$. (no proof necessary)

Problem 3 In the following problems you need to compare several numerical methods for initial value problems on two problems. Compare the results to the actual values (the true solutions are given in B&F 5.2.3)

B&F 5.2.1 a,c (Euler's method)

B&F 5.3.1 a,c (Taylor's method of order 2)

B&F 5.4.1 a,c (Modified Euler's method)

B&F 5.4.9 a,c (Heun's method, use def in p287 of B&F 9th ed. The 8th ed and the class notes define Heun's method differently!)

B&F 5.4.13 a,c (Runge-Kutta method of order four)

Problem 4 B&F 5.5.3 a,b (Runge-Kutta-Fehlberg method)

Note: For Problems 3 and 4, I suggest you use the stubs and sample drivers in the class website.