

Vectors & Scalars

Vectors

\mathbb{R}^2 is the set of all pairs of real numbers. In the context of drawing graphs, the objects in \mathbb{R}^2 are called points, and pairs are written left-to-right, so that $(3, 2)$ is the point in \mathbb{R}^2 whose x -coordinate equals 3 and whose y -coordinate equals 2.

In the context of linear algebra, the objects in \mathbb{R}^2 are called *vectors*, and instead of being written left-to-right, they are usually written top-to-bottom. Written in this way, the vector in \mathbb{R}^2 whose x -coordinate is 3 and whose y -coordinate is 2 is

$$\begin{pmatrix} 3 \\ 2 \end{pmatrix}$$

\mathbb{R}^3 is the set of all “triples” of real numbers. An object in \mathbb{R}^3 – also called a vector – has an x -coordinate, a y -coordinate, and a z -coordinate. When writing vectors in \mathbb{R}^3 , the x -coordinate is on top, the y -coordinate is directly below, and the z -coordinate is on the bottom. Thus

$$\begin{pmatrix} 5 \\ 0 \\ -1 \end{pmatrix}$$

is the vector in \mathbb{R}^3 where $x = 5$, $y = 0$, and $z = -1$.

Vector addition

To add two vectors in \mathbb{R}^2 – or two vectors in \mathbb{R}^3 – add each of their coordinates.

Examples.

$$\begin{pmatrix} -5 \\ 1 \end{pmatrix} + \begin{pmatrix} 4 \\ 2 \end{pmatrix} = \begin{pmatrix} -5 + 4 \\ 1 + 2 \end{pmatrix} = \begin{pmatrix} -1 \\ 3 \end{pmatrix}$$

and

$$\begin{pmatrix} 4 \\ 2 \\ 6 \end{pmatrix} + \begin{pmatrix} 3 \\ -8 \\ 0 \end{pmatrix} = \begin{pmatrix} 4 + 3 \\ 2 - 8 \\ 6 + 0 \end{pmatrix} = \begin{pmatrix} 7 \\ -6 \\ 6 \end{pmatrix}$$

Scalar multiplication

In linear algebra, real numbers are often called *scalars*. You cannot multiply two vectors, but you can multiply a scalar and a vector. To do so, multiply every coordinate in the vector by the scalar.

Examples.

$$2 \begin{pmatrix} 7 \\ -3 \end{pmatrix} = \begin{pmatrix} 2(7) \\ 2(-3) \end{pmatrix} = \begin{pmatrix} 14 \\ -6 \end{pmatrix}$$

and

$$5 \begin{pmatrix} -1 \\ 0 \\ 4 \end{pmatrix} = \begin{pmatrix} 5(-1) \\ 5(0) \\ 5(4) \end{pmatrix} = \begin{pmatrix} -5 \\ 0 \\ 20 \end{pmatrix}$$

* * * * *

Exercises

For #1-8, perform the vector arithmetic indicated.

$$1.) \begin{pmatrix} -5 \\ 1 \end{pmatrix} + \begin{pmatrix} 4 \\ 2 \end{pmatrix} \quad 2.) \begin{pmatrix} 4 \\ 2 \\ 6 \end{pmatrix} + \begin{pmatrix} 3 \\ -8 \\ 0 \end{pmatrix} \quad 3.) \begin{pmatrix} 2 \\ 8 \end{pmatrix} + \begin{pmatrix} -3 \\ 1 \end{pmatrix}$$

$$4.) \begin{pmatrix} 3 \\ 5 \\ -3 \end{pmatrix} + \begin{pmatrix} -2 \\ -1 \\ 3 \end{pmatrix} \quad 5.) 2 \begin{pmatrix} 7 \\ -3 \end{pmatrix} \quad 6.) 5 \begin{pmatrix} -1 \\ 0 \\ 4 \end{pmatrix}$$

$$7.) -1 \begin{pmatrix} 2 \\ 4 \end{pmatrix} \quad 8.) 3 \begin{pmatrix} 2 \\ 3 \\ -1 \end{pmatrix}$$

Completely factor the cubic polynomials below.

$$9.) -3x^3 + 9x^2 - 12$$

$$10.) 2x^3 - 4x^2 - 4x - 6$$