


1. Let  $\mathbf{v} = \langle 2, 1 \rangle$ .
  - (a) Rotate  $\mathbf{v}$  by  $90^\circ$  counter-clockwise.
  - (b) Rotate  $\mathbf{v}$  by  $90^\circ$  clockwise.
2. Let  $\mathbf{F} = y \sin x \mathbf{i} - z \cos x \mathbf{j} + xz^2 \mathbf{k}$ , and let  $\varphi = 2e^{xyz}$ .
  - (a) Compute  $\operatorname{div} \mathbf{F}$ .
  - (b) Compute  $\operatorname{grad} \varphi$
  - (c) Compute  $\operatorname{curl} \mathbf{F}$  and  $\nabla \times \nabla \varphi$ .
3. Evaluate the path integral

$$\oint_C (xy + 1)dx + (x^2 + y^2)dy$$

where  $C$  is the triangle with vertices  $(0, 0)$ ,  $(1, 0)$  and  $(\frac{1}{2}, 1)$ .

4. Circle "T" for True or "F" for False. In the following assume  $\mathbf{F} = \langle M, N, P \rangle$  is a vector field where  $M, N$  and  $P$  have continuous first-order partial derivatives on an open set  $R$ , and let  $\varphi$  be a potential function on the same region.
  - (a) T    F    If  $\int_C \mathbf{F} \cdot d\mathbf{r}$  is independent of path then  $\oint_C \mathbf{F} \cdot d\mathbf{r}$  is zero for every closed path  $C$ .
  - (b) T    F    Consider  $\varphi$  on the unit disk  $D$ . If  $\nabla^2 \varphi = 0$ , then the max of  $\varphi$  occurs at the origin  $(0, 0)$ .
  - (c) T    F    The shear flow in the figure below has zero curl.



5. Use Green's Theorem to compute the area of any region  $S$  in the plane where the boundary of  $S$ ,  $\partial S$  is a simple closed curve. That is show

$$A(S) = \oint_{\partial S} \mathbf{F} \cdot \mathbf{T} \, ds.$$

where  $\mathbf{F} = -\frac{1}{2}y\mathbf{i} + \frac{1}{2}x\mathbf{j}$ . (Hint: See Example 2 and Example 5 in §14.4 of your textbook)

6. Consider the force field  $\mathbf{F}(x, y) = \langle 2xy, x^2 \rangle$ . Find

$$W = \int_{\Gamma} \mathbf{F} \cdot d\mathbf{r},$$

the work done in moving an object in this field from the origin  $(0, 0)$  to  $(1, 1)$  along the parabolic arc  $\Gamma$  described by the graph of  $y = x^2$  connecting these two points. (Hint: Determine whether this force is conservative, and use a potential function to find the integral if it is. Otherwise, parameterize the arc and compute it directly.)

7. Let  $\Omega \subset \mathbb{R}^2$  be the disk of radius 2,  $\Omega = \{(x, y) \in \mathbb{R}^2 : x^2 + y^2 \leq 4\}$ . Consider the fluid velocity field  $\mathbf{v}(x, y) = \langle -y^3, x^3 \rangle$  on  $\Omega$ .

- (a) Find the curl of the fluid velocity,  $\nabla \times \mathbf{v}$ .  
 (b) Use Green's Theorem to evaluate the circulation of  $\mathbf{v}$  around the boundary of  $\Omega$ . That is, find the line integral

$$\oint_{\partial\Omega} \mathbf{v} \cdot d\mathbf{r}.$$

around the circle of radius 2 forming the boundary  $\partial\Omega$  of the disk  $\Omega$ , traversed in the counterclockwise direction, by using your result from (a) in the area integral from Green's Theorem.

8. Suppose that an object of mass  $m$  is moving along a smooth curve  $C$  given by

$$\mathbf{r} = \mathbf{r}(t) = x(t)\mathbf{i} + y(t)\mathbf{j} + z(t)\mathbf{k}, \quad a \leq t \leq b.$$

under the influence of a conservative force  $\mathbf{F}(\mathbf{r}) = \nabla f(\mathbf{r})$ . From physics, we learn three facts about the object at time  $t$

- (a)  $\mathbf{F}(\mathbf{r}(t)) = m\mathbf{r}''(t)$  (Newton's Second Law)  
 (b)  $KE = \frac{1}{2}m\|\mathbf{r}'(t)\|^2$  (kinetic energy)  
 (c)  $PE = -f(\mathbf{r})$  (potential energy)

Use the above to prove

$$\frac{d}{dt}(KE + PE) = 0.$$

That is that energy is conserved over time. (Hint: See page 747 of your text.)

9. Use the Divergence Theorem to evaluate the flux integral

$$\iint_{\partial\Omega} \mathbf{F} \cdot \mathbf{n} \, dS$$

where  $\mathbf{F} = (x - 2xy)\mathbf{i} + y^2\mathbf{j} + 3z\mathbf{k}$  and  $\Omega$  is the solid sphere of radius 3 centered at the origin.

10. Compute the flux through the unit sphere for the vector field  $\mathbf{F}(\mathbf{r}) = \mathbf{r}$ .
11. Solve Laplace's equation explicitly in one dimension,

$$u_{xx} = \frac{d^2u}{dx^2} = 0.$$

Consider a harmonic function  $u(x)$  solving the above equation on the interval  $[a, b]$ . Demonstrate by graphing  $u(x)$  that it attains its maximum and minimum values on the boundary of  $[a, b]$ .

12. Evaluate the surface integral

$$\iint_G (x^2 + y^2) \, dS$$

where  $G$  is the part of the paraboloid  $z = 1 - x^2 - y^2$  that projects onto the region  $R = \{(x, y) : x^2 + y^2 \leq 1\}$ .

13. Let  $S$  be the solid determined by  $1 \leq x^2 + y^2 + z^2 \leq 4$ , and let  $\mathbf{F} = x\mathbf{i} + (2y + z)\mathbf{j} + (z + x^2)\mathbf{k}$ . Evaluate

$$\iint_{\partial S} \mathbf{F} \cdot \mathbf{n} \, dS$$

14. Consider the potential energy of a harmonic oscillator in three dimensions given by

$$\varphi(x, y, z) = \frac{k}{2} (x^2 + y^2 + z^2).$$

- (a) Find the force  $\mathbf{F}(x, y, z) = -\nabla\varphi$  at a given position  $\mathbf{r} = (x, y, z)$ .
- (b) What are the equipotential surfaces (level sets)?
- (c) Find the divergence and curl of  $\mathbf{F}$ , that is, find  $\nabla \cdot \mathbf{F}$  and  $\nabla \times \mathbf{F}$ .
- (d) Use the divergence theorem for  $\mathbf{F}$  over  $\Omega = \{(x, y, z) \in \mathbb{R}^3 : x^2 + y^2 + z^2 \leq a^2\}$ , the ball of radius  $a$ , and your result from (b), to find the flux

$$\iint_{\partial\Omega} \mathbf{F} \cdot \mathbf{n} \, dS$$

of  $\mathbf{F}$  through the spherical surface  $\partial\Omega = \{(x, y, z) \in \mathbb{R}^3 : x^2 + y^2 + z^2 = a^2\}$ .