


## Vector Spaces and Subspaces

- Vector Space  $V$
- Subspaces  $S$  of Vector Space  $V$ 
  - The Subspace Criterion
  - Subspaces are Working Sets
  - The Kernel Theorem
  - Not a Subspace Theorem
- Independence and Dependence in Abstract spaces
  - Independence test for two vectors  $\mathbf{v}_1, \mathbf{v}_2$ . An Illustration.
  - Geometry and Independence
  - Rank Test and Determinant Test for Fixed Vectors
  - Sampling Test and Wronskian Test for Functions.
  - Independence of Atoms

## Vector Space $V$

It is a **data set**  $V$  plus a **toolkit** of eight (8) algebraic properties. The data set consists of packages of data items, called **vectors**, denoted  $\vec{X}$ ,  $\vec{Y}$  below.

Closure	The operations $\vec{X} + \vec{Y}$ and $k\vec{X}$ are defined and result in a new vector which is also in the set $V$ .	
Addition	$\vec{X} + \vec{Y} = \vec{Y} + \vec{X}$ $\vec{X} + (\vec{Y} + \vec{Z}) = (\vec{Y} + \vec{X}) + \vec{Z}$ Vector $\vec{0}$ is defined and $\vec{0} + \vec{X} = \vec{X}$ Vector $-\vec{X}$ is defined and $\vec{X} + (-\vec{X}) = \vec{0}$	commutative associative zero negative
Scalar multiply	$k(\vec{X} + \vec{Y}) = k\vec{X} + k\vec{Y}$ $(k_1 + k_2)\vec{X} = k_1\vec{X} + k_2\vec{X}$ $k_1(k_2\vec{X}) = (k_1k_2)\vec{X}$ $1\vec{X} = \vec{X}$	distributive I distributive II distributive III identity



**Figure 1.** A *Vector Space* is a data set, operations  $\oplus$  and  $\odot$ , and the 8-property toolkit.

## Definition of Subspace

---

A **subspace**  $S$  of a vector space  $V$  is a nonvoid subset of  $V$  which under the operations  $+$  and  $\cdot$  of  $V$  forms a vector space in its own right.

## Subspace Criterion

---

Let  $S$  be a subset of  $V$  such that

1. Vector  $\mathbf{0}$  is in  $S$ .
2. If  $\vec{X}$  and  $\vec{Y}$  are in  $S$ , then  $\vec{X} + \vec{Y}$  is in  $S$ .
3. If  $\vec{X}$  is in  $S$ , then  $c\vec{X}$  is in  $S$ .

Then  $S$  is a subspace of  $V$ .

Items **2**, **3** can be summarized as *all linear combinations of vectors in  $S$  are again in  $S$* . In proofs using the criterion, items 2 and 3 may be replaced by

$$c_1\vec{X} + c_2\vec{Y} \text{ is in } S.$$

## Subspaces are Working Sets

---

We call a subspace  $S$  of a vector space  $V$  a **working set**, because the purpose of identifying a subspace is to shrink the original data set  $V$  into a smaller data set  $S$ , customized for the application under study.

**A Key Example.** Let  $V$  be ordinary space  $R^3$  and let  $S$  be the plane of action of a planar kinematics experiment. The data set for the experiment is all 3-vectors  $v$  in  $V$  collected by a data recorder. Detected and recorded is the 3D position of a particle which has been constrained to a plane. The plane of action  $S$  is computed as a homogeneous equation like  $2x+3y+1000z=0$ , the equation of a plane, from the recorded data set in  $V$ . After least squares is applied to find the optimal equation for  $S$ , then  $S$  replaces the larger data set  $V$ . The customized smaller set  $S$  is the *working set* for the kinematics problem.

## The Kernel Theorem

---

### Theorem 1 (Kernel Theorem)

Let  $V$  be one of the vector spaces  $\mathbb{R}^n$  and let  $A$  be an  $m \times n$  matrix. Define a smaller set  $S$  of data items in  $V$  by the kernel equation

$$S = \{\mathbf{x} : \mathbf{x} \text{ in } V, \quad A\mathbf{x} = \mathbf{0}\}.$$

Then  $S$  is a subspace of  $V$ .

In particular, operations of addition and scalar multiplication applied to data items in  $S$  give answers back in  $S$ , and the 8-property toolkit applies to data items in  $S$ .

**Proof:** Zero is in  $V$  because  $A\mathbf{0} = \mathbf{0}$  for any matrix  $A$ . To verify the subspace criterion, we verify that  $\mathbf{z} = c_1\mathbf{x} + c_2\mathbf{y}$  for  $\mathbf{x}$  and  $\mathbf{y}$  in  $V$  also belongs to  $V$ . The details:

$$\begin{aligned} A\mathbf{z} &= A(c_1\mathbf{x} + c_2\mathbf{y}) \\ &= A(c_1\mathbf{x}) + A(c_2\mathbf{y}) \\ &= c_1A\mathbf{x} + c_2A\mathbf{y} \\ &= c_1\mathbf{0} + c_2\mathbf{0} \\ &= \mathbf{0} \end{aligned}$$

Because  $A\mathbf{x} = A\mathbf{y} = \mathbf{0}$ , due to  $\mathbf{x}, \mathbf{y}$  in  $V$ .

Therefore,  $A\mathbf{z} = \mathbf{0}$ , and  $\mathbf{z}$  is in  $V$ .

The proof is complete.

## Not a Subspace Theorem

---

### Theorem 2 (Testing $S$ not a Subspace)

Let  $V$  be an abstract vector space and assume  $S$  is a subset of  $V$ . Then  $S$  is not a subspace of  $V$  provided one of the following holds.

- (1) The vector  $0$  is not in  $S$ .
- (2) Some  $x$  and  $-x$  are not both in  $S$ .
- (3) Vector  $x + y$  is not in  $S$  for some  $x$  and  $y$  in  $S$ .

**Proof:** The theorem is justified from the *Subspace Criterion*.

1. The criterion requires  $0$  is in  $S$ .
2. The criterion demands  $cx$  is in  $S$  for all scalars  $c$  and all vectors  $x$  in  $S$ .
3. According to the subspace criterion, the sum of two vectors in  $S$  must be in  $S$ .

## Definition of Independence and Dependence

---

A list of vectors  $\mathbf{v}_1, \dots, \mathbf{v}_k$  in a vector space  $V$  are said to be **independent** provided every linear combination of these vectors is uniquely represented. **Dependent** means **not independent**.

## Unique representation

---

An equation

$$a_1\mathbf{v}_1 + \dots + a_k\mathbf{v}_k = b_1\mathbf{v}_1 + \dots + b_k\mathbf{v}_k$$

implies matching coefficients:  $a_1 = b_1, \dots, a_k = b_k$ .

## Independence Test

---

Form the system in unknowns  $c_1, \dots, c_k$

$$c_1\mathbf{v}_1 + \dots + c_k\mathbf{v}_k = \mathbf{0}.$$

Solve for the unknowns [how to do this depends on  $V$ ]. Then the vectors are independent if and only if the unique solution is  $c_1 = c_2 = \dots = c_k = 0$ .

### **Independence test for two vectors $v_1, v_2$**

---

In an abstract vector space  $V$ , form the equation

$$c_1v_1 + c_2v_2 = 0.$$

Solve this equation for  $c_1, c_2$ .

Then  $v_1, v_2$  are independent in  $V$  if and only if the system has unique solution  $c_1 = c_2 = 0$ .



## Geometry and Independence

---

- One fixed vector is independent if and only if it is nonzero.
- Two fixed vectors are independent if and only if they form the edges of a parallelogram of positive area.
- Three fixed vectors are independent if and only if they are the edges of a parallelepiped of positive volume.

In an abstract vector space  $V$ , two vectors [two data packages] are independent if and only if one is not a scalar multiple of the other.

## Illustration

---

Vectors  $\mathbf{v}_1 = \cos x$  and  $\mathbf{v}_2 = \sin x$  are two data packages [graphs] in the vector space  $V$  of continuous functions. They are independent because one graph is not a scalar multiple of the other graph.

## An Illustration of the Independence Test

---

Two column vectors are tested for independence by forming the system of equations  $c_1\mathbf{v}_1 + c_2\mathbf{v}_2 = \mathbf{0}$ , e.g,

$$c_1 \begin{pmatrix} -1 \\ 1 \end{pmatrix} + c_2 \begin{pmatrix} 2 \\ 1 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}.$$

This is a homogeneous system  $\mathbf{A}\mathbf{c} = \mathbf{0}$  with

$$\mathbf{A} = \begin{pmatrix} -1 & 2 \\ 1 & 1 \end{pmatrix}, \quad \mathbf{c} = \begin{pmatrix} c_1 \\ c_2 \end{pmatrix}.$$

The system  $\mathbf{A}\mathbf{c} = \mathbf{0}$  can be solved for  $\mathbf{c}$  by frame sequence methods. Because  $\text{rref}(\mathbf{A}) = \mathbf{I}$ , then  $c_1 = c_2 = 0$ , which verifies independence.

If the system  $\mathbf{A}\mathbf{c} = \mathbf{0}$  is square, then  $\det(\mathbf{A}) \neq 0$  applies to test independence.

There is **no chance to use determinants** when the system is not square, e.g., consider the homogeneous system

$$c_1 \begin{pmatrix} -1 \\ 1 \\ 0 \end{pmatrix} + c_2 \begin{pmatrix} 2 \\ 1 \\ 0 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}.$$

It has vector-matrix form  $\mathbf{A}\mathbf{c} = \mathbf{0}$  with  $3 \times 2$  matrix  $\mathbf{A}$ , for which  $\det(\mathbf{A})$  is undefined.

## Rank Test

---

In the vector space  $\mathbf{R}^n$ , the independence test leads to a system of  $n$  linear homogeneous equations in  $k$  variables  $c_1, \dots, c_k$ . The test requires solving a matrix equation  $\mathbf{A}\mathbf{c} = \mathbf{0}$ . The signal for independence is **zero free variables**, or nullity zero, or equivalently, maximal rank. To justify the various statements, we use the relation  $\text{nullity}(\mathbf{A}) + \text{rank}(\mathbf{A}) = k$ , where  $k$  is the column dimension of  $\mathbf{A}$ .

### Theorem 3 (Rank-Nullity Test)

Let  $\mathbf{v}_1, \dots, \mathbf{v}_k$  be  $k$  column vectors in  $\mathbf{R}^n$  and let  $\mathbf{A}$  be the augmented matrix of these vectors. The vectors are independent if  $\text{rank}(\mathbf{A}) = k$  and dependent if  $\text{rank}(\mathbf{A}) < k$ . The conditions are equivalent to  $\text{nullity}(\mathbf{A}) = 0$  and  $\text{nullity}(\mathbf{A}) > 0$ , respectively.

## Determinant Test

---

In the unusual case when the system arising in the independence test can be expressed as  $\mathbf{A}\mathbf{c} = \mathbf{0}$  and  $\mathbf{A}$  is square, then  $\det(\mathbf{A}) = 0$  detects dependence, and  $\det(\mathbf{A}) \neq 0$  detects independence. The reasoning is based upon the adjugate formula  $\mathbf{A}^{-1} = \mathbf{adj}(\mathbf{A}) / \det(\mathbf{A})$ , valid exactly when  $\det(\mathbf{A}) \neq 0$ .

### Theorem 4 (Determinant Test)

Let  $\mathbf{v}_1, \dots, \mathbf{v}_n$  be  $n$  column vectors in  $\mathbf{R}^n$  and let  $\mathbf{A}$  be the augmented matrix of these vectors. The vectors are independent if  $\det(\mathbf{A}) \neq 0$  and dependent if  $\det(\mathbf{A}) = 0$ .

## Sampling Test

---

Let functions  $f_1, \dots, f_n$  be given and let  $x_1, \dots, x_n$  be distinct  $x$ -sample values. Define

$$A = \begin{pmatrix} f_1(x_1) & f_2(x_1) & \cdots & f_n(x_1) \\ f_1(x_2) & f_2(x_2) & \cdots & f_n(x_2) \\ \vdots & \vdots & \cdots & \vdots \\ f_1(x_n) & f_2(x_n) & \cdots & f_n(x_n) \end{pmatrix}.$$

Then  $\det(A) \neq 0$  implies  $f_1, \dots, f_n$  are independent functions.

### Proof

---

We'll do the proof for  $n = 2$ . Details are similar for general  $n$ . Assume  $c_1 f_1 + c_2 f_2 = 0$ . Then for all  $x$ ,  $c_1 f_1(x) + c_2 f_2(x) = 0$ . Choose  $x = x_1$  and  $x = x_2$  in this relation to get  $Ac = 0$ , where  $c$  has components  $c_1, c_2$ . If  $\det(A) \neq 0$ , then  $A^{-1}$  exists, and this in turn implies  $c = A^{-1}Ac = 0$ . We conclude  $f_1, f_2$  are independent.

## Wronskian Test

---

Let functions  $f_1, \dots, f_n$  be given and let  $x_0$  be a given point. Define

$$W = \begin{pmatrix} f_1(x_0) & f_2(x_0) & \cdots & f_n(x_0) \\ f_1'(x_0) & f_2'(x_0) & \cdots & f_n'(x_0) \\ \vdots & \vdots & \cdots & \vdots \\ f_1^{(n-1)}(x_0) & f_2^{(n-1)}(x_0) & \cdots & f_n^{(n-1)}(x_0) \end{pmatrix}.$$

Then  $\det(W) \neq 0$  implies  $f_1, \dots, f_n$  are independent functions. The matrix  $W$  is called the **Wronskian Matrix** of  $f_1, \dots, f_n$  and  $\det(W)$  is called the **Wronskian determinant**.

### Proof

---

We'll do the proof for  $n = 2$ . Details are similar for general  $n$ . Assume  $c_1 f_1 + c_2 f_2 = 0$ . Then for all  $x$ ,  $c_1 f_1(x) + c_2 f_2(x) = 0$  and  $c_1 f_1'(x) + c_2 f_2'(x) = 0$ . Choose  $x = x_0$  in this relation to get  $Wc = 0$ , where  $c$  has components  $c_1, c_2$ . If  $\det(W) \neq 0$ , then  $W^{-1}$  exists, and this in turn implies  $c = W^{-1}Wc = 0$ . We conclude  $f_1, f_2$  are independent.

## Atoms

---

**Definition.** A function is called an **atom** provided it has coefficient **1** and is obtained as the real or imaginary part of the expression

$$x^k e^{ax} (\cos bx + i \sin bx).$$

The powers **1**,  $x$ ,  $x^2$ , ... are atoms (select  $a = b = 0$ ). Multiples of these powers by  $\cos bx$ ,  $\sin bx$  are also atoms. Finally, multiplying all these atoms by  $e^{ax}$  expands and completes the list of atoms.

## Illustration

---

We show the powers  $1, x, x^2, x^3$  are independent atoms by applying the Wronskian Test:

$$W = \begin{pmatrix} 1 & x_0 & x_0^2 & x_0^3 \\ 0 & 1 & 2x_0 & 3x_0^2 \\ 0 & 0 & 2 & 6x_0 \\ 0 & 0 & 0 & 6 \end{pmatrix}.$$

Then  $\det(W) = 12 \neq 0$  implies the functions  $1, x, x^2, x^3$  are linearly independent.



## Subsets of Independent Sets are Independent

---

Suppose  $v_1, v_2, v_3$  make an independent set and consider the subset  $v_1, v_2$ . If

$$c_1v_1 + c_2v_2 = 0$$

then also

$$c_1v_1 + c_2v_2 + c_3v_3 = 0$$

where  $c_3 = 0$ . Independence of the larger set implies  $c_1 = c_2 = c_3 = 0$ , in particular,  $c_1 = c_2 = 0$ , and then  $v_1, v_2$  are independent.

### Theorem 5 (Subsets and Independence)

- A non-void subset of an independent set is also independent.
- Non-void subsets of dependent sets can be independent or dependent.

## Atoms and Independence

---

### **Theorem 6 (Independence of Atoms)**

Any list of distinct atoms is linearly independent.

### **Unique Representation**

---

The theorem is used to extract equations from relations involving atoms. For instance:

$$(c_1 - c_2) \cos x + (c_1 + c_3) \sin x + c_1 + c_2 = 2 \cos x + 5$$

implies

$$c_1 - c_2 = 2,$$

$$c_1 + c_3 = 0,$$

$$c_1 + c_2 = 5.$$

## Atoms and Differential Equations

---

It is known that solutions of linear constant coefficient differential equations of order  $n$  and also systems of linear differential equations with constant coefficients have a general solution which is a linear combination of atoms.

- The harmonic oscillator  $y'' + b^2y = 0$  has general solution  $y(x) = c_1 \cos bx + c_2 \sin bx$ . This is a linear combination of the two atoms  $\cos bx, \sin bx$ .
- The third order equation  $y''' + y' = 0$  has general solution  $y(x) = c_1 \cos x + c_2 \sin x + c_3$ . The solution is a linear combination of the independent atoms  $\cos x, \sin x, 1$ .
- The linear dynamical system  $x'(t) = y(t), y'(t) = -x(t)$  has general solution  $x(t) = c_1 \cos t + c_2 \sin t, y(t) = -c_1 \sin t + c_2 \cos t$ , each of which is a linear combination of the independent atoms  $\cos t, \sin t$ .