

## How to Solve Linear Differential Equations

- Definition: Base Atom, Atom
- Independence of Atoms
- Construction of the General Solution from a List of Distinct Atoms
- Euler's Theorems
  - Euler's Basic Theorem
  - Euler's Multiplicity Theorem
  - A Shortcut Method
- Examples
- Main Theorems on Atoms and Linear Differential Equations

## Atoms

---

A **base atom** is one of  $1$ ,  $e^{ax}$ ,  $\cos bx$ ,  $\sin bx$ ,  $e^{ax} \cos bx$ ,  $e^{ax} \sin bx$ , with  $b > 0$  and  $a \neq 0$ .

An **atom** equals  $x^n$  times a base atom, where  $n \geq 0$  is an integer.

## Details and Remarks

---

- Euler's formula  $e^{i\theta} = \cos \theta + i \sin \theta$  implies that an atom is constructed from the complex expression  $x^n e^{ax+ibx}$  by taking real and imaginary parts.
- The powers  $1, x, x^2, \dots, x^k$  are atoms.
- The term that makes up an atom has coefficient 1, therefore  $2e^x$  is not an atom, but the 2 can be stripped off to create the atom  $e^x$ . Zero is not an atom. Linear combinations like  $2x + 3x^2$  are not atoms, but the individual terms  $x$  and  $x^2$  are indeed atoms. Terms like  $-e^x$ ,  $e^{-x^2}$ ,  $x^{5/2} \cos x$ ,  $\ln |x|$  and  $x/(1 + x^2)$  are not atoms.

## Independence

---

Linear algebra defines a list of functions  $f_1, \dots, f_k$  to be **linearly independent** if and only if the representation of the zero function as a linear combination of the listed functions is uniquely represented, that is,

$$0 = c_1 f_1(x) + c_2 f_2(x) + \dots + c_k f_k(x) \text{ for all } x$$

implies  $c_1 = c_2 = \dots = c_k = 0$ .

## Independence and Atoms

---

### Theorem 1 (Atoms are Independent)

A list of finitely many distinct atoms is linearly independent.

### Theorem 2 (Powers are Independent)

The list of distinct atoms  $1, x, x^2, \dots, x^k$  is linearly independent. And all of its sublists are linearly independent.

## Construction of the General Solution from a List of Distinct Atoms

- **Picard's theorem** says that the homogeneous constant-coefficient linear differential equation

$$y^{(n)} + p_{n-1}y^{(n-1)} + \cdots + p_1y' + p_0y = 0$$

has solution space  $S$  of dimension  $n$ . Picard's theorem reduces the general solution problem to finding  $n$  linearly independent solutions.

- **Euler's theorem** *infra* says that the required  $n$  independent solutions can be taken to be atoms. The theorem explains how to construct a list of distinct atoms, each of which is a solution of the differential equation, from the roots of the characteristic equation [**characteristic polynomial**=left side]

$$r^n + p_{n-1}r^{n-1} + \cdots + p_1r + p_0 = 0.$$

- The **Fundamental Theorem of Algebra** states that there are exactly  $n$  roots  $r$ , real or complex, for an  $n$ th order polynomial equation. The result implies that the characteristic equation has exactly  $n$  roots, counting multiplicities.
- **General Solution.** Because the list of atoms constructed by Euler's theorem has  $n$  distinct elements, then this list of independent atoms forms a **basis** for the general solution of the differential equation, giving

$$y = c_1(\text{atom } 1) + \cdots + c_n(\text{atom } n).$$

Symbols  $c_1, \dots, c_n$  are arbitrary coefficients. In particular, each atom listed is itself a solution of the differential equation.

## Euler's Basic Theorem

---

### Theorem 3 (L. Euler)

The exponential  $y = e^{r_1 x}$  is a solution of a constant-coefficient linear homogeneous differential of the  $n$ th order if and only if  $r = r_1$  is a root of the characteristic equation.

---

- If  $r_1 = a$  is a real root, then one atom  $e^{ax}$  is constructed by Euler's Theorem.
- If  $r_1 = a + ib$  is a complex root ( $b > 0$ ), then Euler's Theorem gives a complex solution

$$e^{r_1 x} = e^{ax} \cos bx + ie^{ax} \sin bx.$$

The real and imaginary parts of this complex solutions are real solutions of the differential equation. Therefore, one complex root  $r_1 = a + ib$  produces *two atoms*

$$e^{ax} \cos bx, \quad e^{ax} \sin bx.$$

The conjugate root  $a - ib$  produces the same two atoms, hence it is ignored.

## Euler's Multiplicity Theorem

---

**Definition.** A root  $r = r_1$  of a polynomial equation  $p(r) = 0$  has **multiplicity  $k$**  provided  $(r - r_1)^k$  divides  $p(r)$  but  $(r - r_1)^{k+1}$  does not divide  $p(r)$ . The calculus equivalent is  $p^{(j)}(r_1) = 0$  for  $j = 0, \dots, k - 1$  and  $p^{(k)}(r_1) \neq 0$ .

### Theorem 4 (L. Euler)

The expression  $y = x^k e^{r_1 x}$  is a solution of a constant-coefficient linear homogeneous differential of the  $n$ th order if and only if  $(r - r_1)^{k+1}$  divides the characteristic polynomial.

## A Shortcut for using Euler's Theorems

---

Let root  $r_1$  of the characteristic equation have multiplicity  $m + 1$ .

- Find a base atom from Euler's Basic Theorem.
- Multiply the base atom by  $1, x, \dots, x^m$ .

This process constructs, from the base atom, exactly  $m + 1$  atoms. The atom count  $m + 1$  equals the multiplicity of root  $r_1$ .

- A real root  $r_1 = a$  will produce one base atom  $e^{ax}$ , to which this process is applied.
- A complex root  $r_1 = a + ib$  will produce 2 base atoms  $e^{ax} \cos bx, e^{ax} \sin bx$ . This process is applied to both.

## Atom List Examples

---

1. If root  $r = -3$  has multiplicity 4, then the atom list is

$$e^{-3x}, xe^{-3x}, x^2e^{-3x}, x^3e^{-3x}.$$

The list is constructed by multiplying the base atom  $e^{-3x}$  by powers  $1, x, x^2, x^3$ . The multiplicity 4 of the root equals the number of constructed atoms.

2. If  $r = -3 + 2i$  is a root of the characteristic equation, then the base atoms for this root (both  $-3 + 2i$  and  $-3 - 2i$  counted) are

$$e^{-3x} \cos 2x, \quad e^{-3x} \sin 2x.$$

If root  $r = -3 + 2i$  has multiplicity 3, then the two real atoms are multiplied by  $1, x, x^2$  to obtain a total of 6 atoms

$$e^{-3x} \cos 2x, \quad xe^{-3x} \cos 2x, \quad x^2e^{-3x} \cos 2x, \\ e^{-3x} \sin 2x, \quad xe^{-3x} \sin 2x, \quad x^2e^{-3x} \sin 2x.$$

The number of atoms generated for each base atom is 3, which equals the multiplicity of the root  $-3 + 2i$ .

### **Theorem 5 (Homogeneous Solution $y_h$ and Atoms)**

Linear homogeneous differential equations with constant coefficients have general solution  $y_h(x)$  equal to a linear combination of atoms.

### **Theorem 6 (Particular Solution $y_p$ and Atoms)**

A linear non-homogeneous differential equation with constant coefficients having forcing term  $f(x)$  equal to a linear combination of atoms has a particular solution  $y_p(x)$  which is a linear combination of atoms.

### **Theorem 7 (General Solution $y$ and Atoms)**

A linear non-homogeneous differential equation with constant coefficients having forcing term

$$f(x) = \text{a linear combination of atoms}$$

has general solution

$$y(x) = y_h(x) + y_p(x) = \text{a linear combination of atoms.}$$

### **Proofs**

---

The first theorem follows from Picard's theorem, Euler's theorem and independence of atoms. The second follows from the method of undetermined coefficients, *infra*. The third theorem follows from the first two.