

January 2007

Name \_\_\_\_\_ Class Time \_\_\_\_\_

Project 8. Solve problems L8.1 to L8.6. The problem headers:

\_\_\_\_\_ PROBLEM L8.1. BUILDING MODEL FOR AN EARTHQUAKE.  
 \_\_\_\_\_ PROBLEM L8.2. TABLE OF NATURAL FREQUENCIES AND PERIODS.  
 \_\_\_\_\_ PROBLEM L8.3. UNDETERMINED COEFFICIENTS STEADY-STATE PERIODIC SOLUTION.  
 \_\_\_\_\_ PROBLEM L8.4. PRACTICAL RESONANCE.  
 \_\_\_\_\_ PROBLEM L8.5. EARTHQUAKE DAMAGE.  
 \_\_\_\_\_ PROBLEM L8.6. SIX FLOORS.

```
> with(linalg):
```

## 1. BUILDING MODEL FOR AN EARTHQUAKE.

Refer to the textbook of Edwards-Penney, section 7.4, page 437. Consider a building with 7 floors.

Let the mass in slugs of each story be  $m = 1000.0$  and let the spring constant be  $k = 10000$  lbs/foot. Define the 7 by 7 mass matrix  $M$  and Hooke's matrix  $K$  for this system and convert  $Mx'' = Kx$  into the system  $x'' = Ax$  where  $A$  is defined by textbook equation (1) , page 437.

## PROBLEM L8.1

Find the eigenvalues of the matrix  $A$  to six digits, using the Maple command `eigenvals(A)`. Justify in particular that all seven eigenvalues are negative by direct computation.

# Sample Maple code for a model with 4 floors.

# Use maple help to learn about evalf and eigenvals.

```
A:=matrix([ [-20,10,0,0], [10,-20,10,0], [0,10,-20,10], [0,0,10,-10]]);
```

```
evalf(eigenvals(A));
```

```
> # Problem L8.1
```

## 2. TABLE OF NATURAL FREQUENCIES AND PERIODS.

Refer to figure 7.4.17, page 437.

## PROBLEM L8.2.

Find the natural angular frequencies  $\omega = \sqrt{-\lambda}$  for the seven story building and also the corresponding periods  $2\pi/\omega$ , accurate to six digits. Display the answers in a table . The answers appear in Figure 7.4.17, page 437.

# Sample code for a 4x3 table.

# Use maple help to learn about nops and printf.

```
ev:=[-10,-1.206147582,-35.32088886,-23.47296354]: n:=nops(ev):
```

```
Omega:=lambda->sqrt(-lambda):
```

```
format:=" %10.6f %10.6f %10.6f\n":
```

```
seq(printf(format,ev[i],Omega(ev[i]),2*evalf(Pi)/Omega(ev[i])),i=1..n );
```

```
> # Problem L8.2
```

### 3. UNDETERMINED COEFFICIENTS STEADY-STATE PERIODIC SOLUTION.

Consider the forced equation  $x' = Ax + \cos(wt)b$  where  $b$  is a constant vector. The earthquake's ground vibration is accounted for by the extra term  $\cos(wt)b$ , which has period  $T = 2\pi/w$ . The solution  $x(t)$  is the 7-vector of excursions from equilibrium of the corresponding 7 floors. Sought here is not the general solution, which certainly contains transient terms, but rather the steady-state periodic solution, which is known from the theory to have the form  $x(t) = \cos(wt)c$  for some vector  $c$  that depends only on  $A$  and  $b$ .

#### PROBLEM L8.3.

Define `b:=0.25*w*w*vector([1,1,1,1,1,1,1]):` in Maple and find the vector  $c$  in the undetermined coefficients solution  $x(t) = \cos(wt)c$ . Vector  $c$  depends on  $w$ . As outlined in the textbook, vector  $c$  can be found by solving the linear algebra problem  $-w^2c = Ac + b$ ; see page 433. Don't print  $c$ , as it is too complex; instead, print `c[1]` as an illustration.

# Sample code for defining  $b$  and  $A$ , then solving for  $c$  in the 4-floor case.

# See maple help to learn about vector and linsolve.

w:='w': u:=w\*w: b:=0.25\*u\*vector([1,1,1,1]):

Au:=matrix([ [-20+u,10,0,0], [10,-20+u,10,0], [0,10,-20+u,10], [0,0,10,-10+u]]);

c:=linsolve(Au,-b):

evalf(c[1],2);

> # PROBLEM L8.3

### 4 PRACTICAL RESONANCE.

Consider the forced equation  $x' = Ax + \cos(wt)b$  of L8.3 above. Practical resonance can occur if a component of  $x(t)$  has large amplitude compared to the vector norm of  $b$ . For example, an earthquake might cause a small 3-inch excursion on level ground, but the building's floors might have 50-inch excursions, enough to destroy the building.

#### PROBLEM L8.4.

Let  $\text{Max}(c)$  denote the maximum modulus of the components of vector  $c$ . Plot  $g(T) = \text{Max}(c(w))$  with  $w = 2\pi/T$  for periods  $T = 0$  to  $T = 6$ , ordinates  $\text{Max} = 0$  to  $\text{Max} = 10$ , the vector  $c(w)$  being the answer produced in L8.3 above. Compare your figure to the textbook Figure 7.4.18, page 438.

# Sample maple code to define the function  $\text{Max}(c)$ , 4-floor building.

# Use maple help to learn about norm, vector, subs and linsolve.

with(linalg):

w:='w': Max:= c -> norm(c,infinity); u:=w\*w:

b:=0.25\*w\*w\*vector([1,1,1,1]):

Au:=matrix([ [-20+u,10,0,0], [10,-20+u,10,0], [0,10,-20+u,10], [0,0,10,-10+u]]);

C:=ww -> subs(w=ww,linsolve(Au,-b)):

plot(Max(C(2\*Pi/r)),r=0..10,numpoints=150);

> # PROBLEM L8.4. WARNING: Save your file

> often!!!

### 5. EARTHQUAKE DAMAGE.

The maximum amplitude plot of L8.4 can be used to detect the likelihood of earthquake damage for a given ground vibration of period  $T$ . A ground vibration  $(1/4)\cos(wt)$ ,  $T = 2\pi/w$ , will be assumed, as in L8.4.

#### PROBLEM L8.5.

- Replot the amplitudes in L8.4 for periods 1.14 to 4 and amplitudes 5 to 10. There will be four spikes.
- Create four zoom-in plots, one for each spike, choosing a  $T$ -interval that shows the full spike.
- Determine from the four zoom-in plots approximate intervals for the period  $T$  such that some floor in the building will undergo excursions from equilibrium in excess of 5 feet.

```

# Example: Zoom-in on a spike for amplitudes 5 feet to 10 feet, periods 1.97 to 2.01.
with(linalg): w:= 'w': Max:= c -> norm(c,infinity); u:=w*w:
Au:=matrix([ [-20+u,10,0,0], [10,-20+u,10,0], [0,10,-20+u,10], [0,0,10,-10+u]]);
b:=0.25*w*w*vector([1,1,1,1]):
C:=ww -> subs(w=ww,linsolve(Au,-b)):
plot(Max(C(2*Pi/r)),r=1.97..2.01,5..10,numpoints=150);
> # PROBLEM L5.5. WARNING: Save your file often!!
> # (a) Plot four spikes on separate graphs
> # (b) Plot four zoom-in graphs.
> # (c) Print period ranges.

```

## 6. SIX FLOORS.

Consider a building with six floors each weighing 50 tons. Assume each floor corresponds to a restoring Hooke's force with constant  $k = 5$  tons/foot. Assume that ground vibrations from the earthquake are modeled by  $(1/4)\cos(wt)$  with period  $T = 2\pi/w$  (same as the 7-floor model above).

### PROBLEM L8.6.

Model the 6-floor problem in Maple. Plot the maximum amplitudes against the period 0 to 6 and amplitude 4 to 10. Determine from the graphic the period ranges which cause the amplitude plot to spike above 4 feet. Sanity check:  $m=3125$ , and the 6x6 matrix contains fraction  $16/5$ . There are five spikes.

```

> # PROBLEM L8.6. WARNING: Save your file
> often!!
> # Define k, m and the 6x6 matrix A.
> # Amplitude plot for T=0..6,C=4..10
> # Plot five zoom-in graphs
> # From the graphics, five T-ranges give
> amplitude # spikes above 4 feet. These are determined by left
> # mouse-clicks on the graph, so they are approximate values only.

```