

Laplace Ch10

Lerch's Theorem

If $\mathcal{L}(f) = \mathcal{L}(g)$ for $s \geq s_0$, Then $f(t) = g(t)$

Example. If $\mathcal{L}(y(t)) = \mathcal{L}(te^{-t})$, Then $y(t) = te^{-t}$
This is the basic cancellation law for solving equations.

Laplace Table

$f(t)$	$\int_0^{\infty} e^{-st} f(t) dt$
1	$\frac{1}{s}$
t^n	$\frac{n!}{s^{n+1}}$
e^{at}	$\frac{1}{s-a}$
$\cos(\omega t)$	$\frac{s}{s^2 + \omega^2}$
$\sin(\omega t)$	$\frac{\omega}{s^2 + \omega^2}$
$H(t-a)$	$\frac{e^{-as}}{s} \quad (a \geq 0)$

Laplace Rules

$\mathcal{L}(f) = \int_0^{\infty} e^{-st} f(t) dt$ (Direct transform)

$\mathcal{L}(c_1 f_1 + c_2 f_2) = c_1 \mathcal{L}(f_1) + c_2 \mathcal{L}(f_2)$ (Linearity)

$\mathcal{L}(-t)f(t) = \frac{d}{ds} [\mathcal{L}(f)]$ (s-diff)

$\mathcal{L}(tf'(t)) = s \mathcal{L}(f) - f(0)$ (t-diff)

$\mathcal{L}(e^{at} f(t)) = \mathcal{L}(f) |_{s \rightarrow s-a}$ (shift)

$\mathcal{L}(f(t-a)H(t-a)) = e^{-as} \mathcal{L}(f)$, $\mathcal{L}(g(t)H(t-a)) = e^{-as} \mathcal{L}(g(t+a))$

Other rules:

- periodic rule
- Convolution
- Integral

Evaluate $\mathcal{L}(5e^{-t})$

$$\mathcal{L}(5e^{-t}) = \mathcal{L}(5e^{-t})$$

$$= 5e \mathcal{L}(e^{-t})$$

$$= 5e \frac{1}{s-(-1)}$$

$$= \boxed{\frac{5e}{s+1}}$$

- Exponential rule
 $e^a e^b = e^{a+b}$
- Linearity of \mathcal{L}
- $\mathcal{L}(e^{at}) = \frac{1}{s-a}$

Evaluate $\mathcal{L}(e^{-t/3} + \sinh(-t/3))$

$$f(t) = e^{-t/3} + \sinh(-t/3)$$

$$= e^{-t/3} + \frac{1}{2}(e^{-t/3} - e^{t/3})$$

$$= \frac{3}{2}e^{-t/3} - \frac{1}{2}e^{t/3}$$

$$\mathcal{L}(f) = \frac{3}{2} \mathcal{L}(e^{-t/3}) - \frac{1}{2} \mathcal{L}(e^{t/3})$$

$$= \frac{3}{2} \frac{1}{s-(-1/3)} - \frac{1}{2} \frac{1}{s-1/3}$$

$$= \boxed{\frac{1.5}{s+1/3} - \frac{0.5}{s-1/3}}$$

• $\sinh(u) \equiv \frac{1}{2}(e^u - e^{-u})$

• Linearity of \mathcal{L}

• $\mathcal{L}(e^{at}) = \frac{1}{s-a}$

Maple might return the answer as a single fraction, e.g.,

$$\frac{s - 2/3}{s^2 - 1/9}$$

Generally, transform answers are best left unsimplified and unchanged from the first instance of a valid expression. Only when checking answers does the issue of other forms of the same answer become an issue.

Given $f(t) = t^2 \cos(3t)$, find $\mathcal{L}\{f(t)\}$.

$$\mathcal{L}\{f(t)\} = \mathcal{L}\{(-t)^2 \cos(3t)\}$$

$$= \frac{d}{ds} \frac{d}{ds} \mathcal{L}\{\cos 3t\}$$

$$= \left(\frac{d}{ds}\right)^2 \left(\frac{s}{s^2+9}\right)$$

$$= \boxed{\frac{8s^3}{(s^2+9)^3} - \frac{6s}{(s^2+9)^2}}$$

Given $f(t) = \sin^2(3t)$, find $\mathcal{L}\{f(t)\}$

$$\mathcal{L}\{f(t)\} = \mathcal{L}\{\sin^2(3t)\}$$

$$= \mathcal{L}\left\{\frac{1}{2} - \frac{1}{2} \cos(6t)\right\}$$

$$= \mathcal{L}\left\{\frac{1}{2}\right\} - \mathcal{L}\left\{\frac{1}{2} \cos(6t)\right\}$$

$$= \frac{1}{2} \mathcal{L}\{1\} - \frac{1}{2} \mathcal{L}\{\cos(6t)\}$$

$$= \boxed{\frac{1}{2} \frac{1}{s} - \frac{1}{2} \frac{s}{s^2+36}}$$

- Prepare to use s-diff rule
- Apply s-diff rule twice
- Table

• Found second derivative. Verified in Maple. See details at bottom of page.

• Given

• Use $\cos(2\theta) = 1 - 2\sin^2(\theta)$.

• Identity

• Identity again

• Table

Details:

$$\frac{d}{ds} \left(\frac{s}{s^2+9}\right) = \frac{1}{s^2+9} - \frac{2s^2}{(s^2+9)^2}$$

$$\frac{d^2}{ds^2} \left(\frac{s}{s^2+9}\right) = \frac{d}{ds} \left(\frac{1}{s^2+9}\right) - \frac{d}{ds} \left(\frac{2s^2}{(s^2+9)^2}\right)$$

$$= \frac{-2s}{(s^2+9)^2} - \frac{4s}{(s^2+9)^2} + \frac{(2s^2)(2)(2s)}{(s^2+9)^3}$$

$$= \frac{8s^3}{(s^2+9)^3} - \frac{6s}{(s^2+9)^2}$$

• Prod rule
(uv)' = u'v + uv'
Applied to u = s,
v = (s^2+9)^-1.

Given $f(t) = e^{at} \sin t - e^{2t} \cos(4t)$, find $\mathcal{L}\{f(t)\}$.

$$\mathcal{L}\{e^{at} \sin t\} = \mathcal{L}\{\sin t\} \Big|_{s \rightarrow s-a}$$

$$= \frac{1}{s^2+1} \Big|_{s \rightarrow s-a}$$

$$= \frac{1}{(s-a)^2+1}$$

$$\mathcal{L}\{e^{2t} \cos(4t)\} = \mathcal{L}\{\cos(4t)\} \Big|_{s \rightarrow s-2}$$

$$= \frac{s}{s^2+4^2} \Big|_{s \rightarrow s-2}$$

$$= \frac{s-2}{(s-2)^2+16}$$

• Apply the shifting rule see below.

• Apply the shifting rule

• Identity of \mathcal{L}

$$= \boxed{\frac{1}{(s-1)^2+1} - \frac{s-2}{(s-2)^2+16}}$$

Shifting Theorem

$\mathcal{L}\{e^{at} f(t)\}$ equals $\mathcal{L}\{f(t)\}$ with s replaced by $s-a$.

$$\mathcal{L}\{e^{at} f(t)\} = \mathcal{L}\{f(t)\} \Big|_{s \rightarrow s-a}$$

Proof:

$$\mathcal{L}\{e^{at} f(t)\} = \int_0^{\infty} e^{-st} e^{at} f(t) dt$$

$$= \int_0^{\infty} e^{-(s-a)t} f(t) dt$$

$$= \int_0^{\infty} e^{-s't} f(t) dt \text{ with } s \rightarrow s-a$$

$$= \mathcal{L}\{f(t)\} \Big|_{s \rightarrow s-a}$$

• Direct transform
• $e^a e^b = e^{a+b}$

Solve for $f(t)$ in the equation $\mathcal{L}(f(t)) = \frac{3}{s^2} + \frac{s+1}{s^2+1}$

$$\mathcal{L}(f(t)) = \frac{3}{s^2} + \frac{s+1}{s^2+1} \quad \cdot \text{given}$$

$$= 3 \left(\frac{1}{s^2} \right) + (1) \left(\frac{s}{s^2+1} \right) + (1) \left(\frac{1}{s^2+1} \right) \quad \cdot \text{Arrange for table usage}$$

$$= 3 \mathcal{L}(t) + (1) \mathcal{L}(\cos t) + (1) \mathcal{L}(\sin t) \quad \cdot \text{Use table}$$

$$= \mathcal{L}(3t + \cos t + \sin t) \quad \cdot \text{linearity of } \mathcal{L}$$

$$f(t) = \boxed{3t + \cos t + \sin t} \quad \cdot \text{Apply Lerch's cancellation}$$

Evaluate $\mathcal{L}(1.1 - (t-1)\mathcal{L}(t+1)(t-2))$

$$\text{Let } F(s) = 1.1 - (t-1)\mathcal{L}(t+1)(t-2) \quad \cdot \text{Given}$$

$$= 1.1 - (t^2-1)(t-2) \quad \cdot \text{Multiply}$$

$$= 1.1 - t^3 + 2t^2 + t - 2$$

$$= -t^3 + 2t^2 + t - 0.9$$

Then

$$\mathcal{L}(f(t)) = \mathcal{L}(-t^3 + 2t^2 + t - 0.9)$$

$$= -\mathcal{L}(t^3) + 2\mathcal{L}(t^2) + \mathcal{L}(t) - 0.9\mathcal{L}(1) \quad \cdot \text{Linearity}$$

$$= -\frac{3!}{s^4} + 2\frac{2!}{s^3} + \frac{1}{s^2} - \frac{0.9}{s} \quad \cdot \text{Table}$$

$$= \boxed{-\frac{6}{s^4} + \frac{4}{s^3} + \frac{1}{s^2} - \frac{9/10}{s}}$$

Solve for $f(t)$ in the equality $\mathcal{L}(f(t)) = \frac{s+1}{(s-1)(s-2)}$

$$\mathcal{L}(f(t)) = \frac{s+1}{(s-1)(s+2)} \quad \cdot \text{Given}$$

$$= \frac{A}{s-1} + \frac{B}{s+2} \quad \cdot \text{Theory of partial fraction}$$

$$= A\mathcal{L}(e^t) + B\mathcal{L}(e^{-2t}) \quad \cdot \mathcal{L}(e^{at}) = \frac{1}{s-a}$$

$$= \mathcal{L}(Ae^t + Be^{-2t}) \quad \cdot \text{linearity of } \mathcal{L}$$

$$f(t) = Ae^t + Be^{-2t} \quad \cdot \text{Lerch's cancellation}$$

$$= \boxed{\left(\frac{2}{3}\right)e^t + \left(\frac{1}{3}\right)e^{-2t}}$$

\cdot Solve partial fraction problem, $A = \frac{2}{3}$, $B = \frac{1}{3}$.
(see below)

Solve $\frac{s+1}{(s-1)(s+2)} = \frac{A}{s-1} + \frac{B}{s+2}$ for A, B

cross-multiply by $s-1$ and then let $s-1 = 0$:

$$\frac{s+1}{s+2} = A + \frac{B}{s-1}$$

$$\frac{1+1}{1+2} = A + \frac{B}{1-1} \quad \cdot \text{Set } s-1 = 0$$

$$= A$$

Then $A = \frac{2}{3}$. Similarly, cross-multiply by $s+2$ and set $s+2 = 0$:

$$\frac{s+1}{s-1} = \frac{A}{s-1} + B$$

$$\frac{-2+1}{-2-1} = \frac{A}{-2-1} + B$$

Then $B = \frac{1}{3}$

Solve for $f(t)$ in the equation $\mathcal{L}(f(t)) = \arctan(1/s)$

The answer via $f(t) = \frac{\sin t}{t}$, obtained as follows.

$$\begin{aligned} \mathcal{L}((-t)f(t)) &= \frac{d}{ds} \mathcal{L}(f(t)) \\ &= \frac{d}{ds} (\arctan(1/s)) \\ &= \frac{-s^{-2}}{1 + (1/s)^2} \\ &= \frac{-1}{s^2 + 1} \\ &= -\mathcal{L}(\sin t) \\ &= \mathcal{L}(-\sin t) \end{aligned}$$

- s-diff rule
- Do the differentiation
- Tables
- Invariant of \mathcal{L}
- Apply Lebniz' cancellation law.
- Divide to find f .

s-diff rule. Multiplication of $f(t)$ by $(-t)$ differentiates the transform: $\mathcal{L}((-t)f(t)) = \frac{d}{ds} \mathcal{L}(f(t))$.

Proof:

$$\begin{aligned} \mathcal{L}((-t)f(t)) &= \int_0^{\infty} (-t)f(t)e^{-st} dt \\ &= \int_0^{\infty} f(t) (-t)e^{-st} dt \\ &= \int_0^{\infty} f(t) \frac{d}{ds} (e^{-st}) dt \\ &= \frac{d}{ds} \int_0^{\infty} f(t) e^{-st} dt \\ &= \frac{d}{ds} \mathcal{L}(f(t)). \end{aligned}$$

- Direct transform
- Possible because of convergence properties of the integrals.

Solve $\begin{cases} x'' + 3x' + 2x = \frac{1}{2} + e^{-3t} \\ x(0) = 0, x'(0) = 0 \end{cases}$ by the Laplace method.

Step 1

$$x'' + 3x' + 2x = \frac{1}{2} + e^{-3t}$$

$$\mathcal{L}(x'' + 3x' + 2x) = \mathcal{L}\left(\frac{1}{2} + e^{-3t}\right)$$

$$\mathcal{L}(x'') + 3\mathcal{L}(x') + 2\mathcal{L}(x) = \mathcal{L}\left(\frac{1}{2}\right) + \mathcal{L}(e^{-3t})$$

$$s^2 \mathcal{L}(x) - sx(0) - x'(0) = \frac{1}{2s} + \frac{1}{s+3}$$

$$+ 3[s\mathcal{L}(x) - x(0)] + 2\mathcal{L}(x) = \frac{1}{2s} + \frac{1}{s+3}$$

$$[s^2 + 3s + 2]\mathcal{L}(x) = sx(0) + x'(0) + 3x(0) + \frac{1}{2s} + \frac{1}{s+3}$$

Char Equation appears!

$$[s^2 + 3s + 2]\mathcal{L}(x) = \frac{1}{2s} + \frac{1}{s+3}$$

$$\mathcal{L}(x) = \frac{3s + 3}{(2s)(s+3)(s^2 + 3s + 2)}$$

$$= \frac{3/2}{s(s+3)(s+2)}$$

End of step 1. Found $\mathcal{L}(x)$ explicitly.

Given DE

Take Laplace of both sides
[Mult both sides by e^{-st} and integrate $t=0$ to $t=\infty$]

$$\mathcal{L}(ax+by) = a\mathcal{L}(x) + b\mathcal{L}(y)$$

[Integral of a sum = sum of integrals; const go through the integral \mathcal{L}]

$$\text{Derivative Theorem} \\ \mathcal{L}(y') = s\mathcal{L}(y) - y(0)$$

move terms to right, collect factor $\mathcal{L}(x)$ on LHS

$$\text{use } x(0)=0, x'(0)=0, \\ \mathcal{L}(1) = \frac{1}{s}, \mathcal{L}(e^{at}) = \frac{1}{s-a}$$

Divide to isolate $\mathcal{L}(x)$ on the left.

Factor, cancel $s+1$ top and bottom.

Step 2.

The objective is to leave $f(s)$ on the left unchanged, but change P_2 this to look like $f(s)$ (something)

$$\begin{aligned}
 f(x) &= \frac{3/2}{s(s+3)(s+2)} \\
 &= \frac{A}{s} + \frac{B}{s+3} + \frac{C}{s+2} \\
 &= \frac{1/4}{s} + \frac{+4/2}{s+3} + \frac{-3/4}{s+2} \\
 &= \frac{1}{4} f(1) + \frac{1}{2} f(-3) - \frac{3}{4} f(-2) \\
 &= f\left(\frac{1}{4} + \frac{1}{2} e^{-3t} - \frac{3}{4} e^{-2t}\right)
 \end{aligned}$$

Step 3.

Apply Lerch's Theorem to find x

$$x(t) = \frac{1}{4} + \frac{1}{2} e^{-3t} - \frac{3}{4} e^{-2t}$$

Transient and steady-state

$$x_{ss}(t) = \frac{1}{4}$$

$$x_{tr}(t) = \frac{1}{2} e^{-3t} - \frac{3}{4} e^{-2t}$$

The end!

Lerch's Theorem will be applied:
 $f(x) = f(x_1) \Rightarrow x_1 = x_2 = 1^2$, the f cancels on both sides.

Theory of partial fraction from college algebra and calculus I.

By Heaviside's method

$$By f(e^{at}) = \frac{1}{s-a} !$$

By Linearity (again).

The f cancels on each side of the previous relation [equivalent to the inverse Laplace transform]

Def:

$$\begin{aligned}
 x &= x_{ss} + x_{tr} \\
 \text{and } x_{tr} &\rightarrow 0 \text{ as } t \rightarrow \infty.
 \end{aligned}$$

10.1-3 Find $f(t)$ for $F(s) = e^{-2t+\pi}$

Example.

Find $f(t)$ for $F(s) = e^{-2t+\pi}$

$$\begin{aligned}
 f(s) &= f(e^{-2t} e^{\pi}) \\
 &= e^{\pi} f(e^{-2t}) \\
 &= e^{\pi} \frac{1}{s-(-2)} \\
 &= \frac{e^{\pi}}{s+2}
 \end{aligned}$$

use $e^{at+b} = e^a e^{bt}$
 Linearity of f
 Tables

10.1-5 Find $f(t)$ for $F(s) = \sinh(t)$.

Hint: $\sinh(u) = \frac{1}{2} e^u - \frac{1}{2} e^{-u}$ by definition.

10.1-17 Find $f(t)$ for $F(s) = \cos^2(2t)$

Hint: The table contains cosines and sines but not $\cos^2(2t)$. By trig identities,

$$\begin{aligned}
 \cos(2\theta) &= \cos^2 \theta - \sin^2 \theta \\
 &= 2 \cos^2 \theta - 1.
 \end{aligned}$$

Hence,

$$\cos(4t) = 2 \cos^2(2t) - 1$$

This identity implies $f(t) = \frac{1}{2} + \frac{1}{2} \cos(4t)$, or sum of terms already present in the table.

10.1-27 Find $f(t)$ given $\mathcal{L}(f) = \frac{3}{s-4}$

Example: Find $f(t)$ given $\mathcal{L}(f) = \frac{1}{s} - \frac{2}{s^2} + \frac{4}{s-16}$

$$\mathcal{L}(f) = \frac{1}{s} + \frac{-2}{s^2} + \frac{4}{s-16}$$

$$= \mathcal{L}(1) + (-2)\mathcal{L}(t) + 4\mathcal{L}(e^{16t}) \quad \text{by tables}$$

$$= \mathcal{L}(1 - 2t + 4e^{16t}) \quad \text{Linearity}$$

$$f(t) = 1 - 2t + 4e^{16t} \quad \text{Levi's Thm applied}$$

10.2-7 Solve by Laplace method $\begin{cases} x'' + x = \cos(3t) \\ x(0) = 1, x'(0) = 0 \end{cases}$

Details: Apply \mathcal{L} across the DE and use Laplace rules to obtain the equation (see Ex 2 in 10.1)

$$\mathcal{L}(x) = \frac{1}{s^2+1} [s + \mathcal{L}(\cos 3t)] \quad \text{fill in the details!}$$

$$= \frac{1}{s^2+1} [s + \frac{s}{s^2+9}]$$

$$= \frac{s}{s^2+1} + \frac{s}{(s^2+1)(s^2+9)}$$

$$= \mathcal{L}(\cos t) + \frac{s}{(s^2+1)(s^2+9)}$$

To finish, expand the fraction on the right as partial fractions

$$\frac{as+b}{s^2+1} + \frac{cs+3d}{s^2+9} \quad \text{which equals}$$

$$\mathcal{L}(a \cos t + b \sin t + c \sin 3t + d \cos 3t).$$

10.2-11 Solve by the Laplace method

$$\begin{cases} x' = 2x + y \\ y' = 6x + 3y \\ x(0) = 1, y(0) = -2 \end{cases}$$

solution details: Transform each DE to obtain equations

$$\begin{cases} s\mathcal{L}(x) - 1 = 2\mathcal{L}(x) + \mathcal{L}(y) \\ s\mathcal{L}(y) + 2 = 6\mathcal{L}(x) + 3\mathcal{L}(y) \end{cases}$$

write as a linear system $A\mathbf{z} = \mathbf{b}$ where $\mathbf{z} = \begin{bmatrix} \mathcal{L}(x) \\ \mathcal{L}(y) \end{bmatrix}$ and then solve it to obtain

$$\mathcal{L}(x) = \frac{s-5}{s(s-5)} = \frac{1}{s}$$

$$\mathcal{L}(y) = \frac{-2}{s}$$

Apply Table methods to get $x = 1, y = -2$.

10.2-15 project Solve by the Laplace method

$$\begin{cases} x'' + x' + y' + 2x - y = 0 \\ y'' + x' + y' + 4x - 2y = 0 \\ x(0) = y(0) = 1, x'(0) = y'(0) = 0 \end{cases}$$

Hint: Transform the DEs to obtain the system

$$\begin{bmatrix} s^2+s+2 & s-1 \\ s+4 & s^2+s-2 \end{bmatrix} \begin{bmatrix} \mathcal{L}(x) \\ \mathcal{L}(y) \end{bmatrix} = \begin{bmatrix} s+1+1 \\ s+1+1 \end{bmatrix}$$

Solve by Cramer's rule or equivalent to get $\mathcal{L}(x) = \frac{s^2+3s+1}{s(s^2+3s)}$
 $\mathcal{L}(y) = \text{similar}$. Expand in partial fractions to get the book's answer, e.g., $x = \frac{2}{3} + \frac{1}{3}e^{-3t/2}(\cos \frac{\sqrt{3}t}{2} + \sqrt{3} \sin \frac{\sqrt{3}t}{2})$

10.3-3 Find $f(t)$ for $F(s) = e^{-2t} \sin(3\pi t)$

Example. Find $f(t)$ for $f(t) = e^{-\pi t} \cos(\pi t)$

$$\begin{aligned}
 f(t) &= \mathcal{L}(e^{-\pi t} \cos \pi t) \\
 &= \mathcal{L}(\cos \pi t) \Big|_{s \mapsto s+\pi} \quad \text{Shift Theorem} \\
 &= \frac{s}{s^2 + \pi^2} \Big|_{s \mapsto s+\pi} \quad \text{Table} \\
 &= \frac{s+\pi}{(s+\pi)^2 + \pi^2}
 \end{aligned}$$

10.3-7 Find $f(t)$ given $\mathcal{L}(f) = \frac{1}{s^2 + 4s + 4}$

Example. Find $f(t)$ given $\mathcal{L}(f) = \frac{1}{s^2 + 5s + 4}$

$$\begin{aligned}
 f(t) &= \frac{1}{s^2 + 5s + 4} \\
 &= \frac{1}{(s+1)(s+4)} \\
 &= \frac{A}{s+1} + \frac{B}{s+4} \quad \text{partial fractions} \\
 &= \mathcal{L}(Ae^{-t} + Be^{-4t}) \quad \text{Table}
 \end{aligned}$$

$$\begin{aligned}
 f(t) &= Ae^{-t} + Be^{-4t} \\
 &= \frac{1}{3}e^{-t} - \frac{1}{3}e^{-4t}
 \end{aligned}$$

$A = \frac{1}{3}$, $B = -\frac{1}{3}$ by Heaviside's method.

10.3-19 Find $f(t)$ given $\mathcal{L}(f) = \frac{s^2 + 2s}{s^4 + 5s^2 + 4}$

$$\begin{aligned}
 \text{Hint: } \mathcal{L}(f) &= \frac{s^2 + 2s}{(s^2 + 4)(s^2 + 1)} \\
 &= \frac{as + ab}{s^2 + 4} + \frac{cs + d}{s^2 + 1} \quad \text{partial fraction theory} \\
 &= \mathcal{L}(a \cos 2t + b \sin 2t + c \cos t + d \sin t)
 \end{aligned}$$

The problem thus reduces to computing constants a, b, c, d .

10.3-29 Solve by the Laplace method

$$x'' - 4x = 3t, \quad x(0) = e, \quad x'(0) = e$$

Hint. Transform the DE to get

$$\begin{aligned}
 \mathcal{L}(x) &= \frac{3}{s^2(s^2 - 4)} \\
 &= \frac{3}{s^2(s-2)(s+2)} \\
 &= \frac{a}{s^2} + \frac{b}{s} + \frac{c}{s-2} + \frac{d}{s+2} \quad \text{partial fractions} \\
 &= \mathcal{L}(at + b + ce^{2t} + de^{-2t})
 \end{aligned}$$

It remains to show details about the formula for $\mathcal{L}(x)$ and College algebra to find the constants a, b, c, d . By Heaviside's Theorem,

$$x = at + b + ce^{2t} + de^{-2t}$$

The book answer computes $\sin h(2t) = \frac{1}{2}e^{2t} - \frac{1}{2}e^{-2t}$.

Convolution Theorem

Given two functions $f(x)$, $g(x)$ of exponential order,
Then

$$\mathcal{L}(f(x)) \mathcal{L}(g(x)) = \mathcal{L}\left(\int_0^t f(x)g(t-x)dx\right)$$

Example. Solve for $y(x)$ in the equation $f(y(x)) = \frac{1}{s^2(s-1)}$

Solution by convolution:

$$\mathcal{L}(y(x)) = \frac{1}{s^2} \frac{1}{s-1}$$

$$= f(x) g(x)$$

$$= \mathcal{L}\left(\int_0^t x e^{-x} dx\right)$$

$$= \mathcal{L}\left(e^t \int_0^t x e^{-x} dx\right)$$

$$= \mathcal{L}\left(e^t (1 - e^{-t} - t e^{-t})\right)$$

$$= \mathcal{L}\left(e^t - 1 - t\right)$$

$$y(x) = e^x - 1 - x$$


by Laplace's convolution law.

check: $f(y) = \frac{1}{s-1} - \frac{1}{s} - \frac{1}{s^2}$

$$= \frac{-s+1-s^2+s+s^2}{s^2(s-1)}$$

$$= \frac{1}{s^2(s-1)}$$

Example. Calculate $\mathcal{L}(f)$ for the periodic extension of $H(t) + H(t-2)$ on $0 \leq t \leq 4$, of period 4.



The base function is a step, $f=1$ on $0 \leq t \leq 2$, $f=2$ on $2 \leq t \leq 4$.

$$\mathcal{L}(f) = \frac{\int_0^p e^{-st} f(t) dt}{1 - e^{-ps}}$$

$$p=4$$

$$f(t) = 1 + H(t-2)$$

$$= \frac{\int_0^4 e^{-st} (1 + H(t-2)) dt}{1 - e^{-4s}}$$

$$= \left(\int_0^2 e^{-st} dt + \int_2^4 e^{-st} dt \right) / (1 - e^{-4s})$$

$$= \frac{1 + \frac{e^{-2s} - 2e^{-4s}}{s}}{s(1 - e^{-4s})}$$

Final answer

Remark

The integration above used

$$\int_0^4 e^{-st} f(t) dt = \int_0^2 e^{-st} f(t) dt + \int_2^4 e^{-st} f(t) dt$$

$$= \int_0^2 e^{-st} (1) dt + \int_2^4 e^{-st} (2) dt$$

Generally, step function integrations require such a splitting of integrals, and subsequent replacement of simple expressions within the integrand, in order to be successfully integrated.

Historical Origins of The Laplace Transform

1822

Jean-Baptiste Joseph Fourier published "The analytical theory of heat" in Paris.

Studied heat conduction, for an insulated bar placed on the side with ice, the heat $u(x,t)$ at position x ($0 \leq x \leq \pi$) and time $t \geq 0$ is given as

$$u(x,t) = \sum_{n=1}^{\infty} C_n \sin(nx) e^{-n^2 t}$$

Fourier claimed that any initial heat distribution $u(x,0) = f(x)$ could be written as

$$f(x) = \sum_{n=0}^{\infty} a_n \cos(nx) + b_n \sin(nx)$$

Fourier-series

Dirichlet made the work rigorous by providing hypotheses on f that made it true (1804-1859).

Fourier's ideas were applied to vibrations of strings, with the vibration excursion from equilibrium $u(x,t)$ again being represented in Fourier's natural sine-cosine coordinate system.

The Fourier Integral was invented to handle continuous spectra and non-periodic behavior.

$$f(x) = \int_0^{\infty} (A(\omega) \cos(\omega x) + B(\omega) \sin(\omega x)) d\omega$$

$$A(\omega) = \frac{1}{\pi} \int_{-\infty}^{\infty} f(v) \cos(\omega v) dv, \quad B(\omega) = \frac{1}{\pi} \int_{-\infty}^{\infty} f(v) \sin(\omega v) dv$$

Historical origins-2

1890

Heaviside's operational calculus evolved into modern Laplace Theory.

The Complex Fourier Integral was derived from Euler's classic formula $e^{i\theta} = \cos\theta + i\sin\theta$.

$$f(x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} F(\omega) e^{i\omega x} d\omega$$

$$F(\omega) = \int_{-\infty}^{\infty} f(x) e^{-i\omega x} dx$$

Fourier Transform

Oliver Heaviside invented an operational method for solving differential equations, very mysterious. The explanation of why it worked leads to the following mathematical object:

$$\int_0^{\infty} e^{-st} f(t) dt$$

The Laplace Transform

For a function $f(t) = 0$ for $t < 0$, this is the same as

$$\int_{-\infty}^{\infty} e^{-i\omega t} f(t) dt$$

The Fourier Transform revisited

with $i\omega$ replaced by s .

Lerch proved a cancellation law that allowed the special transform to be used as an alternate method for a differential equation. It reads:

$$\mathcal{L}\{f\} \mathcal{L}\{g\} = \mathcal{L}\{fg\} \Rightarrow f(t) = g(t)$$