

## ANSWERS

1. (5 points) What is the magnitude of the vector  $\begin{pmatrix} 1 \\ 2 \end{pmatrix}$ ?

**Answer:**

$$\left\| \begin{pmatrix} 1 \\ 2 \end{pmatrix} \right\| = \sqrt{\begin{pmatrix} 1 \\ 2 \end{pmatrix} \cdot \begin{pmatrix} 1 \\ 2 \end{pmatrix}} = \sqrt{5}$$

2. (5 points) What is the unit vector that has the same direction as  $\begin{pmatrix} 1 \\ 2 \end{pmatrix}$ ?

**Answer:**

$$\frac{\begin{pmatrix} 1 \\ 2 \end{pmatrix}}{\left\| \begin{pmatrix} 1 \\ 2 \end{pmatrix} \right\|} = \begin{pmatrix} \frac{1}{\sqrt{5}} \\ \frac{2}{\sqrt{5}} \end{pmatrix}$$

3. (5 points) What is the coefficient matrix corresponding to the following system of linear equations?

$$\begin{aligned} x_1 + x_2 + x_3 + x_4 &= b_1 \\ 4x_1 + 2x_2 + 3x_3 &= b_2 \\ 4x_3 + 5x_4 &= b_3 \end{aligned}$$

**Answer:**

$$\begin{pmatrix} 1 & 1 & 1 & 1 \\ 4 & 2 & 3 & 0 \\ 0 & 0 & 4 & 5 \end{pmatrix}$$

4. (5 points) Let  $A$  be a  $3 \times 4$  matrix. What is the elimination matrix that replaces Row 2 of  $A$  with  $Row2 - Row1$  and replaces  $Row3$  of  $A$  by  $Row3 - 2Row1$ ?

Answer:

$$\begin{pmatrix} 1 & 0 & 0 \\ -1 & 1 & 0 \\ -2 & 0 & 1 \end{pmatrix}$$

5. (5 points) What is the inverse of the following matrix?

$$E = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & -5 & 1 & 0 \\ 0 & 0 & 1 & 0 & 1 \end{pmatrix}$$

Answer:

$E$  is an elimination matrix. It replaces Row 4 with Row 4 - 5 Row 3 and replaces Row 5 with Row 3 + Row 5. The inverse is the elimination matrix that reverses these steps, replacing Row 4 with Row 4 + 5Row 3 and replacing Row 5 with Row 5 - Row 3.

$$E^{-1} = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 5 & 1 & 0 \\ 0 & 0 & -1 & 0 & 1 \end{pmatrix}$$

6. (5 points) Describe in word the effect of multiplying  $EA$  where  $E$  is the matrix in the previous problem and  $A$  is a  $5 \times 5$  matrix.

Answer:

$E$  is an elimination matrix. It replaces Row 4 with Row 4 - 5 Row 3 and replaces Row 5 with Row 3 + Row 5.

7. (15 points) Let  $A = \begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix}$ . Let  $B = \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}$ . Let  $C = \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix}$ . Calculate the following:

$$A(BC)$$

$$\begin{aligned}
 &(AB)C \\
 &C^2 \\
 &2A + B - C \\
 &A(B - C)
 \end{aligned}$$

**Answer:**

$$\begin{aligned}
 A(BC) &= \begin{pmatrix} 6 & 3 \\ 14 & 7 \end{pmatrix} \\
 (AB)C &= \begin{pmatrix} 6 & 3 \\ 14 & 7 \end{pmatrix} \\
 C^2 &= \begin{pmatrix} 2 & 1 \\ 1 & 1 \end{pmatrix} \\
 2A + B - C &= \begin{pmatrix} 2 & 4 \\ 6 & 9 \end{pmatrix} \\
 A(B - C) &= \begin{pmatrix} 0 & 2 \\ 0 & 4 \end{pmatrix}
 \end{aligned}$$

**8. (20 points)** Prove or disprove that the following sets of vectors form a basis for the indicated vector space:

$$\begin{pmatrix} 1 \\ 2 \end{pmatrix}, \begin{pmatrix} 2 \\ 2 \end{pmatrix} \text{ in } \mathbb{R}^2$$

$$\begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 2 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ -1 \\ 1 \end{pmatrix}, \begin{pmatrix} -1 \\ 1 \\ 1 \end{pmatrix} \text{ in } \mathbb{R}^3$$

**Answer:**

$$\begin{pmatrix} 1 \\ 2 \end{pmatrix}, \begin{pmatrix} 2 \\ 2 \end{pmatrix} \text{ in } \mathbb{R}^2$$

These two vectors form a basis for  $\mathbb{R}^2$ . These two vectors are not scalar multiples of each other, so they are linearly independent.  $\mathbb{R}^2$  is two dimensional, so an two linearly independent vectors form a basis.

$$\begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 2 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ -1 \\ 1 \end{pmatrix}, \begin{pmatrix} -1 \\ 1 \\ 1 \end{pmatrix} \text{ in } \mathbb{R}^3$$

These four vectors do not form a basis. A basis consists of linearly independent vectors.  $\mathbb{R}^3$  is three dimensional, so no set of more than three vectors can ever be linearly independent in  $\mathbb{R}^3$ .

**9. (25 points)** Find all solutions to the equation  $A\bar{x} = \bar{b}$  for

$$A = \begin{pmatrix} 1 & 2 & 3 & 4 \\ 5 & 6 & 7 & 8 \end{pmatrix}$$

$$\bar{b} = \begin{pmatrix} 0 \\ 4 \end{pmatrix}$$

**Answer:**

The augmented matrix for this system of equations is:

$$\begin{pmatrix} 1 & 2 & 3 & 4 & 0 \\ 5 & 6 & 7 & 8 & 4 \end{pmatrix}$$

Put this is reduced row echelon form by performing Gauss-Jordan Elimination:

$$\begin{pmatrix} 1 & 0 \\ -5 & 1 \end{pmatrix} \begin{pmatrix} 1 & 2 & 3 & 4 & 0 \\ 5 & 6 & 7 & 8 & 4 \end{pmatrix} = \begin{pmatrix} 1 & 2 & 3 & 4 & 0 \\ 0 & -4 & -8 & -12 & 4 \end{pmatrix}$$

$$\begin{pmatrix} 1 & \frac{1}{2} \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 2 & 3 & 4 & 0 \\ 0 & -4 & -8 & -12 & 4 \end{pmatrix} = \begin{pmatrix} 1 & 0 & -1 & -2 & 2 \\ 0 & -4 & -8 & -12 & 4 \end{pmatrix}$$

$$\begin{pmatrix} 1 & 0 \\ 0 & -\frac{1}{4} \end{pmatrix} \begin{pmatrix} 1 & 0 & -1 & -2 & 2 \\ 0 & -4 & -8 & -12 & 4 \end{pmatrix} = \begin{pmatrix} 1 & 0 & -1 & -2 & 2 \\ 0 & 1 & 2 & 3 & -1 \end{pmatrix}$$

Solutions to the original equation are the same as solutions to the new equation :

$$\begin{pmatrix} 1 & 0 & -1 & -2 \\ 0 & 1 & 2 & 3 \end{pmatrix} \bar{x} = \begin{pmatrix} 2 \\ -1 \end{pmatrix}$$

Columns 1 and 2 are pivot columns; columns 3 and 4 are free columns, so  $x_3$  and  $x_4$  are the free variables.

A particular solution is obtained by setting  $x_3 = x_4 = 0$  and solving for  $x_1$  and  $x_2$ .

$$\begin{pmatrix} 1 & 0 & -1 & -2 \\ 0 & 1 & 2 & 3 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ 0 \\ 0 \end{pmatrix} = \begin{pmatrix} 2 \\ -1 \end{pmatrix}$$

The particular solution is

$$\bar{x}_p = \begin{pmatrix} 2 \\ -1 \\ 0 \\ 0 \end{pmatrix}$$

The other solutions differ from  $\bar{x}_p$  by vectors in the nullspace of  $A$ . A basis for the nullspace is given by the “special solutions” obtained by setting one free variable equal to 1, the others equal to 0, and solving for the non-free variables.

$\bar{s}_3$  is the special solution obtained by setting  $x_3 = 1$  and  $x_4 = 0$  and solving

$$\begin{pmatrix} 1 & 0 & -1 & -2 \\ 0 & 1 & 2 & 3 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ 1 \\ 0 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

$$\text{So } \bar{s}_3 = \begin{pmatrix} 1 \\ -2 \\ 1 \\ 0 \end{pmatrix}$$

Similarly,  $s_4$  is obtained by setting  $x_3 = 0$  and  $x_4 = 1$  and solving

$$\begin{pmatrix} 1 & 0 & -1 & -2 \\ 0 & 1 & 2 & 3 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ 0 \\ 1 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

$$\text{So } \bar{s}_4 = \begin{pmatrix} 2 \\ -3 \\ 0 \\ 1 \end{pmatrix}$$

Finally, all solutions are of the form  $\bar{x}_p + x_3\bar{s}_3 + x_4\bar{s}_4$ , so all solutions to the equation are all vectors:

$$\bar{x} = \begin{pmatrix} 2 \\ -1 \\ 0 \\ 0 \end{pmatrix} + x_3 \begin{pmatrix} 1 \\ -2 \\ 1 \\ 0 \end{pmatrix} + x_4 \begin{pmatrix} 2 \\ -3 \\ 0 \\ 1 \end{pmatrix}$$

where  $x_3$  and  $x_4$  are arbitrary real numbers.

**10. (20 points)** Prove that a system of 4 linear equations in 5 unknowns has either 0 or infinitely many solutions.

**Answer:**

There are more unknowns than equations. Either the equations are inconsistent and there are no solutions, or there is a free variable. We may choose this free variable to be any real number and find a corresponding solution, so this is infinitely many solutions.

No new questions beyond this point.