

Examples: Solving n th Order Equations

- Euler Solution Atoms and Euler Base Atoms
- L. Euler's Theorem
- The Atom List
- First Order. Solve $2y' + 5y = 0$.
- Second Order. Solve the equations $y'' + 2y' + y = 0$, $y'' + 3y' + 2y = 0$ and $y'' + 2y' + 5y = 0$.
- Third Order. Solve $y''' - y' = 0$ and $y''' - y'' = 0$.
- Fourth Order. Solve $y^{(4)} - y^{(2)} = 0$.

Atoms

Assume symbols a and b are real constants, with $a \neq 0$ and $b > 0$.

An Euler **base atom** is one of the functions

$$1, \cos bx, \sin bx, \\ e^{ax}, e^{ax} \cos bx, e^{ax} \sin bx.$$

An Euler solution **atom** is a base atom multiplied by a power x^n , where $n \geq 0$ is an integer.

$$\text{Euler solution atom} = x^n (\text{Euler base atom})$$

Examples

Atoms: $1, \cos(3x/2), x^3, x \cos(2x), x^2 e^{2x} \sin(3x), \cos(\pi x), e^{e^2 x}$

Not an atom: $2, -x, x^{3/2}, \tan x, \sin(x^2), e^{x^2}, \frac{x}{1+x^2}, \ln|x|, \sinh x$

Euler's Theorem

Theorem 1 (L. Euler)

The function $y = x^m e^{r_1 x}$ is a solution of a constant-coefficient linear homogeneous differential equation of the n th order if and only if $(r - r_1)^{m+1}$ divides the characteristic polynomial.

Euler's theorem is used to construct solutions of the n th order differential equation. The solutions so constructed are n distinct Euler solution atoms, hence independent. Picard's theorem implies the list of Euler solution atoms is a basis for the solution space.

Euler's Theorem in Words

To construct solutions of homogeneous constant-coefficient differential equations, use Euler's Theorem as follows.

- Find the roots of the characteristic equation.
 - For each real root r , the exponential solution e^{rx} is an Euler base atom solution.
 - For each complex conjugate pair of roots $a \pm bi$, $b > 0$, the functions $e^{ax} \cos bx$, $e^{ax} \sin bx$ are Euler base atom solutions.
- Multiply each base atom by the powers $1, x, x^2, \dots, x^{k-1}$, where k is the multiplicity of the root. The atoms so found are a basis of solutions for the differential equation.

The Atom List

1. If r_1 is a real root, then the Euler base atom for r_1 is $e^{r_1 x}$. The root r_1 has multiplicity k provided factor $(r - r_1)^k$ divides the characteristic polynomial, but factor $(r - r_1)^{k+1}$ does not. Multiply the base atom by powers $1, x, \dots, x^{k-1}$ (a total of k terms) to obtain the atom list

$$e^{r_1 x}, x e^{r_1 x}, \dots, x^{k-1} e^{r_1 x}.$$

2. If $r_1 = a + ib$ ($b > 0$ assumed) and its conjugate $r_2 = a - ib$ are roots of the characteristic equation, then the Euler base atoms for this pair of roots (both r_1 and r_2 counted) are

$$e^{ax} \cos bx, \quad e^{ax} \sin bx.$$

For a root of multiplicity k , these real atoms are multiplied by atoms $1, \dots, x^{k-1}$ to obtain the complete list of $2k$ atoms

$$\begin{aligned} &e^{ax} \cos bx, \quad x e^{ax} \cos bx, \quad \dots, \quad x^{k-1} e^{ax} \cos bx, \\ &e^{ax} \sin bx, \quad x e^{ax} \sin bx, \quad \dots, \quad x^{k-1} e^{ax} \sin bx. \end{aligned}$$

1 Example (First Order) Solve $2y' + 5y = 0$ by Euler's method, verifying $y_h = c_1 e^{-5x/2}$.

Solution

$2y' + 5y = 0$ Given differential equation.

$2r + 5 = 0$ Characteristic equation. Use the shortcut $y^{(n)} \rightarrow r^n$.

$r = -5/2$ Exactly one real root.

Atom = $e^{-5x/2}$ For a real root r , the Euler base atom is e^{rx} .

$y_h = c_1 e^{-5x/2}$ The general solution y_h is written by multiplying the atom list by constant c_1 .

2 Example (Second Order I) Solve $y'' + 2y' + y = 0$ by Euler's method, showing that $y_h = c_1e^{-x} + c_2xe^{-x}$.

Solution

$$y'' + 2y' + y = 0$$

Given differential equation.

$$r^2 + 2r + 1 = 0$$

Characteristic equation. Use shortcut $y^{(n)} \rightarrow r^n$.

$$r = -1, -1$$

Exactly two real roots.

$$\text{Base Atom} = e^{-x}$$

For a real root r_1 , the Euler base atom is e^{r_1x} .

$$\text{Atoms} = e^{-x}, xe^{-x}$$

For a root of multiplicity 2, multiply the base atom by powers 1, x , to create 2 Euler solution atoms.

$$y_h = c_1e^{-x} + c_2xe^{-x}$$

The general solution y_h is written by multiplying the atom list by constants c_1, c_2 .

3 Example (Second Order II) Solve $y'' + 3y' + 2y = 0$ by Euler's method, showing $y_h = c_1e^{-x} + c_2e^{-2x}$.

Solution

$$y'' + 3y' + 2y = 0$$

$$r^2 + 3r + 2 = 0$$

$$r = -1, -2$$

$$\text{Base Atoms} = \begin{matrix} e^{-x} \\ e^{-2x} \end{matrix},$$

$$\text{Atoms} = e^{-x}, e^{-2x}$$

$$y_h = c_1e^{-x} + c_2e^{-2x}$$

Given differential equation.

Characteristic equation. Use shortcut $y^{(n)} \rightarrow r^n$.

Factorization $(r + 2)(r + 1) = 0$.

For a real root r_1 , the Euler base atom is e^{r_1x} .

For a root of multiplicity 1, multiply the base atom by 1, to create one term. The number of terms created from one base atom always equals the multiplicity of the root.

The general solution y_h is written by multiplying the atom list by constants c_1, c_2 .

4 Example (Second Order III) Solve $y'' + 2y' + 5y = 0$ by Euler's method, showing $y_h = c_1 e^{-x} \cos 2x + c_2 x e^{-x} \sin 2x$.

Solution

$$y'' + 2y' + 5y = 0$$

$$r^2 + 2r + 5 = 0$$

$$r = -1 + 2i, -1 - 2i$$

$$\text{Base Atoms} = \begin{matrix} e^{-x} \cos 2x, \\ e^{-x} \sin 2x \end{matrix}$$

$$\text{Atoms} = e^{-x} \cos 2x, e^{-x} \sin 2x$$

$$y_h = c_1 e^{-x} \cos 2x + c_2 e^{-x} \sin 2x$$

Given differential equation.

Characteristic equation.

Shortcut $y^{(n)} \rightarrow r^n$ used.

Factorization $(r + 1)^2 + 4 = 0$ implies $r + 1 = \pm 2\sqrt{-1} = \pm 2i$.

For complex conjugate roots $r = a \pm ib$, the Euler base atoms are $e^{ax} \cos bx$ and $e^{ax} \sin bx$.

Multiply the base atoms by 1, because the multiplicity of root $1 + 2i$ is 1. In general, multiply by 1, x , \dots , x^{k-1} , where k is the root multiplicity.

Multiply the atom list by constants c_1 , c_2 . This is the general solution.

5 Example (Third Order I) Solve $y''' - y' = 0$ by Euler's method, showing $y_h = c_1 + c_2e^x + c_3e^{-x}$.

Solution

$$y''' - y' = 0$$

$$r^3 - r = 0$$

$$r = 0, 1, -1$$

$$\text{Base Atoms} = 1, e^{-x}, e^x$$

$$\text{Atoms} = 1, e^{-x}, e^x$$

$$y_h = c_1 + c_2e^{-x} + c_3e^x$$

Given differential equation.

Characteristic equation. Use $y^{(n)} \rightarrow r^n$.

Factorization $r(r + 1)(r - 1) = 0$.

For a real root r_1 , the Euler base atom is e^{r_1x} .

Each root has multiplicity 1. Multiply each base atom by 1. Generally, multiply by $1, x, \dots, x^{k-1}$, where k is the root multiplicity.

The general solution y_h is written by multiplying the atom list by constants c_1, c_2, c_3 .

6 Example (Third Order II) Solve $y''' - y'' = 0$ by Euler's method, showing $y_h = c_1 + c_2x + c_3e^x$.

Solution

$$y''' - y'' = 0$$

$$r^3 - r^2 = 0$$

$$r = 0, 0, 1$$

$$\text{Base Atoms} = 1, e^x$$

$$\text{Atoms} = 1, x, e^x$$

$$y_h = c_1 + c_2x + c_3e^x$$

Given differential equation.

Characteristic equation. Use $y^{(n)} \rightarrow r^n$.

Factorization $r^2(r - 1) = 0$.

For a real root r_1 , the Euler base atom is e^{r_1x} . Then the Euler base atoms are e^{0x}, e^{1x} , written as $1, e^x$.

Because $r = 0$ has multiplicity **2**, then multiply Euler base atom $1 (=e^{0x})$ by powers $1, x$. Root $r = 1$ with multiplicity **1** implies base atom e^x is multiplied by **1**. The total number of atoms created is $2 + 1 = 3$, which is the sum of the root multiplicities.

The general solution y_h is written by multiplying the atom list by constants c_1, c_2, c_3 .

7 Example (Fourth Order) Solve $y^{(4)} - y^{(2)} = 0$ by Euler's method, showing that the general solution is $y_h = c_1 + c_2x + c_3e^x + c_4e^{-x}$.

Solution

$$y^{(4)} - y^{(2)} = 0$$

$$r^4 - r^2 = 0$$

$$r = 0, 0, 1, -1$$

$$\text{Base Atoms} = 1, e^x, e^{-x}$$

$$\text{Atoms} = 1, x, e^x, e^{-x}$$

$$y_h = c_1 + c_2x + c_3e^x + c_4e^{-x}$$

Given differential equation.

Characteristic equation. Use $y^{(n)} \rightarrow r^n$.

Factorization $r^2(r - 1)(r + 1) = 0$.

For a real root r_1 , the Euler base atom is e^{r_1x} . The base atoms e^{0x} , e^{1x} , e^{-1x} are written 1 , e^x , e^{-x} .

Multiply the multiplicity 2 base atom 1 by powers 1 , x and each of the multiplicity 1 base atoms e^x , e^{-x} by 1 . The total number of atoms created is $2 + 1 + 1 = 4$, which is the sum of the root multiplicities.

Multiply the atom list by constants c_1 , c_2 , c_3 , c_4 and add. This is the general solution.