

3150 S 2014
Sample Exam 2

SOLUTIONS

Fourier
series
problem 1

1. (Periodic Functions)

- (a) [30%] Find the period of $f(x) = \sin(x)\cos(2x) + \sin(2x)\cos(x)$.
- (b) [40%] Let $p = 5$. If $f(x)$ is the odd $2p$ -periodic extension to $(-\infty, \infty)$ of the function $f_0(x) = 100xe^{10x}$ on $0 \leq x \leq p$, then find $f(11.3)$. The answer is not to be simplified or evaluated to a decimal.
- (c) [30%] Mark the expressions which are periodic with letter **P**, those odd with **O** and those even with **E**.

$$\sin(\cos(2x)) \quad \ln|2 + \sin(x)| \quad \sin(2x)\cos(x) \quad \frac{1 + \sin(x)}{2 + \cos(x)}$$

Answer:

- (a) $f(x) = \sin(x + 2x)$ by a trig identity. Then period = $2\pi/3$.
- (b) $f(11.3) = f(11.3 - p - p) = f(1.3) = f_0(1.3) = 130e^{13}$.
- (c) All are periodic of period 2π , satisfying $f(x + 2\pi) = f(x)$. The first is even and the third is odd.

Blackboard photos week 13, 4 April

2 (Fourier Series)

Let $f_0(x) = x$ on the interval $0 < x < 2$, $f_0(x) = -x$ on $-2 < x < 0$, $f_0(x) = 0$ for $x = 0$, $f_0(x) = 2$ at $x = \pm 2$. Let $f(x)$ be the periodic extension of f_0 to the whole real line, of period 4.

(a) [80%] Compute the Fourier coefficients for the terms $\sin(67\pi x)$ and $\cos(2\pi x)$. Leave tedious integrations in integral form, but evaluate the easy ones like the integral of the square of sine or cosine.

(b) [20%] Which values of x in $|x| < 12$ might exhibit Gibb's phenomenon?

Answer:

(a) Because $f_0(x)$ is even, then $f(x)$ is even. Then the coefficient of $\sin(67\pi x)$ is zero, without computation, because all sine terms in the Fourier series of f have zero coefficient. The coefficient of $\cos(n\pi x/2)$ for $n > 0$ is given by the formula

$$a_n = \frac{1}{2} \int_{-2}^2 f_0(x) \cos(n\pi x/2) dx = \int_0^2 x \cos(n\pi x/2) dx.$$

For $\cos(2\pi x)$, we select $n\pi x/2 = 2\pi x$, or index $n = 4$.

(b) There are no jump discontinuities, f is continuous, so no Gibbs overshoot.

3 (Cosine and Sine Series)

Find the first nonzero term in the sine series expansion of $f(x)$, formed as the odd 2π -periodic extension of the function $\sin(x)\cos(x)$ on $0 < x < \pi$. Leave the Fourier coefficient in integral form, unevaluated, unless you can compute the value in a minute or two.

Answer:

Because $\sin(x)\cos(x) = (1/2)\sin(2x)$ is odd and 2π -periodic, this is the Fourier series of f . This term is for coefficient b_2 , so $b_2 = 1/2$ is the first nonzero Fourier coefficient. The first nonzero term is $(1/2)\sin(2x)$.

4 (Convergence of Fourier Series)

(a) [30%] Dirichlet's kernel formula can be used to evaluate the sum $\cos(2x) + \cos(4x) + \cos(6x) + \cos(8x)$. Report its value according to that formula.

(b) [40%] The Fourier Convergence Theorem for piecewise smooth functions applies to continuously differentiable functions of period $2p$. State the theorem for this special case, by translating the results when f is smooth and the interval $-\pi \leq x \leq \pi$ is replaced by $-p \leq x \leq p$.

(c) [30%] Give an example of a function $f(x)$ periodic of period 2 that has a Gibb's overshoot at the integers $x = 0, \pm 2, \pm 4, \dots$, (all $\pm 2n$) and nowhere else.

Answer:

(a) $\frac{1}{2} + \cos(x) + \dots + \cos(nx) = \frac{\sin(nx + x/2)}{2 \sin(x/2)}$ is used with x replaced by $2x$ and $n = 4$ to obtain the answer $0.5 \sin(8x + x)/\sin(x) - 0.5$.

(b) Let f be a p -periodic smooth function on $(-\infty, \infty)$. Then for all values of x ,

$$f(x) = a_0 + \sum_{n=1}^{\infty} (a_n \cos(n\pi x/p) + b_n \sin(n\pi x/p)),$$

where the Fourier coefficients a_0, a_n, b_n are given by the Euler formulas:

Problem 4
Fourier
Series

$$a_0 = \frac{1}{2p} \int_{-p}^p f(x) dx, \quad a_n = \frac{1}{p} \int_{-p}^p f(x) \cos(n\pi x/p) dx,$$

$$b_n = \frac{1}{p} \int_{-p}^p f(x) \sin(n\pi x/p) dx.$$

(c) Any 2-periodic continuous function f will work, if we alter the values of f at the desired points to produce a jump discontinuity. For example, define $f(x) = \sin(\pi x)$ except at the points $\pm 2n$, where $f(x) = 2$ ($f(2n) = 2$ for $n = 0, \pm 1, \pm 2, \pm 3, \dots$).

From
S2013
final Exam
3150

5. (Fourier Series)

(a) [30%] Find and display the nonzero terms in the Fourier series expansion of $f(x)$, formed as the even 2π -periodic extension of the function $f_0(x) = \sin^2(x) + 4 \cos(2x)$ on $0 < x < \pi$.

(b) [50%] Compute the Fourier sine series coefficients b_n for the function $g(x)$, defined as the period 2 odd extension of the function $g_0(x) = 1$ on $0 \leq x \leq 1$. Draw a representative graph for the partial Fourier sum for five terms of the infinite series.

(c) [20%] Define $h_0(x) = \begin{cases} \sin(2x) & 0 \leq x < \pi, \\ x - \pi & \pi \leq x \leq 2\pi, \end{cases}$ and let $f(x)$ be the 4π odd periodic extension of $h_0(x)$ to the whole real line. Compute the sum $f(-5.25\pi) + f(1.5\pi)$.

5(a) $f(x) = \sin^2(x) + 4 \cos(2x) = \frac{1 - \cos 2x}{2} + 4 \cos(2x) = \frac{1}{2} + \frac{7}{2} \cos(2x)$

It is even, so equals its periodic even extension.

Answer: $a_0 = \frac{1}{2}, a_2 = \frac{7}{2}$, all others are zero

5(b) Since g is odd, then all $a_n = 0$. Compute $b_n = \frac{1}{1} \int_{-1}^1 f(x) \sin\left(\frac{n\pi x}{1}\right) dx$

$b_n = \frac{4}{n\pi}$ for n odd, others zero.

5(c) $f(-5.25\pi) = f(4\pi - 5.25\pi)$


$= f(-1.25\pi)$

$= f(1.25\pi)$ odd f

$= 1.25\pi - \pi$ by $h_0(x)$ definition

$f(1.5\pi) = 1.5\pi - \pi$ "

Sum $= 2.75\pi - 2\pi = \pi/4$



wave
equation
problem 1

1. (Vibration of a Finite String)

The normal modes for the string equation $u_{tt} = c^2 u_{xx}$ are given by the functions

$$\sin\left(\frac{n\pi x}{L}\right) \cos\left(\frac{n\pi ct}{L}\right), \quad \sin\left(\frac{n\pi x}{L}\right) \sin\left(\frac{n\pi ct}{L}\right).$$

It is known that each normal mode is a solution of the string equation and that the problem below has solution $u(x, t)$ equal to an infinite series of constants times normal modes.

Solve the finite string vibration problem on $0 \leq x \leq 2, t > 0$,

$$\begin{aligned} u_{tt} &= c^2 u_{xx}, \\ u(0, t) &= 0, \\ u(2, t) &= 0, \\ u(x, 0) &= 0, \\ u_t(x, 0) &= -11 \sin(5\pi x). \end{aligned}$$

Answer:

Because the wave initial shape is zero, then the only normal modes are sine times sine. The initial wave velocity is already a Fourier series, using orthogonal set $\{\sin(n\pi x/2)\}_{n=1}^{\infty}$. The 1-term Fourier series $-11 \sin(5\pi x)$ can be modified into a solution by inserting the missing sine factor present in the corresponding normal mode. Then $u(x, t) = -11 \sin(5\pi x) \sin(5\pi ct)/(5\pi)$. We check it is a solution.

Blackboard photos Wed 13, 4 April

KEY

Partial Differential Equations 3150

Midterm Exam 2

Exam Date: Monday, 22 April 2013

Instructions: This exam is timed for 50 minutes. You will be given extra time to complete the exam. No calculators, notes, tables or books. Problems use only chapters 1 to 4 of the textbook. No answer check is expected. Details count 3/4, answers count 1/4.

2. (CH3. Finite String: Fourier Series Solution)

(a) [75%] Display the series formula without derivation details for the finite string problem

$$\begin{cases} u_{tt}(x,t) = c^2 u_{xx}(x,t), & 0 < x < L, \quad t > 0, \\ u(0,t) = 0, & t > 0, \\ u(L,t) = 0, & t > 0, \\ u(x,0) = f(x), & 0 < x < L, \\ u_t(x,0) = g(x), & 0 < x < L. \end{cases}$$

Symbols f and g should not appear explicitly in the series for $u(x,t)$. Expected in the formula for $u(x,t)$ are product solutions times constants.

(b) [25%] Display an explicit formula for the Fourier coefficients which contains the symbols L , $f(x)$, $g(x)$.

(a) Normal modes: $\sin(n\pi x/L) \cos(n\pi ct/L)$,
 $\sin(n\pi x/L) \sin(n\pi ct/L)$

ans 1 $\rightarrow u(x,t) =$ superposition of the normal modes

$$= \sum_{n=1}^{\infty} a_n \sin(n\pi x/L) \cos(n\pi ct/L) + \sum_{n=1}^{\infty} b_n \sin(n\pi x/L) \sin(n\pi ct/L)$$

(b) $f(x) = u(x,0) = \sum_{n=1}^{\infty} a_n \sin(n\pi x/L)$

ans 2 $\rightarrow a_n = \frac{2}{L} \int_0^L f(x) \sin(n\pi x/L) dx$ by \perp relations
for $\{\sin(n\pi x/L)\}_{n=1}^{\infty}$
 $g(x) = u_t(x,0) = \sum_{n=1}^{\infty} \frac{n\pi c}{L} b_n \sin(n\pi x/L)$

ans 3 $\rightarrow b_n = \frac{L}{n\pi c} \frac{2}{L} \int_0^L g(x) \sin(n\pi x/L) dx$ by \perp relations

Use this page to start your solution. Attach extra pages as needed, then staple.

3. (CH4. Rectangular Membrane)

Consider the general membrane problem

$$\begin{cases} u_{tt}(x, y, t) = c^2(u_{xx}(x, y, t) + u_{yy}(x, y, t)), & 0 < x < a, 0 < y < b, t > 0, \\ u(x, y, t) = 0 & \text{on the boundary,} \\ u(x, y, 0) = f(x, y), & 0 < x < a, 0 < y < b, \\ u_t(x, y, 0) = g(x, y), & 0 < x < a, 0 < y < b. \end{cases} \quad 100$$

Solve the problem for $a = b = c = 1$, $f(x, y) = 1$, $g(x, y) = 0$. Expected are displays for the normal modes, a superposition formula for $u(x, y, t)$, and explicit numerical values for the generalized Fourier coefficients.

The solution is a superposition of the normal modes obtained from separation of variables as:

$$\sin\left(\frac{m\pi x}{a}\right) \sin\left(\frac{n\pi y}{b}\right) \left[B_{mn} \cos(\lambda_{mn} t) + B_{mn}^* \sin(\lambda_{mn} t) \right]$$

With $\lambda_{mn} = c\pi \sqrt{\left(\frac{m}{a}\right)^2 + \left(\frac{n}{b}\right)^2}$

$$u(x, y, t) = \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \sin\left(\frac{m\pi x}{a}\right) \sin\left(\frac{n\pi y}{b}\right) \left[B_{mn} \cos(\lambda_{mn} t) + B_{mn}^* \sin(\lambda_{mn} t) \right]$$

where the Fourier coefficients are:

$$B_{mn} = 4 \int_0^b \int_0^a f(x, y) \sin\left(\frac{m\pi x}{a}\right) \sin\left(\frac{n\pi y}{b}\right) dx dy$$

$$\lambda_{mn} B_{mn}^* = 4 \int_0^b \int_0^a g(x, y) \sin\left(\frac{m\pi x}{a}\right) \sin\left(\frac{n\pi y}{b}\right) dx dy$$

turn
→
(answer continued
on next page)

Use this page to start your solution. Attach extra pages as needed, then staple.

Problem 3 continued

plugging in:

$$a=b=c=1 \quad f(x,y) = 1 \quad g(x,y) = 0$$

normal modes:

$$\sin(m\pi x) \sin(n\pi y) [B_{mn} \cos(\lambda_{mn} t) + B_{mn}^* \sin(\lambda_{mn} t)]$$

$$\lambda_{mn} = \pi \sqrt{m^2 + n^2}$$

$$U(x,y,t) = \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \sin(m\pi x) \sin(n\pi y) [B_{mn} \cos(\lambda_{mn} t) + B_{mn}^* \sin(\lambda_{mn} t)]$$

Where the Fourier coefficients are:

$$B_{mn} = 4 \int_0^1 \int_0^1 \sin(m\pi x) \sin(n\pi y) dx dy$$

$$= 4 \int_0^1 -\cos(m\pi x) \sin(n\pi y) \left(\frac{1}{m\pi}\right) \Big|_0^1 dy$$

$$= 4 \int_0^1 \left[-\cos(m\pi) \sin(n\pi y) \left(\frac{1}{m\pi}\right) - \sin(n\pi y) \left(\frac{1}{m\pi}\right) \right] dy$$

$$= 4 \left[\cos(m\pi) \cos(n\pi y) \left(\frac{1}{m\pi^2}\right) + \cos(n\pi y) \left(\frac{1}{m\pi^2}\right) \right] \Big|_0^1$$

$$B_{mn} = 4 \left[\cos(m\pi) \cos(n\pi) \left(\frac{1}{m\pi^2}\right) + \cos(n\pi) \left(\frac{1}{m\pi^2}\right) - \cos(m\pi) \left(\frac{1}{m\pi^2}\right) - \left(\frac{1}{m\pi^2}\right) \right]$$

$$B_{mn} = \frac{4}{mn\pi^2} \left[\cos(m\pi) \cos(n\pi) + \cos(n\pi) - \cos(m\pi) - 1 \right]$$

turn
→
(answer continued
on next page)

Problem 3 (continued)

$$\begin{aligned} B_{mn}^x &= 4 \int_0^1 \int_0^1 0 \sin\left(\frac{m\pi x}{a}\right) \sin\left(\frac{n\pi y}{b}\right) dx dy \\ &= 4 \int_0^1 \int_0^1 0 dx dy = 0 \end{aligned}$$

$$\boxed{B_{mn}^x = 0}$$

4. (Finite String: Fourier Series Solution)

(a) [50%] Display the series formula, complete with derivation details, for the solution $u(x, t)$ of the finite string problem

$$\begin{cases} u_{tt}(x, t) = \frac{1}{4}u_{xx}(x, t), & 0 < x < 2, & t > 0, \\ u(0, t) = 0, & & t > 0, \\ u(2, t) = 0, & & t > 0, \\ u(x, 0) = f(x), & 0 < x < 2, \\ u_t(x, 0) = g(x), & 0 < x < 2. \end{cases}$$

Symbols f and g should not appear explicitly in the series for $u(x, t)$. Expected in the formula for $u(x, t)$ are product solutions times constants.

(b) [25%] Display explicit formulas for the Fourier coefficients which contains the symbols $f(x)$, $g(x)$.

(c) [25%] Evaluate the Fourier coefficients when $f(x) = 100$ and $g(x) = 0$.

Duplicate of problem 2, but evaluate coefficients.

See 3150 final exam, 5/2013, <http://math.utah.edu/~ngustafso/s2013/>

Fourier Transform

1. (Fourier Transform Theory)

Blackboard
photo
4 Apr
Week 13

(a) [40%] Define Haberman's Fourier transform pair. Give an example of $f(x)$ and $F(w)$ which satisfy these equations.

(b) [60%] The heat equation on the line $-\infty < x < \infty$ can be solved by Fourier transform methods. Outline the method, called Fourier's Method, for the example

$$u_t = 4u_{xx}, \quad -\infty < x < \infty, \quad t > 0, \quad u(x, 0) = f(x).$$

2. (Fourier's Method)

photo
7 Apr
Week 14

Use the Heat kernel, the convolution theorem and the shift theorem to solve the diffusion-convection equation

$$u_t(x, t) = ku_{xx}(x, t) + cu_x(x, t), \quad t > 0, \quad -\infty < x < \infty, \quad u(x, 0) = f(x).$$

Answer:
$$u(x, t) = \frac{1}{\sqrt{4\pi kt}} \int_{-\infty}^{\infty} f(v) e^{-\frac{(x+ct-v)^2}{4kt}} dv$$

3. (Heat Equation and Gauss' Heat Kernel)

Solve the insulated rod heat conduction problem

photo
7 Apr
Week 14

$$\begin{cases} u_t(x, t) = \frac{1}{4}u_{xx}(x, t), & -\infty < x < \infty, \quad t > 0, \\ u(x, 0) = f(x), & -\infty < x < \infty, \\ f(x) = \begin{cases} 50 & 0 < x < 1, \\ 100 & -1 < x < 0 \\ 0 & \text{otherwise} \end{cases} \end{cases}$$

Hint: Use the heat kernel $g_t(x, v) = \frac{\sqrt{k}}{\sqrt{kt}} e^{-\frac{(x-v)^2}{4kt}}$, the error function $\text{erf}(x) = \frac{2}{\sqrt{\pi}} \int_0^x e^{-z^2} dz$,