

Solution Set Basis for Linear Differential Equations

- Nth Order Linear Differential Equation
- Atoms
- Examples of Atoms
- Theorems about Atoms
 - Atoms are independent
 - Euler's Theorem
 - Basis of the solution set
- How to use Euler's Theorem
- Examples

Linear Differential Equations

The solution set of a homogeneous constant coefficient linear differential equation

$$y^{(n)} + a_{n-1}y^{(n-1)} + \cdots + a_0y = 0$$

is known to be a vector space of functions of dimension n , consisting of special linear combinations

$$(1) \quad y = c_1f_1 + \cdots + c_nf_n,$$

where f_1, \dots, f_n are elementary functions known as **atoms**.

Definition of Atom

A **base atom** is defined to be one of

$$1, e^{ax}, \cos bx, \sin bx, e^{ax} \cos bx, e^{ax} \sin bx,$$

with real $a \neq 0$, $b > 0$.

An **atom** equals a base atom multiplied by x^n , where $n = 0, 1, 2 \dots$ is an integer.

An atom has coefficient 1, and the zero function is not an atom.

Examples of Atoms

$$1, x, x^2, e^x, xe^{-x}, x^{15}e^{2x} \cos 3x, \cos 3x, \sin 2x, x^2 \cos 2x, x^6 \sin \pi x, \\ x^{10}e^{\pi x} \sin 0.1x$$

Functions that are not Atoms

$$x/(1+x), \ln|x|, e^{x^2}, \sin(x+1), 0, 2x, \sin(1/x), \sqrt{x}$$

Theorems about Atoms

Theorem 1 (Independence)

Any finite list of atoms is linearly independent.

Theorem 2 (Euler)

The *real* characteristic polynomial $p(r) = r^n + a_{n-1}r^{n-1} + \dots + a_0$ has a factor $(r - a - ib)^{k+1}$ if and only if

$$x^k e^{ax} \cos bx, \quad x^k e^{ax} \sin bx$$

are real solutions of the differential equation (1). If $b > 0$, then both are atoms. If $b = 0$, then only the first is an atom.

Theorem 3 (Real Solutions)

If u and v are real and $u + iv$ is a solution of equation (1), then u and v are real solutions of equation (1).

Theorem 4 (Basis)

The solution set of equation (1) has a basis of n solution atoms which are determined by Euler's theorem.

Euler's Theorem Translated

Theorem 5 (How to Apply Euler's Theorem)

Factor dividing $p(r)$	Solution Atom(s)
$(r - 5)$	e^{5x}
$(r + 7)^2$	e^{-7x}, xe^{-7x}
$(r + 7)^3$	$e^{-7x}, xe^{-7x}, x^2e^{-7x}$
r	e^{0x}
r^2	e^{0x} and xe^{0x}
r^3	$1, x$ and x^2 [$e^{0x} = 1$]
$(r - 5i)$	$\cos 5x$ and $\sin 5x$
$(r + 3i)^2$	$\cos 3x, x \cos 3x, \sin 3x, x \sin 3x$
$(r - 2 + 3i)^2$	$e^{2x} \cos 3x, xe^{2x} \cos 3x, e^{2x} \sin 3x, xe^{2x} \sin 3x$

Example 1. Solve $y''' = 0$. _____

Solution: $p(r) = r^3$ implies 1 is a base atom and then $1, x, x^2$ are solution atoms. They are independent, hence form a basis for the 3-dimensional solution space. Then $y = c_1 + c_2x + c_3x^2$.

Example 2. Solve $y'' + 4y = 0$. _____

Solution: $p(r) = r^2 + 4$ implies base atoms $\cos 2x$ and $\sin 2x$. They are a basis for the 2-dimensional solution space with $y = c_1 \cos 2x + c_2 \sin 2x$.

Example 3. Solve $y'' + 2y' = 0$. _____

Solution: $p(r) = r^2 + 2r$ implies $1, e^{-2x}$ are base solution atoms. These independent atoms form a basis. Then $y = c_1 + c_2e^{-2x}$.

Example 4. Solve $y^{(4)} + 4y'' = 0$. _____

Solution: $p(r) = r^4 + 4r^2 = r^2(r^2 + 4)$ implies the four atoms $1, x, \cos 2x, \sin 2x$ are solutions. Then $y = c_1 + c_2x + c_3 \cos 2x + c_4 \sin 2x$.

Example 5. Solve the differential equation if $p(r) = (r^3 - r^2)(r^2 - 1)(r^2 + 4)^2$.

Solution: The distinct factors of $p(r)$ are $r^2, (r - 1)^2, r + 1, (r - 2i)^2, (r + 2i)^2$. Euler's theorem implies the DE has nine solution atoms $1, x, e^x, xe^x, e^{-x}, \cos 2x, x \cos 2x, \sin 2x, x \sin 2x$. Then y is a linear combination of these atoms.