

Sample Quiz S2014 Math 2250

problem 1

Answers: (1)  $u = 68 + 117 e^{-ht}$

(2)  $h = (-1/50) \ln\left(\frac{113}{117}\right) \cong 0.0006957$

(3)  $t \cong 345.52$  seconds, about 6 min.

Details(1). Because  $u$  = degrees F and  $t$  = seconds, then the model is  $\begin{cases} u' = -h(u - 68), \\ u(0) = 185, \quad u(50) = 181. \end{cases}$

The DE is solved by superposition  $u = u_h + u_p$ . The equilibrium solution is  $u_p = 68$ . Then  $u_h = \frac{c}{\text{integ. factor}} = c e^{-ht}$ , using standard linear form  $u' + hu = 68h$ . Condition  $u(0) = 185$  is used on  $u = u_h + u_p = ce^{-ht} + 68$  to evaluate  $185 = ce^0 + 68$ , then  $c = 185 - 68 = 117$ .

Details(2). Start with answer (1) and use  $u(50) = 181$ .

Then

$$181 = 68 + 117 e^{-ht} \quad \text{when } t=50$$

$$113 = 117 e^{-ht}$$

$$e^{-ht} = \frac{113}{117} \Rightarrow -ht = \ln\left(\frac{113}{117}\right)$$

$$\Rightarrow h = \frac{-1}{50} \ln\left(\frac{113}{117}\right)$$

Details(3). As in details(2),

$$160 = 68 + 117 e^{-ht}$$

$$e^{-ht} = \frac{160 - 68}{117}$$

$$-ht = \ln\left(\frac{92}{117}\right) \quad \text{take log across eqtn.}$$

$$t = \left(\frac{-1}{h}\right) \ln\left(\frac{92}{117}\right)$$

$$t = 50 \frac{\ln(92/117)}{\ln(113/117)} \cong 345.52 \text{ seconds}$$

Sample Quiz 2      Spring 2014      Math 2250

problem 2

Answers (1)  $x' = 0.8x(1 - \frac{x}{780500})$ ,  $x=0, 780500$

$$(2) x^2 - Kx + \frac{HK}{r} = 0, \text{ roots} = \frac{K}{2} \pm \frac{1}{2} \sqrt{K^2 - \frac{4HK}{r}}$$




$$a = \frac{K}{2} + \frac{1}{2}\sqrt{D}$$

$$b = \frac{K}{2} - \frac{1}{2}\sqrt{D}$$

$$D = K^2 - \frac{4HK}{r}$$

$$\text{Choose } H = \frac{Kr}{4} - 5000$$



choose  $H = \frac{kr}{4} + 5000$   
All sols decrease  
to zero (and beyond),  
meaning extinction

Details (1).  $x' = rx(1 - \frac{x}{K})$ , substitute  $r=0.8$ ,  $K=780500$

Details (2).  $x' = rx(1 - \frac{x}{K}) - H = rx - \frac{r}{K}x^2 - H$

$$x' = -\frac{r}{K}(x^2 - Kx + \frac{HK}{r})$$

Apply quadratic formula to  $x^2 - Kx + \frac{HK}{r} = 0$  to find roots, reported in both (2), (3) above.

Details (3). A double real root is when the discriminant  $D = K^2 - \frac{4HK}{r} = 0$ , requiring  $H = \frac{Kr}{4}$ . For  $H < \frac{Kr}{4}$ , there are 2 real roots  $a, b$  as given in the answer. For  $H > \frac{Kr}{4}$  there are no real roots, therefore  $x' < 0$  and  $x$  decreases to zero (extinction) and beyond.

The larger root  $a = \frac{K}{2} + \frac{1}{2}\sqrt{D}$  is a stable funnel.

The smaller root  $b = \frac{K}{2} - \frac{1}{2}\sqrt{D}$  is an unstable spout.

When  $D=0$ , then  $a=b=\frac{K}{2}$  is a node. When  $a, b$  are real, extinction for  $x < b$ , carrying capacity =  $a$ , sustainable harvest for  $x > b$ .

```

> F:=x->r*x*(1-x/K):G:=x->r*x*(1-x/K)-H:
> solve(G(x)=0,x);

$$\frac{1}{2} \frac{Kr + \sqrt{K^2 r^2 - 4 H K r}}{r}, \frac{1}{2} \frac{Kr - \sqrt{K^2 r^2 - 4 H K r}}{r} \quad (1)$$


```


```

> de:=diff(x(t),t)=G(x(t)):r:=0.8:K:=780500:H0:=K*r/4:H:=H0-5000:
a:=K/2+(1/2)*sqrt(K^2-4*H*K/r);b:=K/2-(1/2)*sqrt(K^2-4*H*K/r);
ic:=[[0,0.9*b],[0,0.94*b],[0,0.98*b],[0,b],[0,(a+b)/2],[0,(3*a+b)/4],
[0,a],[0,1.2*a],[0,1.4*a],[0,1.6*a]]:
opts:=dirfield=[10,10],arrows=line,color=gray,linecolor=black,
thickness=2:
> DEtools[DEmplot](de,x(t),t=0..36,x=0..K,ic,opts);

```

$$a := 4.600935752 \cdot 10^5$$

$$b := 3.204064248 \cdot 10^5$$



```

> de:=diff(x(t),t)=G(x(t)):
r:=0.8:K:=780500:H0:=K*r/4:
H:=H0+6000:a:=K/2;b:=K/2;
ic:=[[0,0.9*b],[0,0.94*b],[0,0.98*b],[0,b],[0,a],[0,1.2*a],[0,1.4*a],
[0,1.6*a],[0,1.8*a],[0,2*a]]:

```

```

opts:=dirfield=[10,10],arrows=line,color=gray,linecolor=black,
thickness=2:
DEtools[DEplot](de,x(t),t=0..36,x=0..K,ic,opts);
H := 1.621000000 105
a := 390250
b := 390250

```

