

Systems of Second Order Differential Equations

Cayley-Hamilton-Ziebur

- Characteristic Equation
- Cayley-Hamilton
 - Cayley-Hamilton Theorem
 - An Example
- Euler's Substitution for $\vec{u}'' = A\vec{u}$
- The Cayley-Hamilton-Ziebur Method for $\vec{u}'' = A\vec{u}$

Characteristic Equation

Definition 1 (Characteristic Equation)

Given a square matrix A , the **characteristic equation** of A is the polynomial equation

$$\det(A - \lambda I) = 0.$$

The determinant $\det(A - \lambda I)$ is formed by subtracting λ from the diagonal of A . The polynomial $p(x) = \det(A - xI)$ is called the **characteristic polynomial** of matrix A .

- If A is 2×2 , then $p(x)$ is a quadratic.
- If A is 3×3 , then $p(x)$ is a cubic.
- The determinant is expanded by the cofactor rule, in order to preserve factorizations.

Characteristic Equation Examples

Create $\det(A - xI)$ by subtracting x from the diagonal of A .

Evaluate by the cofactor rule.

$$A = \begin{pmatrix} 2 & 3 \\ 0 & 4 \end{pmatrix}, \quad p(x) = \begin{vmatrix} 2-x & 3 \\ 0 & 4-x \end{vmatrix} = (2-x)(4-x)$$

$$A = \begin{pmatrix} 2 & 3 & 4 \\ 0 & 5 & 6 \\ 0 & 0 & 7 \end{pmatrix}, \quad p(x) = \begin{vmatrix} 2-x & 3 & 4 \\ 0 & 5-x & 6 \\ 0 & 0 & 7-x \end{vmatrix} = (2-x)(5-x)(7-x)$$

Cayley-Hamilton

Theorem 1 (Cayley-Hamilton)

A square matrix A satisfies its own characteristic equation.

If $p(x) = (-x)^n + a_{n-1}(-x)^{n-1} + \dots + a_0$, then the result is the equation

$$(-A)^n + a_{n-1}(-A)^{n-1} + \dots + a_1(-A) + a_0I = 0,$$

where I is the $n \times n$ identity matrix and 0 is the $n \times n$ zero matrix.

The 2×2 Case

Then $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ and for $a_1 = \text{trace}(A)$, $a_0 = \det(A)$ we have $p(x) = x^2 + a_1(-x) + a_0$. The Cayley-Hamilton theorem says

$$A^2 + a_1(-A) + a_0 \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}.$$

Cayley-Hamilton Example

Assume

$$A = \begin{pmatrix} 2 & 3 & 4 \\ 0 & 5 & 6 \\ 0 & 0 & 7 \end{pmatrix}$$

Then

$$p(x) = \begin{vmatrix} 2-x & 3 & 4 \\ 0 & 5-x & 6 \\ 0 & 0 & 7-x \end{vmatrix} = (2-x)(5-x)(7-x)$$

and the Cayley-Hamilton Theorem says that

$$(2I - A)(5I - A)(7I - A) = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}.$$

Euler's Substitution and the Characteristic Equation

Definition. Euler's Substitution for the second order equation $\vec{u}'' = A\vec{u}$ is

$$\vec{u} = \vec{v}e^{rt}.$$

The symbol r is a real or complex constant and symbol \vec{v} is a constant vector.

Theorem 2 (Euler Solution Equation from Euler's Substitution)

Euler's substitution applied to $\vec{u}'' = A\vec{u}$ leads directly to the equation

$$|A - r^2I| = 0.$$

This is perhaps the premier method for remembering the characteristic equation for the second order vector-matrix equation $\vec{u}'' = A\vec{u}$.

Proof: Substitute $\vec{u} = \vec{v}e^{rt}$ into $\vec{u}'' = A\vec{u}$ to obtain $r^2e^{rt}\vec{v} = A\vec{v}e^{rt}$. Cancel the exponential, then $r^2\vec{v} = A\vec{v}$. Re-arrange to the homogeneous system $(A - r^2I)\vec{v} = \vec{0}$. This homogeneous linear algebraic equation has a nonzero solution \vec{v} if and only if the determinant of coefficients vanishes: $|A - r^2I| = 0$.

Cayley-Hamilton-Ziebur Method for Second Order Systems _____

Theorem 3 (Cayley-Hamilton-Ziebur Structure Theorem for $\vec{u}'' = A\vec{u}$)

The solution $\vec{u}(t)$ of second order equation $\vec{u}''(t) = A\vec{u}(t)$ is a vector linear combination of Euler solution atoms corresponding to roots of the equation $\det(A - r^2 I) = 0$.

The equation $|A - r^2 I| = 0$ is formed by substitution of $\lambda = r^2$ into the eigenanalysis characteristic equation of A .

In symbols, the structure theorem says

$$\vec{u} = \vec{d}_1 A_1 + \cdots + \vec{d}_k A_k,$$

where A_1, \dots, A_k are Euler solution atoms corresponding to the roots r of the determining equation $|A - r^2 I| = 0$. Therefore, all vectors in the relation have real entries. However, only $2n$ entries of vectors $\vec{d}_1, \dots, \vec{d}_k$ are arbitrary constants, the remaining entries being dependent on them.

Proof of the Cayley-Hamilton-Ziebur Theorem

Consider the case when \mathbf{A} is 2×2 ($n = 2$), because the proof details are similar in higher dimensions. Expand $|\mathbf{A} - x\mathbf{I}| = 0$ to find the characteristic equation $x^2 + cx + d = 0$, for some constants c, d . The Cayley-Hamilton theorem says that $\mathbf{A}^2 + c\mathbf{A} + d\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}$. Let \vec{u} be a solution of $\vec{u}''(t) = \mathbf{A}\vec{u}(t)$. Multiply the Cayley-Hamilton identity by vector \vec{u} and simplify to obtain

$$\mathbf{A}^2\vec{u} + c\mathbf{A}\vec{u} + d\vec{u} = \vec{0}.$$

Using equation $\vec{u}''(t) = \mathbf{A}\vec{u}(t)$ backwards, we compute $\mathbf{A}^2\vec{u} = \mathbf{A}\vec{u}'' = \vec{u}''''$. Replace the terms of the displayed equation to obtain the relation

$$\vec{u}'''' + c\vec{u}'' + d\vec{u} = \vec{0}.$$

Each component y of vector \vec{u} then satisfies the 4th order linear homogeneous equation $y^{(4)} + cy^{(2)} + dy = 0$, which has characteristic equation $r^4 + cr^2 + d = 0$. This equation is the expansion of determinant equation $|\mathbf{A} - r^2\mathbf{I}| = 0$. Therefore y is a linear combination of the Euler solution atoms found from the roots of this equation. It follows then that \vec{u} is a vector linear combination of the Euler solution atoms so identified. This completes the proof.

A 2×2 Illustration

Solve the system $\vec{u}'' = A\vec{u}$, $A = \begin{pmatrix} -75 & 25 \\ 50 & -50 \end{pmatrix}$, which is a spring-mass system with $k_1 = 100$, $k_2 = 50$, $m_1 = 2$, $m_2 = 1$.

Solution: The eigenvalues of A are $\lambda = -25$ and -100 . Then the determining equation $|A - r^2 I| = 0$ has complex roots $r = \pm 5i$ and $\pm 10i$ with corresponding Euler solution atoms $\cos(4t)$, $\sin(5t)$, $\cos(10t)$, $\sin(10t)$. The eigenpairs of A are

$$\left(-25, \begin{pmatrix} 1 \\ 2 \end{pmatrix}\right), \quad \left(-100, \begin{pmatrix} 1 \\ -1 \end{pmatrix}\right).$$

Then \vec{u} is a vector linear combination of the Euler solution atoms

$$\vec{u}(t) = \vec{d}_1 \cos(5t) + \vec{d}_2 \sin(5t) + \vec{d}_3 \cos(10t) + \vec{d}_4 \sin(10t).$$

A 2 × 2 Illustration continued

How to Find \vec{d}_1 to \vec{d}_4

Substitute the formula

$$\vec{u}(t) = \vec{d}_1 \cos(5t) + \vec{d}_2 \sin(5t) + \vec{d}_3 \cos(10t) + \vec{d}_4 \sin(10t)$$

into $\vec{u}'' = A\vec{u}$, then solve for the unknown vectors \vec{d}_j , $j = 1, 2, 3, 4$, by equating coefficients of Euler solution atoms matching left and right:

$$A\vec{d}_1 = -25\vec{d}_1, \quad A\vec{d}_2 = -25\vec{d}_2, \quad A\vec{d}_3 = -100\vec{d}_3, \quad A\vec{d}_4 = -100\vec{d}_4.$$

These eigenpair relationships imply formulas involving the eigenvectors of A . We get, for some constants a_1, a_2, b_1, b_2 ,

$$\vec{d}_1 = a_1 \begin{pmatrix} 1 \\ 2 \end{pmatrix}, \quad \vec{d}_2 = b_1 \begin{pmatrix} 1 \\ 2 \end{pmatrix}, \quad \vec{d}_3 = a_2 \begin{pmatrix} 1 \\ -1 \end{pmatrix}, \quad \vec{d}_4 = b_2 \begin{pmatrix} 1 \\ -1 \end{pmatrix}.$$

Summary for the 2×2 Illustration

$$\vec{u}(t) = \vec{d}_1 \cos(5t) + \vec{d}_2 \sin(5t) + \vec{d}_3 \cos(10t) + \vec{d}_4 \sin(10t)$$

$$\vec{u}(t) = (a_1 \cos(5t) + b_1 \sin(5t)) \begin{pmatrix} 1 \\ 2 \end{pmatrix} + (a_2 \cos(10t) + b_2 \sin(10t)) \begin{pmatrix} 1 \\ -1 \end{pmatrix}$$