

Name _____

Differential Equations 2280

Midterm Exam 1

Exam Date: Friday, 26 February 2016 at 12:50pm

Instructions: This in-class exam is designed for 50 minutes. No calculators, notes, tables or books. No answer check is expected. Details count 3/4, answers count 1/4.

1. (Quadrature Equations)

(a) [40%] Solve $y' = \frac{2x^3}{1+x^2}$.

(b) [60%] Find the position $x(t)$ from the velocity model $\frac{d}{dt}(e^{-t}v(t)) = 2e^t$, $v(0) = 5$ and the position model $\frac{dx}{dt} = v(t)$, $x(2) = 2$.

Solution to Problem 1.

(a) Answer $y = x^2 - \ln(x^2 + 1) + c$. The integral of $F(x) = \frac{2x^3}{1+x^2}$ is found by substitution $u = 1+x^2$, resulting in the new integration problem $\int F dx = \int \frac{u-1}{u} du = \int (1) du - \int \frac{du}{u}$.

(b) Velocity $v(t) = 2e^{2t} + 3e^t$ by quadrature. Integrate $x'(t) = 2e^{2t} + 3e^t$ with $x(0) = 2$ to obtain position $x(t) = e^{2t} + 3e^t + c$, where $c = 2 - e^4 - 3e^2$.

Name. _____

2. (Classification of Equations)

The differential equation $y' = f(x, y)$ is defined to be **separable** provided $f(x, y) = F(x)G(y)$ for some functions F and G .

(a) [40%] The equation $y' + x(y + 3) = ye^x + 3x$ is separable. Provide formulas for $F(x)$ and $G(y)$.

(b) [60%] Apply partial derivative tests to show that $y' = x + y$ is linear but not separable. Supply all details.

Solution to Problem 2.

(a) The equation is $y' = ye^x - xy = (e^x - x)y$. Then $F(x) = e^x - x$, $G(y) = y$.

(b) Let $f(x, y) = x + y$. Then $\partial f / \partial y = 1$, which is independent of y , hence the equation $y' = f(x, y)$ is linear. The negative test is $\frac{\partial f / \partial y}{f}$ depends on x . In this case, the fraction is

$$\frac{\partial f / \partial y}{f} = \frac{1}{f} = \frac{1}{x + y}.$$

At $y = 0$, this reduces to $1/x$, which depends on x , therefore the equation $y' = f(x, y)$ is not separable.

Name. _____

3. (Solve a Separable Equation)Given $(5y + 10)y' = (xe^{-x} + \sin(x) \cos(x)) (y^2 + 3y - 4)$.

Find a non-equilibrium solution in implicit form.

To save time, **do not solve** for y explicitly and **do not solve** for equilibrium solutions.**Solution to Problem 3.**The solution by separation of variables identifies the separated equation $y' = F(x)G(y)$ using

$$F(x) = xe^{-x} + \sin(x) \cos(x), \quad G(y) = \frac{y^2 + 3y - 4}{5y + 10}.$$

The integral of F is done by parts and also by u -substitution.

$$\begin{aligned} \int F dx &= \int xe^{-x} dx + \int \sin(x) \cos(x) dx \\ &= I_1 + I_2. \\ I_1 &= \int xe^{-x} dx \\ &= -xe^{-x} - \int e^{-x} dx, \quad \text{parts } u = x, dv = e^{-x} dx, \\ &= xe^{-x} - e^{-x} + c_1 \\ I_2 &= \int \sin(x) \cos(x) dx \\ &= \int u du, \quad u = \sin(x), du = \cos(x) dx, \\ &= u^2/2 + c_2 \\ &= \frac{1}{2} \sin^2(x) + c_2 \end{aligned}$$

Then $\int F dx = xe^{-x} - e^{-x} + \frac{1}{2} \sin^2(x) + c_3$.The integral of $1/G(y)$ requires partial fractions. The details:

$$\begin{aligned} \int \frac{dx}{G(y(x))} &= \int \frac{5u + 10}{u^2 + 3u - 4} du, \quad u = y(x), du = y'(x) dx, \\ &= \int \frac{5u + 10}{(u + 4)(u - 1)} du \\ &= \int \frac{A}{u + 4} + \frac{B}{u - 1} du, \quad A, B \text{ determined later,} \\ &= A \ln |u + 4| + B \ln |u - 1| + c_4 \end{aligned}$$

The partial fraction problem

$$\frac{5u + 10}{(u + 4)(u - 1)} = \frac{A}{u + 4} + \frac{B}{u - 1}$$

can be solved in a variety of ways, with answer $A = \frac{-20+10}{-5} = 2$ and $B = \frac{15}{5} = 3$. The final implicit solution is obtained from $\int \frac{dx}{G(y(x))} = \int F(x) dx$, which gives the equation

$$2 \ln |y + 4| + 3 \ln |y - 1| = xe^{-x} - e^{-x} + \frac{1}{2} \sin^2(x) + c.$$

Name. _____

4. (Linear Equations)

(a) [60%] Solve the linear model $2x'(t) = -64 + \frac{10}{3t+2}x(t)$, $x(0) = 32$. Show all integrating factor steps.

(b) [20%] Solve $\frac{dy}{dx} - (\cos(x))y = 0$ using the homogeneous linear equation shortcut.

(c) [20%] Solve $5\frac{dy}{dx} - 7y = 10$ using the superposition principle $y = y_h + y_p$ shortcut. Expected are answers for y_h and y_p .

Solution to Problem 4.

(a) The answer is $v(t) = 32 + 48t$. The details:

$$v'(t) = -32 + \frac{5}{3t+2}v(t),$$

$$v'(t) + \frac{-5}{3t+2}v(t) = -32, \quad \text{standard form } v' + p(t)v = q(t)$$

$$p(t) = \frac{-5}{3t+2},$$

$$W = e^{\int p dt}, \quad \text{integrating factor}$$

$$W = e^u, \quad u = \int p dt = -\frac{5}{3} \ln |3t+2| = \ln(|3t+2|^{-5/3})$$

$$W = (3t+2)^{-5/3}, \quad \text{Final choice for } W.$$

Then replace the left side of $v' + pv = q$ by $(vW)'/W$ to obtain

$$v'(t) + \frac{-5}{3t+2}v(t) = -32, \quad \text{standard form } v' + p(t)v = q(t)$$

$$\frac{(vW)'}{W} = -32, \quad \text{Replace left side by quotient } (vW)'/W$$

$$(vW)' = -32W, \quad \text{cross-multiply}$$

$$vW = -32 \int W dt, \quad \text{quadrature step.}$$

The evaluation of the integral is from the power rule:

$$\int -32W dt = -32 \int (3t+2)^{-5/3} dt = -32 \frac{(3t+2)^{-2/3}}{(-2/3)(3)} + c.$$

Division by $W = (3t+2)^{-5/3}$ then gives the general solution

$$v(t) = \frac{c}{W} - \frac{32}{-2}(3t+2)^{-2/3}(3t+2)^{5/3}.$$

Constant c evaluates to $c = 0$ because of initial condition $v(0) = 32$. Then

$$v(t) = \frac{32}{-2}(3t+2)^{-2/3}(3t+2)^{5/3} = 16(3t+2)^{-\frac{2}{3}+\frac{5}{3}} = 16(3t+2).$$


(b) The answer is $y = \text{constant}$ divided by the integrating factor: $y = \frac{c}{W}$. Because $W = e^u$ where $u = \int -\cos(x)dx = -\sin x$, then $y = ce^{\sin x}$.

(c) The equilibrium solution (a constant solution) is $y_p = -\frac{10}{7}$. The homogeneous solution is $y_h = ce^{7x/5} = \text{constant}$ divided by the integrating factor. Then $y = y_p + y_h = -\frac{10}{7} + ce^{7x/5}$.

Name. _____

5. (Stability)

Assume an autonomous equation $x'(t) = f(x(t))$. Draw a phase portrait with at least 12 threaded curves, using the phase line diagram given below. Add these labels as appropriate: funnel, spout, node [neither spout nor funnel], stable, unstable.

**Solution to Problem 5.**

The graphic is drawn using increasing and decreasing curves, which may or may not be depicted with turning points. The rules:

1. A curve drawn between equilibria is increasing if the sign is PLUS.
2. A curve drawn between equilibria is decreasing if the sign is MINUS.
3. Label: FUNNEL, STABLE
The signs left to right are PLUS MINUS crossing the equilibrium point.
4. Label: SPOUT, UNSTABLE
The signs left to right are MINUS PLUS crossing the equilibrium point.
5. Label: NODE, UNSTABLE
The signs left to right are PLUS PLUS crossing the equilibrium point, or
The signs left to right are MINUS MINUS crossing the equilibrium point.

The answer:

- $x = -10$: FUNNEL, STABLE
 $x = -5$: SPOUT, UNSTABLE
 $x = -3$: FUNNEL, STABLE
 $x = 0$: SPOUT, UNSTABLE
 $x = 3$: NODE, UNSTABLE

Name. _____

6. (ch3)

Using Euler's theorem on Euler solution atoms and the characteristic equation for higher order constant-coefficient differential equations, solve (a), (b), (c).

(a) [40%] Find a constant coefficient differential equation $ay'' + by' + cy = 0$ which has particular solutions $-5e^{-x} + xe^{-x}$, $10e^{-x} + xe^{-x}$.

(b) [30%] Given characteristic equation $r(r-2)(r^3+4r)(r^2+2r+37) = 0$, solve the differential equation.

(c) [30%] Given $mx''(t) + cx'(t) + kx(t) = 0$, which represents an unforced damped spring-mass system. Assume $m = 4$, $c = 4$, $k = 129$. Classify the equation as over-damped, critically damped or under-damped. Illustrate in a spring-mass-dashpot drawing the assignment of physical constants m , c , k and the initial conditions $x(0) = 1$, $x'(0) = 0$.

Solution to Problem 6.**6(a)**

Multiply the first solution by 2 and add it to the second solution. Then Euler atom xe^{-x} is a solution, which implies that $r = -1$ is a double root of the characteristic equation. Then the characteristic equation should be $(r - (-1))(r - (-1)) = 0$, or $r^2 + 2r + 1 = 0$. The differential equation is $y'' + 2y' + y = 0$.

6(b)

The characteristic equation factors into $r^2(r-2)(r^2+4)((r+1)^2+36) = 0$ with roots $r = 0, 0, 2; \pm 2i; -1 \pm 6i$. Then y is a linear combination of the Euler solution atoms $1, x, e^{2x}, \cos(2x), \sin(2x); e^{-x} \cos(6x), e^{-x}$.

6(c)

Use $4r^2 + 4r + 129 = 0$ and the quadratic formula to obtain roots $r = -1/2 + 4\sqrt{2}i, -1/2 - 4\sqrt{2}i$ and Euler solution atoms $e^{-x/2} \cos 4\sqrt{2}t, e^{-x/2} \sin 4\sqrt{2}t$. Then y is a linear combination of these two solution atoms, and it oscillates, therefore the classification is **under-damped**. The illustration shows a spring, a dashpot and a mass with labels k, c, m . Initial conditions mean mass elongation $x = 1$, at rest.

Name. _____

7. (ch3)

Determine for $y^{(3)} + y^{(2)} = x + 2e^{-x} + \sin x$ the corrected trial solution for y_p according to the method of undetermined coefficients. **Do not evaluate the undetermined coefficients!** The trial solution should be the one with fewest Euler solution atoms.

Solution to Problem 7.

The homogeneous equation $y^{(3)} + y^{(2)} = 0$ has solution $y_h = c_1 + c_2x + c_3e^{-x}$, because the characteristic polynomial has roots 1.

1 Rule I constructs an initial trial solution y from the list of independent Euler solution atoms

$$e^{-x}, \quad 1, \quad x, \quad \cos x, \quad \sin x.$$

Linear combinations of these atoms are supposed to reproduce, by assignment of constants, all derivatives of $F(x) = x + 2e^{-x} + \sin x$, which is the right side of the differential equation. Each of y_1 to y_4 in the display below is constructed to have the same **base atom**, which is the Euler atom obtained by stripping the power of x . For example, $x = xe^{0x}$ strips to base atom e^{0x} or 1.

$$\begin{aligned} y &= y_1 + y_2 + y_3 + y_4, \\ y_1 &= d_1 e^{-x}, \\ y_2 &= d_2 + d_3 x, \\ y_3 &= d_4 \cos x, \\ y_4 &= d_5 \sin x. \end{aligned}$$

2 Rule II is applied individually to each of y_1, y_2, y_3, y_4 to give the **corrected trial solution**

$$\begin{aligned} y &= y_1 + y_2 + y_3 + y_4, \\ y_1 &= d_1 x e^{-x}, \\ y_2 &= x^2(d_2 + d_3 x), \\ y_3 &= d_4 \cos x, \\ y_4 &= d_5 \sin x. \end{aligned}$$

The powers of x multiplied in each case are selected to eliminate terms in the initial trial solution which duplicate homogeneous equation Euler solution atoms. The factor used is exactly x^s of the Edwards-Penney table, where s is the multiplicity of the characteristic equation root r that produced the related atom in the homogeneous solution y_h . The atoms in y_3, y_4 are not solutions of the homogeneous equation, therefore y_3, y_4 are unaltered.