

Recall: $f(x, y)$ differentiable at (x_0, y_0) means it has a good tangent approximation there:

$$f(x_0+h_1, y_0+h_2) = f(x_0, y_0) + h_1 f_x(x_0, y_0) + h_2 f_y(x_0, y_0) + h_1 \varepsilon_1(\vec{h}) + h_2 \varepsilon_2(\vec{h}), \quad \varepsilon_1, \varepsilon_2 \rightarrow 0 \text{ as } \vec{h} \rightarrow \vec{0}$$

i.e.

$$f(x, y) = \underbrace{f(x_0, y_0) + f_x(x_0, y_0)(x-x_0) + f_y(x_0, y_0)(y-y_0)}_{T(x, y)} + \vec{h} \cdot \vec{\varepsilon}(\vec{h}) \quad \vec{\varepsilon}(\vec{h}) \rightarrow \vec{0} \text{ as } \vec{h} \rightarrow \vec{0}$$

and if we write this in vector notation, we see that the definition makes sense for a fun of 3, 4, or n variables:

$$f(\vec{p}_0 + \vec{h}) = f(\vec{p}_0) + \nabla f(\vec{p}_0) \cdot \vec{h} + \vec{h} \cdot \vec{\varepsilon}(\vec{h})$$

$$\vec{\varepsilon}(\vec{h}) \rightarrow \vec{0} \text{ as } \vec{h} \rightarrow \vec{0}.$$

$\nabla f(\vec{p}_0)$ = the vector of partial derivs of f , at $\vec{p}_0$
called gradient of f at $\vec{p}_0$.

Theorem

- If the partial derivs of f are continuous near $\vec{p}_0$, then f is differentiable at $\vec{p}_0$.

- Theorem: f differentiable at $\vec{p}_0$ implies f continuous there. Converse not true

Reason

$$f(\vec{p}_0 + \vec{h}) - f(\vec{p}_0) = \nabla f(\vec{p}_0) \cdot \vec{h} + \vec{h} \cdot \vec{\varepsilon}(\vec{h})$$

$$\Rightarrow |f(\vec{p}_0 + \vec{h}) - f(\vec{p}_0)| \leq | \quad |$$

$$\leq | \quad | = \|\nabla f(\vec{p}_0)\| \|\vec{h}\| |\cos \theta_1| + \|\vec{h}\| \|\vec{\varepsilon}(\vec{h})\| |\cos \theta_2|$$

$$\leq \|\nabla f(\vec{p}_0)\| \|\vec{h}\| + \|\vec{h}\| \|\vec{\varepsilon}(\vec{h})\|$$

$$\rightarrow 0 \text{ as } \vec{h} \rightarrow \vec{0}.$$

Directional derivatives : Recall

$$f_x(x_0, y_0) = \lim_{h \rightarrow 0} \frac{f(x_0 + h, y_0) - f(x_0, y_0)}{h} = \text{rate of change of } f \text{ in } \hat{i} \text{ direction}$$

$\uparrow$
scalar

$$f_y(x_0, y_0) = \lim_{h \rightarrow 0} \frac{f(x_0, y_0 + h) - f(x_0, y_0)}{h} = \text{rate of change of } f \text{ in } \hat{j} \text{ direction.}$$

Now,
Let $\vec{u}$ be a unit vector in any direction, f a function of several variables, differentiable at $\vec{p}_0$.

$$D_{\vec{u}} f(\vec{p}_0) := \lim_{h \rightarrow 0} \frac{f(\vec{p}_0 + h\vec{u}) - f(\vec{p}_0)}{h}$$

$\uparrow$
scalar

is called the directional derivative of f in direction $\vec{u}$, at $\vec{p}_0$ and generalizes the concept of partial derivative, to any direction

Examples : this includes partial derivatives as a special case :

$$D_{\hat{i}} f(x_0, y_0) =$$
$$D_{\hat{j}} f(x_0, y_0, z_0) =$$
$$D_{\langle 0, 0, 0, 1 \rangle} f(x_0, y_0, z_0, w_0) =$$

Theorem : Directional derivatives are easy to compute, if f is differentiable at $\vec{p}_0$.

proof : Use the tangent approximation formula

$$f(\vec{p}_0 + h\vec{u}) = f(\vec{p}_0) + \nabla f(\vec{p}_0) \cdot h\vec{u} + h\vec{u} \cdot \vec{\epsilon}(h\vec{u})$$

$$\frac{f(\vec{p}_0 + h\vec{u}) - f(\vec{p}_0)}{h} = \frac{\nabla f(\vec{p}_0) \cdot h\vec{u}}{h} + \frac{h\vec{u} \cdot \vec{\epsilon}(h\vec{u})}{h}$$

take $\lim_{h \rightarrow 0}$:

$$D_{\vec{u}} f(\vec{p}_0) = \nabla f(\vec{p}_0) \cdot \vec{u}$$

Because $\vec{u} \cdot \vec{\epsilon}(h\vec{u}) \leq \|\vec{u}\| \|\vec{\epsilon}(h\vec{u})\| \leq \|\vec{\epsilon}(h\vec{u})\|$

$\frac{\|\vec{\epsilon}(h\vec{u})\|}{\|h\vec{u}\|} \rightarrow 0$ as $h \rightarrow 0$

More fun with index cards


3

$$f(x,y) = e^{2x}(1 + \sin 3y)$$


$$(x_0, y_0) = (0, 0)$$

Far view

level curves of $f(x,y)$, at increments of 1 unit




graph $z = f(x,y)$




Medium view

level curves of $f(x,y)$, at increments of 0.1 unit



contours of graph $z = f(x,y)$





Near view

4

$$f(x,y) = e^x + e^{2x} \sin 3y$$

near view

level curves of $f(x,y)$, increments of 0.01 unit



Estimate, and find exact values, for $D_{\vec{u}} f(\vec{0})$: Step 0: Find $\nabla f(0,0)$.

A) $\vec{u} = \langle 1, 0 \rangle$ $D_{\vec{u}} f(\vec{0}) \approx \frac{1.04 - 1}{.02} = 2$

$$f_x(\vec{0}) = \nabla f(\vec{0}) \cdot \langle 1, 0 \rangle = 2!$$

B) $\vec{u} = \langle 0, 1 \rangle$ $D_{\vec{u}} f(\vec{0}) \approx \frac{1.09 - 1}{.03} = 3$

$$f_y(\vec{0}) = 3!$$

C) $\vec{u} = \frac{1}{\sqrt{13}} \langle 2, 3 \rangle$

D) $\vec{u} = \frac{1}{\sqrt{13}} \langle -3, 2 \rangle$

E) $\vec{u} = \frac{1}{\sqrt{5}} \langle 1, -2 \rangle$

F) $\vec{u} = \frac{1}{\sqrt{5}} \langle 2, 1 \rangle$

G) $\vec{u} = -\frac{1}{\sqrt{13}} \langle 2, 3 \rangle$

Conclusions

$$D_{\vec{u}} f(\vec{p}_0) = \nabla f(\vec{p}_0) \cdot \vec{u} = |\nabla f(\vec{p}_0)| |\vec{u}| \cos \theta = |\nabla f(\vec{p}_0)| \cos \theta$$

So


$$-|\nabla f(\vec{p}_0)| \leq D_{\vec{u}} f(\vec{p}_0) \leq |\nabla f(\vec{p}_0)|$$

↑
if $\vec{u} = -\frac{\nabla f(\vec{p}_0)}{|\nabla f(\vec{p}_0)|}$

get this min value

↑
if $\vec{u} = \frac{\nabla f(\vec{p}_0)}{|\nabla f(\vec{p}_0)|}$ get this max value

gradient points
in direction of
maximum increase




$D_{\vec{u}} f(\vec{p}_0) = 0$ if $\vec{u}$ is in direction of level curve
thru $\vec{p}_0$

i.e. $\cos \theta = 0$

i.e. $\nabla f(\vec{p}_0) \perp$ level curve

(level surface in higher dims!)

gradient and level curves for
 $f(x,y)=x^2-y^2$



My directional derivative calculations: (for page 4)

> Digits:=3;		Digits:=3
> .04/.02;	#<1,0> approx	
2;	#exact	2.00
		2
> .06/.02;	#<0,1> approx	
3;	#exact	3.00
		3
> .133/.036;	#<2,3> approx	
	sqrt(13.0);#exact	3.69
		3.61
> .072/.022;	#<2,1> approx;	
	7/sqrt(5.0);#exact	3.27
		3.12
> 0/3.6;	#<-3,2> approx	
0;	#exact	0.
		0
> -.01/.022;	#<-2,1> approx	
	-1/sqrt(5.0);#exact	-4.55
		-4.46
> (.875-1)/.0355;	#<-2,-3> approx	
	-sqrt(13.0);#exact	-3.52
		-3.61
> (.958-1)/.022	#<1,-2> approx	-1.9
	-4/sqrt(5)	exact -1.7