

Name _____

Student I.D. _____

Math 2250-4
Exam #1
October 4, 2013

Please show all work for full credit. This exam is closed book and closed note. You may use a scientific calculator, but not a "graphing calculator" i.e. not one which is capable of integration, taking derivatives, or matrix algebra. **In order to receive full or partial credit on any problem, you must show all of your work and justify your conclusions.** There are 100 points possible. The point values for each problem are indicated in the right-hand margin. **Good Luck!**

	Score	POSSIBLE
1	_____	10
2	_____	25
3	_____	20
4	_____	15
5	_____	20
6	_____	10
TOTAL	_____	100

1) Concepts:

1a) What is a linear differential equation?

(2 points)

1b) What is an autonomous differential equation?

(2 points)

1c) What is an equilibrium solution to an autonomous differential equation?

(2 points)

1d) For a square matrix $A_{n \times n}$ what condition on the reduced row echelon form of A is equivalent to A having an inverse matrix A^{-1} ?

(2 points)

1e) For a general rectangular matrix $A_{m \times n}$ what is the condition on the reduced row echelon form of A that guarantees that each matrix equation $A \mathbf{x} = \mathbf{b}$ will be consistent (i.e. have at least one solutions \mathbf{x}), for every possible choice of the vector \mathbf{b} ?

(2 points)

2) Consider a boat which starts at rest at time $t = 0$ sec, is accelerated in a straight path by an engine that provides a constant $1500 N$ of force, and which is also subject to drag forces of $60 N$ for each $\frac{m}{s}$ of velocity. The boat (containing the pilot) has mass $300 kg$.

2a) Use your modeling ability to show that the boat velocity $v(t)$ (in meters per second) satisfies the initial value problem

$$\begin{aligned}v'(t) &= 5 - .2 v \\v(0) &= 0\end{aligned}$$

(5 points)

2b) Solve the initial value problem in 2a. Use the algorithm for separable differential equations.

(15 points)

2c) How far does the boat travel in the first ten seconds?

(5 points)

3) Consider the following input-output model: A brine tank initially contains 1600 gallons of brine - with an initial concentration is 0.1 lb of salt for each gallon of water. At time $t = 0$ water begins to flow into and out of the tank at a rate of 80 gallons per minute, keeping the water volume constant. The salt concentration of the incoming water varies with time, with concentration $2 e^{-0.05 t}$ pounds per gallon, and the brine solution in the tank remains well mixed so that the concentration of brine leaving the tank is always the average concentration in the tank. (There is space below for you to sketch the configuration, as it may help you with the rest of the problem.)

3a) Use the description above to show the salt amount $x(t)$ pounds at time t minutes solves the differential equation

$$x'(t) = 160 e^{-0.05 t} - .05 x(t) .$$

(6 points)

3b) Solve the initial value problem for the salt amount $x(t)$ in this problem.

(14 points)

4) Consider the differential equation for $x(t)$:

$$x'(t) = -x^3 + 6x^2 - 9x.$$

4a) Find the equilibrium solutions.

(4 points)

4b) Construct the phase diagram for this differential equation, and use it to determine the stability of the equilibrium solutions.

(6 points)

4c) Let $x(t)$ be the solution to the initial value problem for this differential equation, with $x(0) = 4$.

Without solving for $x(t)$ deduce the value of $\lim_{t \rightarrow \infty} x(t)$. Explain.

(5 points)

5) Consider the homogeneous linear system

$$\begin{bmatrix} 2 & 3 & 0 \\ 1 & 2 & -1 \\ 2 & 0 & 6 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}.$$

5a) Compute the determinant of the matrix in this system. (Be careful to check your work!)

(4 points)

5b) What does the value of the determinant in 5a tell you about the possibilities, for the number of solutions to this homogeneous system? Explain.

(4 points)

5c) Find the solution set to the linear system above, by computing the reduced row echelon form of the augmented matrix and backsolving.

(12 points)

6a) Find the inverse matrix to

$$A = \begin{bmatrix} -2 & 3 \\ -3 & 5 \end{bmatrix}.$$

(5 points)

You may use either method we learned.

6b) Use your work from 6a to solve the matrix equation below for the unknown matrix X :

$$X \begin{bmatrix} -2 & 3 \\ -3 & 5 \end{bmatrix} = \begin{bmatrix} 4 & 3 \\ 2 & 1 \end{bmatrix}.$$

(5 points)