

5.5 Finding y_p for non-homogeneous linear differential equations

$$L(y) = f$$

(so that you can use the general solution $y = y_p + y_H$ to solve initial value problems).

There are two methods we will use:

- The method of undetermined coefficients uses guessing algorithms, and works for constant coefficient linear differential equations and certain classes of functions $f(x)$. The method seems magic, but actually relies on vector space theory. This is the main focus of section 5.5.
- The method of variation of parameters is more general, and yields an integral formula for a particular solution y_p , assuming you are already in possession of a basis for the homogeneous solution space. It works for any linear differential equation and any (continuous) function f , but the formulas can get computationally messy especially for differential equations of order $n > 2$. We'll focus on the case $n = 2$.

The easiest way to explain the method of undetermined coefficients is with examples.

Exercise 1) Find a particular solution $y_p(x)$ for the differential equation

$$L(y) := y'' + 4y' - 5y = 10x + 3.$$

Hint: try $y_p(x) = Ax + B$ (because $L(Ax + B) = cx + d$).

Exercise 2) Use your work in 1 and your expertise with homogeneous linear differential equations to find the general solution to

$$y'' + 4y' - 5y = 10x + 3$$

Exercise 3) Find a particular solution to

$$L(y) = y'' + 4y' - 5y = 14e^{2x}.$$

Hint: try $y_p = C e^{2x}$ since $L(C e^{2x}) = d e^{2x}$.

Exercise 4a) Use superposition and your work from the previous exercises to find the general solution to

$$L(y) = y'' + 4y' - 5y = 14e^{2x} - 20x - 6.$$

4b) Solve (or at least set up the problem to solve) the initial value problem

$$y'' + 4y' - 5y = 14e^{2x} - 20x - 6$$

$$y(0) = 4$$

$$y'(0) = -4.$$

4c) Check your answer with technology.

Exercise 5) Find a particular solution to

$$L(y) := y''' + 4y' - 5y = 2 \cos(3x) .$$

Hint: To solve $L(y) = f$ we hope that f is in some finite dimensional subspace V that is preserved by L , i.e. $L : V \rightarrow V$.

- In Exercise 1 $V = \text{span}\{1, x\}$ and so we guessed $y_p = A + Bx$.
- In Exercise 3 $V = \text{span}\{e^{2x}\}$ and so we guessed $y_p = C e^{2x}$.
- What's the smallest subspace V we can take in the current exercise? Can you see why $V = \text{span}\{\cos(3x)\}$ and a guess of $y_p = D \cos(3x)$ won't work?

Method of undetermined coefficients (base case): If f is in a finite dimensional subspace V with the property that L preserves V , and if the only function $g \in V$ for which $L(g) = 0$ is $g = 0$, then there is a unique $y_p \in V$ with $L(y_p) = f$.

proof: This is a special case of an important vector space theorem called the rank+nullity theorem that you will explore in your homework. Among other results, this theorem implies that if V and W are vector spaces with the same finite dimension, and if $L : V \rightarrow W$ is a linear transformation between them, then the equation $L(\underline{v}) = \underline{w}$ has unique solutions $\underline{v} \in V$ for each $\underline{w} \in W$ if and only if the only solution $\underline{v} \in V$ to $L(\underline{v}) = \underline{0}$ is $\underline{v} = \underline{0}$. We already know this fact is true for matrix transformations $L(\underline{x}) = A_{n \times n} \underline{x}$ with $L :$

$\mathbb{R}^n \rightarrow \mathbb{R}^n$ (because each requirement is equivalent to saying that the reduced row echelon form of A is the identity), but the result is really more general than that special case.

Exercise 6) Use the method of undetermined coefficients to guess the form for a particular solution $y_p(x)$ for a constant coefficient differential equation (assuming the only such solution which would solve the homogeneous DE is the zero solution):

6a) $L(y) = x^3 + 6x - 5$

6b) $L(y) = 4e^{2x}\sin(3x)$

6c) $L(y) = x\cos(2x)$

BUT LOOK OUT

Exercise 7a) Find a particular solution to

$$y'' + 4y' - 5y = 4e^x.$$

Hint: since $y_H = c_1e^x + c_2e^{-5x}$, a guess of $y_p = Ae^x$ will not work (and $\text{span}\{e^x\}$ does not satisfy the vector space conditions for undetermined coefficients). Instead try

$$y_p = Ax e^x$$

and factor $L = D^2 + 4D - 5I = (D + 5I) \circ (D - I)$.

7b) check work with technology

The vector space theorem (special case of rank+nullity theorem) alluded to above for $L : V \rightarrow W$, combined with our understanding of how to factor constant coefficient differential operators leads to the more general method of undetermined coefficients algorithm (and superposition), for right hand sides which can be written as sums of functions having the indicated forms. See the discussion in pages 341-346 of the text, and the table on page 346, reproduced here. If L has a factor $(D - rI)^s$ then the guesses you would have made need to be multiplied by x^s , as in Exercise 7.

See discussion p 341-346. Also p. 346 Table: Don't forget to also use superposition for $f_1 + f_2$

$f(x)$	y_p	^{potential} related root of $p(r)$
$P_m(x) = b_0 + b_1x + \dots + b_mx^m$	$x^s (A_0 + A_1x + A_2x^2 + \dots + A_mx^m)$	$r=0$
$a \cos kx + b \sin kx$	$x^s (A \cos kx + B \sin kx)$	$r = \pm ik$
$e^{rx} (a \cos kx + b \sin kx)$	$x^s e^{rx} (A \cos kx + B \sin kx)$	root = $r + \pm ik$
$P_m(x) e^{rx}$	$x^s (A_0 + A_1x + \dots + A_mx^m) e^{rx}$	root = r
$P_m(x) (a \cos kx + b \sin kx)$	$x^s [(A_0 + A_1x + \dots + A_mx^m) \cos kx + (B_0 + B_1x + \dots + B_mx^m) \sin kx]$	root = $\pm ik$

$x^s = 1$ ($s=0$) as long as the potential root is not actually a root of the characteristic polynomial for L (easy case, Exercises 1-4)

Otherwise, s is the power of $(r - \text{root})$ in factored $p(r)$

e.g. in Exercise 5,

$$p(r) = (r+5)^1 (r-1)^1 \quad \swarrow \text{so multiply guess by } x^1.$$