Exercise 2) There are other possibilities. In the two systems below we kept all of the coefficients the same as in Exercise 1, except for a_{33} , and we changed the right side in the third equation, for 2<u>a</u>. Work out what happens in each case.

<u>2a)</u>

$$x + 2y + z = 4$$

3 x + 8 y + 7 z = 20
2 x + 7 y + 8 z = 20.

<u>2b)</u>

$$x + 2y + z = 4$$

3 x + 8 y + 7 z = 20
2 x + 7 y + 8 z = 23

2x + 7y + 8z = 23. <u>2c</u>) What are the possible solution sets (and geometric configurations) for 1, 2, 3, 4,... equations in 3 unknowns?

Summary of the systematic method known as Gaussian elimination for solving systems of linear equations.

We write the linear system (LS) of *m* equations for the vector $\mathbf{x} = [x_1, x_2, \dots, x_n]$ of the *n* unknowns as

$$a_{11} x_1 + a_{12} x_2 + \dots + a_{1n} x_n = b_1$$

$$a_{21} x_1 + a_{22} x_2 + \dots + a_{2n} x_n = b_2$$

$$\vdots$$

$$a_{m1} x_1 + a_{m2} x_2 + \dots + a_{mn} x_n = b_m$$

The matrix that we get by adjoining (augmenting) the right-side <u>**b**</u>-vector to the coefficient matrix $A = \begin{bmatrix} a_{ij} \end{bmatrix}$ is called the *augmented matrix* $\langle A | \underline{b} \rangle$:

Our goal is to find all the solution vectors \underline{x} to the system - i.e. the <u>solution set</u>.

There are three types of *elementary equation operations* that don't change the solution set to the linear system. They are

- interchange two of equations
- multiply one of the equations by a non-zero constant
- replace an equation with its sum with a multiple of a different equation.

And that when working with the augmented matrix $\langle A | \underline{b} \rangle$ these correspond to the three types of *elementary row operations:*

- interchange ("swap") two rows
- multiply one of the rows by a non-zero constant
- replace a row by its sum with a multiple of a different row.

<u>Gaussian elimination</u>: Use elementary row operations and work column by column (from left to right) and row by row (from top to bottom) to first get the augmented matrix for an equivalent system of equations which is in

row-echelon form:

- (1) All "zero" rows (having all entries = 0) lie beneath the non-zero rows.
- (2) The leading (first) non-zero entry in each non-zero row lies strictly to the right of the one above
- it. These entries are called *pivots* in our text, and the corresponding columns are called *pivot columns*. (At this stage you could "backsolve" to find all solutions.)

Next, continue but by working from bottom to top and from right to left instead, so that you end with an augmented matrix that is in

reduced row echelon form: (1),(2), together with

(3) Each leading non-zero row entry has value 1. Such entries are called "leading 1's" and are the

"pivots" for the corresponding pivot columns.

(4) Each pivot column has 0's in all the entries except for the pivot entry of 1.

Finally, read off how to explicitly specify the solution set, by "backsolving" from the reduced row echelon form.

<u>Note:</u> There are lots of row-echelon forms for a matrix, but only one reduced row-echelon form. (We'll see why later.) All mathematical software packages have a command to find the reduced row echelon form of a matrix.

Exercise 3 Find all solutions to the system of 3 linear equations in 5 unknowns

$$\begin{aligned} x_1 &- 2 x_2 + 3 x_3 + 2 x_4 + x_5 &= 10 \\ 2 x_1 &- 4 x_2 + 8 x_3 + 3 x_4 + 10 x_5 &= 7 \\ 3 x_1 &- 6 x_2 + 10 x_3 + 6 x_4 + 5 x_5 &= 27 . \end{aligned}$$

Here's the augmented matrix:

1	-2	3	2	1	10
2	-4	8	3	10	7
3	-6	10	6	5	27

Find the reduced row echelon form of this augmented matrix and then backsolve to explicitly parameterize the solution set. (Hint: it's a two-dimensional plane in \mathbb{R}^5 , if that helps. :-))

Maple says:

with(LinearAlgebra): # matrix and linear algebra library > A := Matrix(3, 5, [1, -2, 3, 2, 1,2, -4, 8, 3, 10, 3, -6, 10, 6, 5]): b := Vector([10, 7, 27]): $\langle A | b \rangle;$ *# the mathematical augmented matrix doesn't actually have* # a vertical line between the end of A and the start of b *ReducedRowEchelonForm*($\langle A|b \rangle$); $\begin{bmatrix} 1 & -2 & 3 & 2 & 1 & 10 \\ 2 & -4 & 8 & 3 & 10 & 7 \\ 3 & -6 & 10 & 6 & 5 & 27 \end{bmatrix}$ $\begin{bmatrix} 1 & -2 & 0 & 0 & 3 & 5 \\ 0 & 0 & 1 & 0 & 2 & -3 \\ 0 & 0 & 0 & 1 & -4 & 7 \end{bmatrix}$ > LinearSolve(A, b);# this command will actually write down the general solution, using # Maple's way of writing free parameters, which actually makes # some sense. Generally when there are free parameters involved, # there will be equivalent ways to express the solution that may # look different. But usually Maple's version will look like yours, # because it's using the same algorithm and choosing the free parameters # the same way too. $5 + 2 _t_{2} - 3 _t_{5}$ $-t_{2}$ $-3 - 2 _t_{5}$ $7 + 4 _t_{5}$ t

(1)

(2)

Math 2270-004

Wed Jan 10

- 1.3 Vectors and vector equations
- Quiz today at end of class, on section 1.1-1.2 material

Announcements:

Warm-up Exercise:

until 12:58