

Math 2280-1

Friday 2/17

§3.6 forced oscillations

most usual periodic forcing function

$$m x'' + c x' + k x = F_0 \cos \omega t$$

Case 1 $c=0, \omega \neq \omega_0$ ($\omega_0 = \sqrt{\frac{k}{m}}$ still).

$$m x'' + k x = F_0 \cos \omega t$$

+ k [try $x_p = A \cos \omega t + B \sin \omega t$ since $c=0$!

+ 0 [$x_p' = -A \omega \sin \omega t$

+ m [$x_p'' = -A \omega^2 \cos \omega t$

$$L(x_p) = \cos \omega t [A] [k - m \omega^2] \stackrel{\text{want}}{=} F_0 \cos \omega t$$

$$A = \frac{F_0}{k - m \omega^2}$$

$$x_p(t) = \frac{F_0}{k - m \omega^2} \cos \omega t$$

Now solve

$$\begin{cases} m x'' + c x' + k x = F_0 \cos \omega t \\ x(0) = x_0 \\ x'(0) = v_0 \end{cases}$$

$$x(t) = x_p + x_H = \frac{F_0}{k - m \omega^2} \cos \omega t + A \cos \omega_0 t + B \sin \omega_0 t$$

$$x(0) = x_0 = \frac{F_0}{k - m \omega^2} + A \Rightarrow A = x_0 - \frac{F_0}{k - m \omega^2}$$

$$x'(0) = v_0 = B \omega_0 \Rightarrow B = \frac{v_0}{\omega_0}$$

hence

$$x(t) = \frac{F_0}{k - m \omega^2} [\cos \omega t - \cos \omega_0 t] + x_0 \cos \omega_0 t + \frac{v_0}{\omega_0} \sin \omega_0 t$$

or
$$x(t) = \frac{F_0}{m} \frac{1}{\omega_0^2 - \omega^2} [\cos \omega t - \cos \omega_0 t] + x_0 \cos \omega_0 t + \frac{v_0}{\omega_0} \sin \omega_0 t$$

HW for Fri 2/24


3.6 (4) (5) 7 (13) (16) (21) (22)

4.1 1 (8, 11, 15, 16, 21, 24, 26)

1

Beating

```
> plot(cos(31/10*t)-cos(3*t),
      t=0..62.8, color=black, title='half a beating period?');
```



Case 2 $c=0, \omega=\omega_0$ ("resonance")

$$m x'' + k x = F_0 \cos \omega_0 t$$

Find A and B so that

$$+ k [\quad x_p(t) = (A \cos \omega_0 t + B \sin \omega_0 t) t$$

$$o [\quad x_p'(t) =$$

$$+ m [\quad x_p''(t) =$$

$$L x_p''(t) =$$

ans $x_p(t) = \frac{F_0}{2m\omega_0} t \sin \omega_0 t$

VP soltn: $x(t) = \frac{F_0}{2m\omega_0} t \sin \omega_0 t + x_0 \cos \omega_0 t + \frac{y_0}{\omega_0} \sin \omega_0 t$

For $0 \leq t \leq M$ this formula can also be obtained from page 2 formula by letting $\omega \rightarrow \omega_0!$

Example :

$$F_0 = 5$$

$$m = 1$$

$$\omega_0 = 3$$


~~etc~~

$$\omega = 2 : \frac{F_0}{m} \frac{1}{\omega_0^2 - \omega^2} [\cos \omega t - \cos \omega_0 t]$$


$$= \frac{50}{5} (\cos 2t - \cos 3t)$$

(page 1 formula)

```
> plot(10*(cos(2*t)-cos(3*t)), t=0..10, color=black);
```




```
> plot(-100/(9-2.9^2)*sin(-.05*t)*sin(2.95*t), t=0..60,
numpoints=200, color=black);
```



$$\omega = 2.9 \quad x(t) = \frac{50(-2)}{(9-(2.9)^2)} \sin(-.05t) \sin(2.95t)$$

(page 2 formula)

```
> plot(25/3*t*sin(3*t), t=0..60, numpoints=200, color=black);
```



$$\omega = 3 \quad (\text{page 3 formula}) \quad x(t) = \frac{25}{3} t \sin 3t$$

Tues: Case 3 $c \neq 0!$