

Math 2280-1

Tuesday April 6

Finish Monday notes ~ classify linear & non-linear equilibria.

Then begin § 6.3: Population models

(today's notes will probably last thru Wednesday.)

6.3

• predator-prey:

$$\text{PP} \quad \begin{cases} x'(t) = ax - pxy = x(a - py) \\ y'(t) = -by + qxy = y(-b + qx) \end{cases}$$

prey
predator
 $a, b, p, q \geq 0$

natural region of interest: $x \geq 0, y \geq 0$

equil solns:

$$\begin{array}{ll} x=0 & y = a/p \\ \Downarrow & \Downarrow \\ y=0 & x = b/q \\ \begin{bmatrix} 0 \\ 0 \end{bmatrix} & \begin{bmatrix} b/q \\ a/p \end{bmatrix} \end{array}$$

in 1st quad.
is this equil stable?

linearize:

$$\begin{bmatrix} F \\ G \end{bmatrix} = \begin{bmatrix} ax - pxy \\ -by + qxy \end{bmatrix}$$

$$\text{Jacobian matrix} = \begin{bmatrix} F_x & F_y \\ G_x & G_y \end{bmatrix} = \begin{bmatrix} a - py & -px \\ qy & -b + qx \end{bmatrix}$$

$$\text{at } \vec{x}_*, \quad J = \begin{bmatrix} 0 & -pb/q \\ aq/p & 0 \end{bmatrix}$$

$$|J - \lambda I| = \lambda^2 + ab; \quad \lambda = \pm i\sqrt{ab}$$

linearization DE is stable center (ellipse trajectories)

this is borderline for the original non-linear system....


method for understanding soln trajectories: (in this case for the non-linear problem)

1210

Chain rule $\frac{dy}{dt} = \frac{dy}{dx} \frac{dx}{dt}$

So $\frac{dy}{dx} = \frac{\frac{dy}{dt}}{\frac{dx}{dt}}$

(use this in 6.9.1 #24 HW too!)



for PP, $\frac{dy}{dx} = \frac{y(-b+qx)}{x(a-py)}$

separable!

$\frac{a-py}{y} dy = \frac{-b+qx}{x} dx$


$a \ln y - py = -b \ln x + qx + C$ (in 1st quadrant!)

$\underbrace{a \ln y + b \ln x - py - qx}_{f(x,y)} = C$

So solution trajectories follow level curves of $f(x,y)$ (See figures 6.3.1 & 6.3.2 page 401)

and this $f(x,y)$ has a local max at $\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} b/q \\ a/p \end{bmatrix} = \bar{x}^*$ (Actually global 1st quadrant max!)

So $\bar{x}^*$ is stable, and general solns are periodic!!



• Competition model

$$\frac{dx}{dt} = \underbrace{a_1 x - b_1 x^2}_{\text{logistic}} - \underbrace{c_1 xy}_{\text{competition}}$$

all params > 0

$$\frac{dy}{dt} = \underbrace{a_2 y - b_2 y^2}_{\text{logistic}} - \underbrace{c_2 xy}_{\text{competition?}}$$

Equil. solns:

$$\begin{aligned} x [a_1 - b_1 x - c_1 y] &= 0 \\ y [a_2 - b_2 y - c_2 x] &= 0 \end{aligned}$$

$x = 0$

$$\Downarrow$$

$$y(a_2 - b_2 y) = 0$$

$$\Downarrow \quad \text{or} \quad \Downarrow$$

$$y = 0 \quad \text{or} \quad y = a_2/b_2$$

$$\begin{bmatrix} 0 \\ 0 \end{bmatrix} \quad \text{or} \quad \begin{bmatrix} 0 \\ a_2/b_2 \end{bmatrix}$$

$x \neq 0$

$$\Downarrow$$

$$a_1 - b_1 x - c_1 y = 0$$

$$\Downarrow \quad \text{or} \quad \Downarrow$$

$$y = 0 \quad \text{or} \quad a_2 - b_2 y - c_2 x = 0$$

$$\Downarrow \quad \Downarrow$$

$$x = a_1/b_1 \quad \text{or} \quad \begin{bmatrix} a_1/b_1 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} b_1 & c_1 \\ c_2 & b_2 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} a_1 \\ a_2 \end{bmatrix} \quad \text{(book type page 540 (8))}$$

$$\begin{bmatrix} x \\ y \end{bmatrix} = \frac{1}{b_1 b_2 - c_1 c_2} \begin{bmatrix} b_2 & -c_1 \\ -c_2 & b_1 \end{bmatrix} \begin{bmatrix} a_1 \\ a_2 \end{bmatrix}$$

$$= \frac{1}{b_1 b_2 - c_1 c_2} \begin{bmatrix} a_1 b_2 - c_1 b_1 \\ -a_1 c_2 + b_1 a_2 \end{bmatrix}$$

call this $\begin{bmatrix} x_E \\ y_E \end{bmatrix}$

Text claims

① If $c_1 c_2 < b_1 b_2$

(relatively low competition between species vs. self-inhibition)

then $\begin{bmatrix} x_E \\ y_E \end{bmatrix}$ is asymptotically stable equil, and $\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} \rightarrow \begin{bmatrix} x_E \\ y_E \end{bmatrix}$

② if $c_1 c_2 > b_1 b_2$, $\begin{bmatrix} x_E \\ y_E \end{bmatrix}$ is unstable, and

either $x(t)$ or $y(t) \rightarrow 0$ as $t \rightarrow \infty$

(with 100% probability)

[this may or may not be in 1st quadrant; depends on parameter values]

← (for any IVP if $x_0 > 0$, $y_0 > 0$, and if $\begin{bmatrix} x_E \\ y_E \end{bmatrix}$ is in 1st quad.)

Discussion of $\begin{bmatrix} x_E \\ y_E \end{bmatrix}$:

(4)

$$J = \begin{bmatrix} F_x & F_y \\ G_x & G_y \end{bmatrix} = \begin{bmatrix} a_1 - 2b_1x - c_1y & -c_1x \\ -c_2y & a_2 - 2b_2y - c_2x \end{bmatrix}$$

at $\begin{bmatrix} x_E \\ y_E \end{bmatrix}$ we have $\begin{bmatrix} b_1 & c_1 \\ c_2 & b_2 \end{bmatrix} \begin{bmatrix} x_E \\ y_E \end{bmatrix} = \begin{bmatrix} a_1 \\ a_2 \end{bmatrix}$

so (trickery!) at this point

$$J = \begin{bmatrix} -b_1x_E & -c_1x_E \\ -c_2y_E & -b_2y_E \end{bmatrix}$$

write $x = x_E$
 $y = y_E$

$$p(\lambda) = |J - \lambda I| = (b_1x + \lambda)(b_2y + \lambda) - c_1c_2xy$$


$$p(\lambda) = \lambda^2 + (b_1x + b_2y)\lambda + (b_1b_2 - c_1c_2)xy$$

$\underbrace{> 0}_{\text{by hypothesis}}$

the graph of $p(\lambda)$ is a concave up parabola

(2) if $c_1c_2 > b_1b_2$ then $p(0) < 0$

so $p(\lambda)$ has a positive and a negative root, by intermediate value thm, so $\begin{bmatrix} x_E \\ y_E \end{bmatrix}$ is (unstable) saddle.




(1) if $c_1c_2 < b_1b_2$, then

vertex of parabola occurs at $\lambda = -\left(\frac{b_1x + b_2y}{2}\right)$ (complete square!)

$$\text{and } p\left(-\frac{b_1x + b_2y}{2}\right) = \frac{1}{4}(b_1x + b_2y)^2 - \frac{1}{2}(b_1x + b_2y)^2 + (b_1b_2 - c_1c_2)xy$$

$$= -\frac{1}{4} \left[\underbrace{b_1^2x^2 + b_2^2y^2 + 2b_1b_2xy}_{(b_1x - b_2y)^2} - 4b_1b_2xy + 4c_1c_2 \right]$$



< 0



since $p(0) > 0$ in this case,
 $p(\lambda)$ has two negative roots.



(this proves the stability claims for the linearized problem, not the global claim for the nonlinear problem)



We didn't consider all possible cases, even of this "simple" competition model:

