

Math 2280-1
Wednesday Feb. 10

postpone 3.4 hw until
next week
(3.1-3.3 due this Friday)

①

- Carefully go through the rest of Monday's notes, which explain how to find a basis for $\ker L$

when
$$L(y) = y^{(n)} + a_{n-1}y^{(n-1)} + \dots + a_1y' + a_0$$
 (a_j constant)
(a_j real)

and the characteristic polynomial

$$p(r) := r^n + a_{n-1}r^{n-1} + \dots + a_1r + a_0$$

has real roots only (either n distinct roots, or repeated roots.)

- Then figure out what to do when the characteristic poly has complex roots:

Theorem 8 If r is a complex root of $p(r)$, $r = a + bi$ (a, b real)
 $i = \sqrt{-1}$

then $y_1 = e^{ax} \cos bx$

$$y_2 = e^{ax} \sin bx$$

are linearly independent soltns to $L(y) = 0$.

proof: you could check this directly, but there's a more natural (and motivated) proof using complex exponentials.

Step 1 Euler's formula:

$$e^{i\theta} := \cos \theta + i \sin \theta$$

(θ real)

motivation: check Taylor series!

not-needed here but good to know Step 1': $e^{i(\alpha+\beta)} = e^{i\alpha} e^{i\beta}$ is true! ($\cos, \sin$ addition angle!)

Step 2 $e^{\alpha+i\beta} := e^\alpha e^{i\beta} = e^\alpha (\cos \beta + i \sin \beta)$

(α, β real)

so $e^{(a+ib)x} = e^{ax+ibx} = e^{ax} (\cos bx + i \sin bx)$

(x real, a, b real)

Step 3 $D_x (f(x) + ig(x)) := f'(x) + ig'(x)$

f, g real fncs

Step 4 $D_x (e^{(a+ib)x}) = (a+ib) e^{(a+ib)x}$

i.e. if r is a complex number $D_x (e^{rx}) = r e^{rx}$.

This needs to be checked!!

proof of Theorem 8 Let $r = a+bi$ a root of the characteristic polynomial p

$$\text{let } y = e^{rx} = e^{ax} (\cos bx) + i e^{ax} \sin bx \\ = y_1 + i y_2$$

by step 4,

$$L(y) = L(e^{rx}) = p(r) e^{rx} = 0.$$

by step 3,
(and linearity)

$$L(y) = L(y_1 + i y_2) = L(y_1) + i L(y_2)$$

$0+0i$

$$\text{so } 0 = L(y_1) + i L(y_2)$$

"
real
function,
since a, b 's
are real

"
imag. fun
since a, b 's
are real.

$$\text{Thus } L(y_1) = 0 \quad \begin{matrix} \text{equate} \\ \text{(real parts)} \end{matrix} \\ L(y_2) = 0 \quad \begin{matrix} \text{equate} \\ \text{(imag. parts)} \end{matrix}$$

$$\text{last step: Show } y_1 = e^{ax} \cos bx \\ y_2 = e^{ax} \sin bx$$

are linearly independent:

Exercise Find the general solution of

$$y'' + 4y' + 5y = 0.$$

3

Using the ideas we've been discussing, you get the following algorithm to find all sol's to $L(y)=0$ for the const. coeff. linear DE of order n :

Theorem 9: Let $L(y) = y^{(n)} + a_{n-1}y^{(n-1)} + \dots + a_1y' + a_0y$ a_j const.
 $p(r) = r^n + a_{n-1}r^{n-1} + \dots + a_1r + a_0$

if r_j is a real root of p , $p(r_j)=0 \Rightarrow y = e^{r_j x}$ solves $L(y)=0$

if $(r-r_j)^{k_j}$ is a factor of $p \Rightarrow$
 $y_1 = e^{r_j x}$
 $y_2 = x e^{r_j x}$
 $\vdots$
 $y_{k_j} = x^{k_j-1} e^{r_j x}$ are k_j lin. ind sol's to $L(y)=0$

if $r_j = a+bi$ is a complex root, $p(a+bi)=0 \Rightarrow$
 $y_1 = e^{ax} \cos bx$
 $y_2 = e^{ax} \sin bx$ are lin. ind sol's to $L(y)=0$

if $(r-r_j)^{k_j}$ is a factor of p

also get

$x e^{ax} \cos bx, x e^{ax} \sin bx$

$\vdots$

$x^{k_j-1} e^{ax} \cos bx, x^{k_j-1} e^{ax} \sin bx$

(gives $2k_j$ sol's for the roots r_j & $\bar{r}_j$)

In this way we can always find a basis for $\ker(L)$, (assuming we can figure out how to factor p !)

Exercise Find the general solution to

$$y''' - 5y'' + 24y' - 20y = 0!$$

Applications on Friday, § 3.4

should be fun
 $\ddot{v}$