

Math 2280-001

Week 12, April 3-7

Mon Apr 3 Finish section 5.7 notes from Friday. We will also discuss questions you may have about the section 5.6 homework due on Tuesday. A lab problem for next week will be to work the input-output problem from the second midterm using (1) matrix exponentials, and (2) the diagonalization method. We may begin that problem in lab format, if time permits.

Wed Apr 5

## 7.1-7.2 Laplace transform, and application to DE IVPs

- The Laplace transform is a linear transformation " $\mathcal{L}$ " that converts piecewise continuous functions  $f(t)$ , defined for  $t \geq 0$  and with at most exponential growth ( $|f(t)| \leq Ce^{Mt}$  for some values of  $C$  and  $M$ ), into functions  $F(s)$  defined by the transformation formula

$$F(s) = \mathcal{L}\{f(t)\}(s) := \int_0^{\infty} f(t)e^{-st} dt.$$

- Notice that the integral formula for  $F(s)$  is only defined for sufficiently large  $s$ , and certainly for  $s > M$ , because as soon as  $s > M$  the integrand is decaying exponentially, so the improper integral from  $t = 0$  to  $\infty$  converges.
- The convention is to use lower case letters for the input functions and (the same) capital letters for their Laplace transforms, as we did for  $f(t)$  and  $F(s)$  above. Thus if we called the input function  $x(t)$  then we would denote the Laplace transform by  $X(s)$ .

Taking Laplace transforms seems like a strange thing to do. And yet, the Laplace transform  $\mathcal{L}$  is just one example of a collection of useful "integral transforms".  $\mathcal{L}$  is especially good for solving IVPs for linear DEs, as we shall see starting today. Other famous transforms - e.g. Fourier series and Fourier transform are extremely important in studying linear differential and partial differential equations. We will discuss Fourier series in about a week. These transforms are also studied in Math 3140, 3150, and in various 5000-level pure and applied math classes.

Exercise 1) Use the definition of Laplace transform

$$F(s) = \mathcal{L}\{f(t)\}(s) := \int_0^{\infty} f(t)e^{-st} dt$$

to check the following facts, which you will also find inside the front cover of your text book.

a)  $\mathcal{L}\{1\}(s) = \frac{1}{s} \quad (s > 0)$

b)  $\mathcal{L}\{e^{\alpha t}\}(s) = \frac{1}{s - \alpha} \quad (s > \alpha \text{ if } \alpha \in \mathbb{R}, s > a \text{ if } \alpha = a + ki \in \mathbb{C})$

c) Laplace transform is linear, i.e.

$$\mathcal{L}\{f_1(t) + f_2(t)\}(s) = F_1(s) + F_2(s).$$

$$\mathcal{L}\{cf(t)\}(s) = cF(s).$$

d) Use linearity and your work above to compute  $\mathcal{L}\{3 - 4e^{-2t}\}(s)$ .

For the linear differential equations and systems of differential equations that we've been studying, the following Laplace transforms are very important:

Exercise 2) Use complex number algebra, including Euler's formula, linearity, and the result from 1b that

$$\mathcal{L}\{e^{(a + ki)t}\}(s) = \frac{1}{s - (a + ki)}$$

to verify that

$$\underline{a)} \quad \mathcal{L}\{\cos(kt)\}(s) = \frac{s}{s^2 + k^2}$$

$$\underline{b)} \quad \mathcal{L}\{\sin(kt)\}(s) = \frac{k}{s^2 + k^2}$$

$$\underline{c)} \quad \mathcal{L}\{e^{at}\cos(kt)\}(s) = \frac{s - a}{(s - a)^2 + k^2}$$

$$\underline{d)} \quad \mathcal{L}\{e^{at}\sin(kt)\}(s) = \frac{k}{(s - a)^2 + k^2} .$$

(Notice that if we tried doing these Laplace transforms directly from the definition, the integrals would be messy but we could attack them via integration by parts or integral tables.)

It's a theorem (hard to prove but true) that a given Laplace transform  $F(s)$  can arise from at most one piecewise continuous function  $f(t)$ . (Well, except that the values of  $f$  at the points of discontinuity can be arbitrary, as they don't affect the integral used to compute  $F(s)$ .) Therefore you can read Laplace transform tables in either direction, i.e. not only to deduce Laplace transforms, but inverse Laplace transforms  $\mathcal{L}^{-1}\{F(s)\}(t) = f(t)$  as well.

Exercise 3) Use the Laplace transforms we've computed and linearity to compute

$$\mathcal{L}^{-1}\left\{\frac{7}{s} + \frac{1}{s^2 + 16} - \frac{10s}{s^2 + 16}\right\}(t) .$$

$f(t)$ $ f(t)  \leq Ce^{Mt}$	$F(s) := \int_0^\infty f(t)e^{-st} dt$ for $s > M$
$c_1 f_1(t) + c_2 f_2(t)$	$c_1 F_1(s) + c_2 F_2(s)$
1	$\frac{1}{s} \quad (s > 0)$
$e^{\alpha t}$	$\frac{1}{s - \alpha} \quad (s > \Re(\alpha))$
$\cos(kt)$	$\frac{s}{s^2 + k^2} \quad (s > 0)$
$\sin(kt)$	$\frac{k}{s^2 + k^2} \quad (s > 0)$
$e^{at}\cos(kt)$	$\frac{(s - a)}{(s - a)^2 + k^2} \quad (s > a)$
$e^{at}\sin(kt)$	$\frac{k}{(s - a)^2 + k^2} \quad (s > a)$
$f'(t)$	$sF(s) - f(0)$
$f''(t)$	$s^2F(s) - sf(0) - f'(0)$

**Laplace transform table**

The integral transforms of DE's and PDE's were designed to have the property that they convert the corresponding linear DE and PDE problems into algebra problems. For the Laplace transform it's because of these facts:

Exercise 4a) Use integration by parts and the definition of Laplace transform to show that

$$\mathcal{L}\{g'(t)\}(s) = s \mathcal{L}\{g(t)\}(s) - g(0) = s G(s) - g(0) .$$

4b) Use the result of a, applied to the function  $f'(t)$  to show that

$$\mathcal{L}\{f''(t)\}(s) = s^2 F(s) - s f(0) - f'(0) .$$

4c) What would you guess is the Laplace transform of  $f'''(t)$  ? Could you check this?

Here's an example of using Laplace transforms to solve DE IVPs, in the context of Chapter 3 and the mechanical (and electrical) application problems we considered there.

Exercise 5) Consider the undamped forced oscillation IVP

$$\begin{aligned} x''(t) + 4x(t) &= 10 \cos(3t) \\ x(0) &= 2 \\ x'(0) &= 1 \end{aligned}$$

If  $x(t)$  is the solution, then both sides of the DE are equal. Thus the Laplace transforms are equal as well... so, equate the Laplace transforms of each side and use algebra to find  $\mathcal{L}\{x(t)\}(s) = X(s)$ . Notice you've computed  $X(s)$  without actually knowing  $x(t)$ ! If you were happy to stay in "Laplace land" you'd be done. In any case, at this point you can use our table entries to find  $x(t) = \mathcal{L}^{-1}\{X(s)\}$ .

(Notice that if your algebra skills are good you've avoided having to use the Chapter 3 algorithm of (i) find  $x_H$  (ii) find an  $x_P$  (iii)  $x = x_P + x_H$  (iv) solve IVP.) Magic! Or, would you have preferred to convert to a first order system and to have used variation of parameters with an FM or  $e^{At}$ , like in Chapter 5 :- ) ? )



Input:

$$\{x''(t) + 4x(t) = 10 \cos(3t), x(0) = 2, x'(0) = 1\}$$

[Open code](#)

ODE classification:

second-order linear ordinary differential equation

Alternate forms: [More](#)

$$\{x''(t) = 10 \cos(3t) - 4x(t), x(0) = 2, x'(0) = 1\}$$


$$\{x''(t) + 4x(t) = 10 \cos(t) (2 \cos(2t) - 1), x(0) = 2, x'(0) = 1\}$$

$$\{x''(t) + 4x(t) = 10 \cos^3(t) - 30 \cos(t) \sin^2(t), x(0) = 2, x'(0) = 1\}$$

Differential equation solution: [Step-by-step solution](#)

$$x(t) = \frac{1}{2} (\sin(2t) + 8 \cos(2t) - 4 \cos(3t))$$

Plots of the solution:



[Enlarge](#) | [Data](#) | [Customize](#) | [Plaintext](#) | [Interactive](#)

Input:

partial fractions

$$10 \times \frac{s}{(s^2 + 9)(s^2 + 4)} + \frac{2s + 1}{s^2 + 4}$$

[Open code](#)

Result: [Step-by-step solution](#)

$$\frac{10s}{(s^2 + 4)(s^2 + 9)} + \frac{2s + 1}{s^2 + 4} = \frac{4s + 1}{s^2 + 4} - \frac{2s}{s^2 + 9}$$

inverse Laplace transform

Assuming "inverse Laplace transform" refers to a computation | Use as [referring to a mathematical definition](#) instead

function to transform:  $(4s+1)/(s^2+4) - 2s/(s^2+9)$

initial variable:

transform variable:

Input:

$$\mathcal{L}_s^{-1} \left[ \frac{4s+1}{s^2+4} - 2 \times \frac{s}{s^2+9} \right] (t)$$

[Open code](#)

$\mathcal{L}_s^{-1}[f(s)](t)$  is the inverse Laplace transform of  $f(s)$  with real variable  $t$

Result:

$$\frac{1}{2} (\sin(2t) + 8 \cos(2t) - 4 \cos(3t))$$

Exercise 6) Use Laplace transform as above, to solve the IVP for the following underdamped, unforced oscillator DE:

$$\begin{aligned}x''(t) + 6x'(t) + 34x(t) &= 0 \\x(0) &= 3 \\x'(0) &= 1\end{aligned}$$



Math 2280-001

Fri Apr 7

7.1-7.4 Laplace transform, and application to DE IVPs, especially those in Chapter 3. Today we'll continue to fill in the Laplace transform table (at the end of the notes). Along the way we'll revisit some of the mechanical oscillation differential equations from Chapter 3.

Exercise 1) (to review) Use the table to compute


1a)  $\mathcal{L}\{4 + 2e^{-4t}\}(s)$

1b)  $\mathcal{L}^{-1}\left\{\frac{2}{s-2} + \frac{6}{s}\right\}(t)$ .

Exercise 2) (to review the definition) Use the definition of Laplace transform,

$$F(s) = \mathcal{L}\{f(t)\}(s) := \int_0^{\infty} f(t)e^{-st} dt$$

to find the Laplace transform of the step function graphed below. (The function is equal to zero for  $t \geq 3$ .)



Exercise 3a) Use the Table entry we proved on Wednesday for derivatives (via integration by parts), namely

$$\mathcal{L}\{g'(t)\}(s) = s \mathcal{L}\{g(t)\}(s) - g(0) = s G(s) - g(0)$$

and math induction to show that for  $n \in \mathbb{N}$

$$\mathcal{L}\{f^{(n)}(t)\}(s) = s^n F(s) - s^{n-1}f(0) - s^{n-2}f'(0) - \dots - s f^{(n-2)}(0) - f^{(n-1)}(0) .$$

3b) (Integrals are "negative" derivatives): Use the Laplace transform first-derivative formula above to show that

$$\begin{aligned} \mathcal{L}\left\{\int_0^t f(\tau) \, d\tau\right\}(s) &= \frac{1}{s} \mathcal{L}\{f(t)\}(s) = \frac{F(s)}{s} \\ \mathcal{L}\left\{\int_0^t \left(\int_0^r f(\tau) \, d\tau\right) dr\right\}(s) &= \frac{F(s)}{s^2} \dots \end{aligned}$$

Exercise 4) Use the result of 3a to show that for  $n \in \mathbb{N}$ ,

$$\mathcal{L}\{t\}(s) = \frac{1}{s^2}$$

$$\mathcal{L}\{t^2\}(s) = \frac{2}{s^3}$$

$$\mathcal{L}\{t^n\}(s) = \frac{n!}{s^{n+1}}.$$

Exercise 5) Find  $\mathcal{L}^{-1}\left\{\frac{1}{s(s^2 + 4)}\right\}(t)$

a) using the result of 3b.

b) using partial fractions.

Exercise 6) (first translation theorem). Use the definition of Laplace transform to show that

$$\mathcal{L}\{e^{at}f(t)\}(s) = \mathcal{L}\{f(t)\}(s-a) = F(s-a)$$

Exercise 7) As a special case of Exercise 6, show

$$\begin{aligned}\mathcal{L}\{te^{at}\}(s) &= \frac{1}{(s-a)^2} . \\ \mathcal{L}\{t^n e^{at}\}(s) &= \frac{n!}{(s-a)^{n+1}}\end{aligned}$$

A harder table entry to understand is the following one - go through this computation and see why it seems reasonable, even though there's one step that we don't completely justify. The table entry is

$tf(t)$	$-F'(s)$
---------	----------

Here's how we get it:

$$\begin{aligned}F(s) &= \mathcal{L}\{f(t)\}(s) := \int_0^\infty f(t)e^{-st} dt \\ \Rightarrow \frac{d}{ds}F(s) &= \frac{d}{ds} \int_0^\infty f(t)e^{-st} dt = \int_0^\infty \frac{d}{ds}f(t)e^{-st} dt .\end{aligned}$$

It's this last step which is true, but needs more justification. We know that the derivative of a sum is the sum of the derivatives, and the integral is a limit of Riemann sums, so this step does at least seem reasonable. The rest is straightforward:

$$\int_0^\infty \frac{d}{ds}f(t)e^{-st} dt = \int_0^\infty f(t)(-t)e^{-st} dt = -\mathcal{L}\{tf(t)\}(s) \quad \square.$$

For resonance and other applications ...

Exercise 8) Use  $\mathcal{L}\{tf(t)\}(s) = -F'(s)$  directly, or Euler's formula and  $\mathcal{L}\{te^{\alpha t}\}(s) = \frac{1}{(s - \alpha)^2}$  to

show

a)  $\mathcal{L}\{t \cos(kt)\}(s) = \frac{s^2 - k^2}{(s^2 + k^2)^2}$

b)  $\mathcal{L}\left\{\frac{1}{2k} t \sin(kt)\right\}(s) = \frac{s}{(s^2 + k^2)^2}$

c) Then use a and the identity

$$\frac{1}{(s^2 + k^2)^2} = \frac{1}{2k^2} \left( \frac{s^2 + k^2}{(s^2 + k^2)^2} - \frac{s^2 - k^2}{(s^2 + k^2)^2} \right)$$

to verify the table entry

$$\mathcal{L}^{-1}\left\{\frac{1}{(s^2 + k^2)^2}\right\}(t) = \frac{1}{2k^2} \left( \frac{1}{k} \sin(kt) - t \cos(kt) \right).$$

Exercise 9) Use Laplace transforms to write down the solution to

$$\begin{aligned}x''(t) + \omega_0^2 x(t) &= F_0 \sin(\omega_0 t) \\ x(0) &= x_0 \\ x'(0) &= v_0.\end{aligned}$$

what phenomena do solutions to this DE illustrate (even though we're forcing with  $\sin(\omega_0 t)$  rather than  $\cos(\omega_0 t)$ )? How would you have tried to solve this problem in Chapter 3?

Exercise 10) Solve the following IVP. Use this example to recall the general partial fractions algorithm.

$$\begin{aligned}x''(t) + 4x(t) &= 8te^{2t} \\ x(0) &= 0 \\ x'(0) &= 1\end{aligned}$$

$f(t), \text{ with }  f(t)  \leq C e^{M t}$	$F(s) := \int_0^\infty f(t) e^{-s t} dt \text{ for } s > M$	$\downarrow$ verified
$c_1 f_1(t) + c_2 f_2(t)$	$c_1 F_1(s) + c_2 F_2(s)$	<input type="checkbox"/>
$1$ $t$ $t^2$ $t^n, n \in \mathbb{N}$	$\frac{1}{s} \quad (s > 0)$ $\frac{1}{s^2}$ $\frac{2}{s^3}$ $\frac{n!}{s^{n+1}}$	<input type="checkbox"/>
$e^{\alpha t}$	$\frac{1}{s - \alpha} \quad (s > \Re(\alpha))$	<input type="checkbox"/>
$\cos(k t)$ $\sin(k t)$ $\cosh(k t)$ $\sinh(k t)$ $e^{a t} \cos(k t)$ $e^{a t} \sin(k t)$	$\frac{s}{s^2 + k^2} \quad (s > 0)$ $\frac{k}{s^2 + k^2} \quad (s > 0)$ $\frac{s}{s^2 - k^2} \quad (s > k)$ $\frac{k}{s^2 - k^2} \quad (s > k)$ $\frac{(s - a)}{(s - a)^2 + k^2} \quad (s > a)$ $\frac{k}{(s - a)^2 + k^2} \quad (s > a)$	<input type="checkbox"/> <input type="checkbox"/>  <input type="checkbox"/> <input type="checkbox"/>
$\frac{f'(t)}{f''(t)}$ $f^{(n)}(t), n \in \mathbb{N}$ $\int_0^t f(\tau) d\tau$	$\frac{s F(s) - f(0)}{s^2 F(s) - s f(0) - f'(0)}$ $\frac{s^n F(s) - s^{n-1} f(0) - \dots - f^{(n-1)}(0)}{F(s)}$ $\frac{1}{s}$	<input type="checkbox"/> <input type="checkbox"/>
$t f(t)$ $t^2 f(t)$ $t^n f(t), n \in \mathbb{Z}$	$\frac{-F'(s)}{F''(s)}$ $(-1)^n F^{(n)}(s)$	

$\frac{f(t)}{t}$	$\int_s^\infty F(\sigma) d\sigma$	
$t \cos(k t)$ $\frac{1}{2 k} t \sin(k t)$ $\frac{1}{2 k^3} (\sin(k t) - k t \cos(k t))$	$\frac{s^2 - k^2}{(s^2 + k^2)^2}$ $\frac{s}{(s^2 + k^2)^2}$ $\frac{1}{(s^2 + k^2)^2}$	
$e^{a t} f(t)$ $t e^{a t}$ $t^n e^{a t}, n \in \mathbb{Z}$	$F(s - a)$ $\frac{1}{(s - a)^2}$ $\frac{n!}{(s - a)^{n+1}}$	

**Laplace transform table**