


### 1. JORDAN LEMMA

Let  $R_0$  be a positive number and  $a$  a real number with  $|a| < R_0$ . Let

$$S = \{z \in \mathbb{C} \mid |z| \geq R_0 \text{ and } \text{Im } z \geq a\}$$

be the region sketched in the following picture:



Let  $f$  be a continuous function on  $S$  such that  $\lim_{z \rightarrow \infty} f(z) = 0$ , i.e., for any  $\epsilon > 0$  there exists  $R \geq R_0$  such that  $z \in S$  and  $|z| > R$  implies that  $|f(z)| < \epsilon$ .

Let  $\gamma_R$  be the positively oriented arc determined as the intersection of the circle of radius  $R$  centered at the origin with  $S$ .

**1.1. Lemma (Jordan Lemma).** *For any  $m > 0$  we have*

$$\lim_{R \rightarrow \infty} \int_{\gamma_R} e^{imz} f(z) dz = 0.$$

Assume first that  $a$  is positive. Then  $\gamma_R$  is parameterized as  $\gamma_R(\varphi) = Re^{i\varphi}$  for  $\varphi \in [\alpha(R), \pi - \alpha(R)]$ . Therefore, we have


$$\begin{aligned} \left| \int_{\gamma_R} e^{imz} f(z) dz \right| &= \left| iR \int_{\alpha(R)}^{\pi - \alpha(R)} e^{imRe^{i\varphi}} f(Re^{i\varphi}) e^{i\varphi} d\varphi \right| \\ &\leq R \int_{\alpha(R)}^{\pi - \alpha(R)} |e^{imR(\cos \varphi + i \sin \varphi)}| |f(Re^{i\varphi})| d\varphi \\ &= R \int_{\alpha(R)}^{\pi - \alpha(R)} e^{-mR \sin \varphi} |f(Re^{i\varphi})| d\varphi \\ &\leq R \max_{z \in \gamma_R^*} |f(z)| \int_{\alpha(R)}^{\pi - \alpha(R)} e^{-mR \sin \varphi} d\varphi \\ &\leq R \max_{z \in \gamma_R^*} |f(z)| \int_0^\pi e^{-mR \sin \varphi} d\varphi = 2R \max_{z \in \gamma_R^*} |f(z)| \int_0^{\frac{\pi}{2}} e^{-mR \sin \varphi} d\varphi. \end{aligned}$$

Now we use the following simple lemma.

**1.2. Lemma.** *Let  $\varphi \in [0, \frac{\pi}{2}]$ . Then*

$$\sin \varphi \geq \frac{2}{\pi} \varphi.$$

*Proof.* This is clear from the following picture representing the graphs of the functions  $y = \sin x$  and  $y = \frac{2}{\pi}x$  which intersect at the origin and the point  $(\frac{\pi}{2}, 1)$ :



To get an analytic proof, consider the function

$$F(x) = \frac{\sin x}{x}$$

on  $(0, \frac{\pi}{2}]$ . Clearly, the function is differentiable on  $(0, \infty)$  and

$$F'(x) = \frac{x \cos x - \sin x}{x^2}.$$

Moreover, if  $G(x) = x \cos x - \sin x$ , we have

$$G'(x) = \cos x - x \sin x - \cos x = -x \sin x.$$

Hence,  $G'(x) \leq 0$  for  $x \in [0, \pi]$ , and the function is decreasing there, i.e.,  $G(x) \leq G(0) = 0$  for  $x \in [0, \pi]$ . This implies that  $F'(x) \leq 0$  for  $x \in (0, \pi]$  and the function is decreasing on this interval. In particular, for  $x \in (0, \frac{\pi}{2}]$ , we have  $F(x) \geq F(\frac{\pi}{2}) = \frac{2}{\pi}$ .  $\square$

Using this result, we see that

$$\begin{aligned} \int_0^{\frac{\pi}{2}} e^{-mR \sin \varphi} d\varphi &\leq \int_0^{\frac{\pi}{2}} e^{-\frac{2mR}{\pi} \varphi} d\varphi \\ &= -\frac{\pi}{2mR} \left[ e^{-\frac{2mR}{\pi} \varphi} \right]_0^{\frac{\pi}{2}} = \frac{\pi}{2mR} \left( 1 - e^{-mR} \right) \leq \frac{\pi}{2mR}. \end{aligned}$$

Combining this with the above we get

$$\left| \int_{\gamma_R} e^{imz} f(z) dz \right| \leq \frac{\pi}{m} \max_{z \in \gamma_R^*} |f(z)|.$$

Therefore, the left side tends to 0 as  $R \rightarrow \infty$ . This proves the lemma in this case.

If  $a < 0$ , the path  $\gamma_R$  is parameterized as  $\gamma_R(\varphi) = Re^{i\varphi}$  where  $\varphi \in [-\alpha(R), \pi + \alpha(R)]$ . Therefore,  $\gamma_R$  consists of three positively oriented arcs: the arc  $\gamma_R''$  for  $\varphi \in [-\alpha(R), 0]$ , followed by  $\gamma_R'$  for  $\varphi \in [0, \pi]$  and  $\gamma_R'''$  for  $\varphi \in [\pi, \pi + \alpha(R)]$ . It follows that

$$\int_{\gamma_R} e^{imz} f(z) dz = \int_{\gamma_R''} e^{imz} f(z) dz + \int_{\gamma_R'} e^{imz} f(z) dz + \int_{\gamma_R'''} e^{imz} f(z) dz.$$

The second integral tends to zero as  $R \rightarrow \infty$  by the first part of the proof. As before, the first integral satisfies

$$\begin{aligned} \left| \int_{\gamma_R''} e^{imz} f(z) dz \right| &= \left| iR \int_{-\alpha(R)}^0 e^{imRe^{i\varphi}} f(Re^{i\varphi}) e^{i\varphi} d\varphi \right| \\ &\leq R \int_{-\alpha(R)}^0 e^{-mR \sin \varphi} |f(Re^{i\varphi})| d\varphi. \end{aligned}$$

From the geometry of our situation, we see that  $\sin \alpha(R) = \frac{|a|}{R}$ . Since  $\alpha(R) < \frac{\pi}{2}$ , we see that

$$-\sin \varphi \leq \sin \alpha(R) = \frac{|a|}{R}$$

for  $\varphi \in [-\alpha(R), 0]$ . It follows that

$$\left| \int_{\gamma_R''} e^{imz} f(z) dz \right| \leq e^{m|a|} \max_{z \in \gamma_R^*} |f(z)| \ell(\gamma_R'') = e^{m|a|} \max_{z \in \gamma_R^*} |f(z)| R \arcsin \left( \frac{|a|}{R} \right).$$

Now, we have

$$\lim_{R \rightarrow \infty} R \arcsin \left( \frac{|a|}{R} \right) = \lim_{s \rightarrow 0} \frac{\arcsin(|a|s)}{s} = |a| \lim_{s \rightarrow 0} \frac{\arcsin s}{s} = |a| \lim_{t \rightarrow 0} \frac{t}{\sin t} = |a|,$$

and this implies that the above integral tends to zero as  $R \rightarrow \infty$ . The argument for  $\gamma_R'''$  is analogous.