

Midterm Exam
Math 21B: Section 001
Summer 2012

Name _____

ID number _____

Directions

- Do not begin until instructed to do so. You will have 90 minutes to complete the exam.
- You may use pencils, pens, and erasers.
- Put away all books, notes, cell phones, calculators, and other electronic devices.
- Show all work for full credit. If in doubt, write it out.
- Keep your work as neat as possible. If we can't read it, we won't grade it!
- Keep your student ID out on your desk while working on your exam.
- You must also present your student ID to the instructor when turning in your exam.

Point totals

Problem	1	2	3	4	5	6	7	8	9	10	11	total
Score	/8	/12	/8	/8	/12	/16	/8	/12	/16	/16	/16	/132

1. (8 Points) Solve the following initial value problem:

$$\frac{ds}{dt} = \cos(t) + \sin(t), \quad s(\pi) = 1.$$

Solution: We know that $s(t)$ is an antiderivative of the given function. Therefore we calculate

$$s(t) = \int (\cos(t) + \sin(t)) dt = \sin(t) - \cos(t) + C. \quad \left. \begin{array}{l} 3 \text{ points for} \\ \text{taking anti-deriv.} \end{array} \right\}$$

We now use the given initial condition to solve for the unknown constant C .

$$1 = s(\pi) = \sin(\pi) - \cos(\pi) + C = 1 + C, \Rightarrow C = 0. \quad \left. \begin{array}{l} 3 \text{ points for setting } s(\pi) = 1 \\ 1 \text{ point} \end{array} \right\}$$

This fully determines the function $s(t) = \sin(t) - \cos(t)$

1 point for writing some function of t for $s(t)$

~~2 points~~
for calculating proper constant.

2. (12 Points) Suppose that f and g are both integrable functions. You are given that

$$\int_1^3 f(x) dx = -4, \quad \int_1^5 f(x) dx = 1, \quad \int_{-1}^3 g(x) dx = 7/2, \quad \int_{-1}^5 g(x) dx = 1/2.$$

Use these facts to calculate the definite integral

$$\int_5^3 (f(x) - 2g(x)) dx.$$

Solution: We use the properties of the definite integral

$$\begin{aligned} \int_5^3 (f(x) - 2g(x)) dx &= - \int_3^5 (f(x) - 2g(x)) dx && \left. \begin{array}{l} 3 \text{ points for switching} \\ \text{limits of integration} \end{array} \right\} \\ &= \int_3^5 (2g(x) - f(x)) dx \\ &= 2 \int_3^5 g(x) dx - \int_3^5 f(x) dx && \left. \begin{array}{l} 3 \text{ points for breaking} \\ \text{Integrand} \end{array} \right\} \\ &= 2 \left(\int_{-1}^5 g(x) dx - \int_{-1}^3 g(x) dx \right) - \left(\int_1^5 f(x) dx - \int_1^3 f(x) dx \right) && \left. \begin{array}{l} 3 \text{ points} \\ \text{for} \\ \text{breaking} \\ \text{Interval} \end{array} \right\} \\ &= 2 \left(\underbrace{1/2 - 7/2} \right) - \left(\underbrace{1 - (-4)} \right) = -11. \\ & \left. \begin{array}{l} 3 \text{ points for using} \\ \text{given information.} \end{array} \right\} \end{aligned}$$

3. (8 Points) Calculate the definite integral

$$\int_0^1 xe^{x^2} dx.$$

Solution: We let $u = x^2$. A quick calculation shows $du = 2xdx$ and thus $x dx = \frac{1}{2} du$. When $x = 0$ we have $u = 0$, and similarly when $x = 1$, $u = 1$. We use these facts to make the substitution

$$\int_0^1 xe^{x^2} dx = \int_0^1 \frac{1}{2} e^u du$$

$$= \frac{1}{2} e^u \Big|_0^1$$

$$= \frac{1}{2} (e^1 - e^0)$$

$$= \frac{e-1}{2}$$

1 point for ~~each~~ a working candidate for u

3 points for properly substituting integral limits

2 points for finding correct anti-derivative

2 points

for evaluating properly

4. (8 Points) Find the indefinite integral

$$\int \frac{t\sqrt[3]{t} - \sqrt[3]{t^2}}{t^2} dt$$

Solution:

$$\int \frac{t\sqrt[3]{t} - \sqrt[3]{t^2}}{t^2} dt = \int \frac{t \cdot t^{1/3} - (t^2)^{1/3}}{t^2} dt$$

$$= \int \frac{t^{4/3} - t^{2/3}}{t^2} dt$$


$$= \int t^{-2/3} - t^{-4/3} dt$$

$$= 3t^{1/3} + 3t^{-1/3} + C.$$

4 points for simplifying Integrand

4 points for proper anti-derivative

5. (12 Points) Sketch the region between the curves $y = 1 + \cos(x)$, $y = 2$, and $x = \pi$. Then calculate the area of this region. Solution: The region in question is pictured in the figure below.



4 points for figure

We calculate its area using the formula $A = \int_a^b h(x) dx$. The function $h(x)$ is the height of the region of interest, which in this case is from the top curve ($y = 2$) to the bottom curve ($y = 1 + \cos(x)$). Therefore

$$h(x) = 2 - (1 + \cos(x)) = 1 - \cos(x).$$

4 points for finding proper function to integrate

As can be seen in the figure, the region spans from $x = 0$ to $x = \pi$. Therefore

$$\begin{aligned} A &= \int_0^{\pi} (1 - \cos(x)) dx \\ &= x - \sin(x) \Big|_0^{\pi} \\ &= (\pi - \sin(\pi)) - (0 - \sin(0)) \\ &= \pi \end{aligned}$$

4 points for calculating integral properly

6. (16 Points) Solve the following initial value problem:

$$\frac{d^2y}{d\theta^2} = 4 \sec^2(2\theta) \tan(2\theta), \quad y'(0) = 4, \quad y(0) = -1.$$

Solution: As before, we begin by finding the general antiderivative. To do this we make the substitution $u = \sec(2\theta)$. A quick calculation shows that $du = 2 \sec(2\theta) \tan(2\theta) d\theta$.

Therefore

$$\int 4 \sec^2(2\theta) \tan(2\theta) d\theta = \int 2u du = u^2 + C = \sec^2(2\theta) + C.$$

} 5 points for first int. ~~derivative~~

We use the initial condition for $y'(\theta)$ to solve for the unknown constant

$$4 = y'(0) = \sec^2(0) + C \Rightarrow C = 3.$$

} 3 points for first constant

This gives us that $\frac{dy}{d\theta} = \sec^2(2\theta) + 3$. We repeat this process to find $y(\theta)$. Taking the antiderivative we have

$$\int \frac{dy}{d\theta} d\theta = \int (\sec^2(2\theta) + 3) d\theta = \frac{1}{2} \tan(2\theta) + 3\theta + C.$$

} 5 points for 2nd Integral

Using our initial condition we see

$$-1 = y(0) = \frac{1}{2} \tan(0) + 3(0) + C \Rightarrow C = -1.$$

Therefore

$$y(\theta) = \frac{1}{2} \tan(2\theta) + 3\theta - 1.$$

} 3 points for 2nd constant & final answer.

7. (8 Points) Find the average value of the function $f(x) = \sin(x)$ over the interval $[0, 7\pi]$.

Solution: We know that the average value of a function over an interval is given by

$\bar{f} = \frac{1}{b-a} \int_a^b f(x) dx$. Therefore we calculate

$$\bar{f} = \frac{1}{7\pi - 0} \int_0^{7\pi} \sin(x) dx$$

} 4 points for setting up Integral

$$= \frac{-1}{7\pi} \cos(x) \Big|_0^{7\pi}$$

$$= \frac{-1}{7\pi} (\cos(7\pi) - \cos(0))$$

$$= \frac{2}{7\pi}$$

} 4 points for calculation.

8. (12 Points) Find the length of the curve given by $y = \frac{2}{3}(x^2 + 1)^{3/2}$ on the interval $0 \leq x \leq 2$.

Solution: We begin by recalling the formula for arc length:

$$L = \int_a^b \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx.$$

In order to use this formula, we calculate

$$\frac{dy}{dx} = 2x(x^2 + 1)^{1/2}.$$

← 2 points for calculating derivative

$$\left(\frac{dy}{dx}\right)^2 = 4x^2(x^2 + 1) = 4x^4 + 4x^2.$$

$$1 + \left(\frac{dy}{dx}\right)^2 = 4x^4 + 4x^2 + 1 = (2x^2 + 1)^2,$$

and therefore

$$\sqrt{1 + \left(\frac{dy}{dx}\right)^2} = 2x^2 + 1.$$

4 points for calculating arc length element.

This allows us to use the arc length formula and calculate


$$\begin{aligned} L &= \int_0^2 (2x^2 + 1) dx \\ &= \left. \frac{2}{3}x^3 + x \right|_0^2 \\ &= \frac{2}{3}(2^3) + 2 \\ &= \frac{22}{3} \end{aligned}$$

← 3 points for setting up integral

3 points for evaluating integral

9. (16 Points) Calculate the volume of a sphere of radius 3 by rotating a semi-circular arc (of radius 3) about the x-axis (HINT: The equation for a circle of radius r in the x-y plane is $y^2 + x^2 = r^2$).

Solution: We begin by drawing the semi-circle (of radius 3) in the top half plane. We define the distance from the x-axis to this curve to be $r(x)$.



4/3 points
for defining
a working ~~area~~ region
& axis of rotation

The formula given in the hint allows us to solve for $r(x) = y = \sqrt{9 - x^2}$. Using the disk method, we know that rotating this semi-circle about the x-axis will produce a volume $V = \int_a^b \pi r(x)^2 dx$. This is the sphere we are interested in. Our limits of integration are where the curve touches the x-axis, which is $x = -3$ and $x = 3$. Using this information, we calculate


$$\begin{aligned}
 V &= \int_{-3}^3 \pi (\sqrt{9 - x^2})^2 dx \\
 &= \int_{-3}^3 \pi (9 - x^2) dx \\
 &= \pi \left(9x - \frac{x^3}{3} \right) \Big|_{-3}^3 \\
 &= \pi \left(27 - \frac{27}{3} \right) - \pi \left(-27 + \frac{27}{3} \right) \\
 &= \pi (27 - 9 + 27 - 9) \\
 &= 36\pi
 \end{aligned}$$

8 points for setting up proper Integral

4 points for evaluating Integral

10. (16 Points) Calculate the volume of the solid that is generated when you take the region bounded by the x-axis and the graph of $f(x) = \sin(x^2)$ (on the interval $0 \leq x \leq \sqrt{\pi}$) and revolve it about the y-axis.

Solution: We will calculate this volume using the method of cylindrical shells. Begin by sketching the region being rotated and drawing a slice through it parallel to the axis of revolution.



This slice has a thickness that we call dx . It is a distance $r(x) = x$ from the axis of revolution. It stretches from the x-axis to the curve $y = \sin(x^2)$. Therefore the height of this slice is $h(x) = \sin(x^2)$. The method of cylindrical shells tells us that the volume of the solid of revolution is

$$V = \int_a^b 2\pi r(x)h(x) dx = \int_0^{\sqrt{\pi}} 2\pi x \sin(x^2) dx.$$

8 points for setting up shells Integral

To evaluate this integral, we make the substitution $u = x^2$. A quick calculation shows us that $du = 2x dx$, when $x = 0$, $u = 0$, and when $x = \sqrt{\pi}$, $u = \pi$. Using this information, we make a U-substitution and get

$$V = \int_0^{\pi} \pi \sin(u) du = -\pi \cos(u) \Big|_0^{\pi} = 2\pi.$$


2 points for Evaluating Integral.

6 points for u-sub

11. (16 Points) Calculate the lateral surface area (*not* including the circular end) of the solid that is generated when you take the graph of the function $g(y) = y^3/3$ on the interval $0 \leq y \leq 1$ and revolve it about the y -axis.

Solution: The figure below shows the curve that will be rotated about the y -axis. Recall that the area for surfaces of revolution is

$$A = \int_a^b 2\pi x(y) \sqrt{1 + \left(\frac{dx}{dy}\right)^2} dy.$$



We begin by calculating the terms needed to evaluate this integral:

$$\frac{dx}{dy} = y^2, \quad \left(\frac{dx}{dy}\right)^2 = y^4, \quad \sqrt{1 + \left(\frac{dx}{dy}\right)^2} = \sqrt{y^4 + 1}.$$

4 points for calculating arc length element.

The problem statement tells us that we will be integrating from $y = 0$ to $y = 1$.

Therefore we must evaluate the integral

$$A = \int_0^1 \frac{2\pi}{3} y^3 \sqrt{y^4 + 1} dy.$$

8 points for setting up proper integral

To do this, we make the substitution that $u = y^4 + 1$. A calculation shows us that $du = 4y^3 dy$, and therefore $\frac{2}{3}y^3 dy = \frac{1}{6} du$. Furthermore, when $y = 0$, we have $u = 1$, and when $y = 1$, $u = 2$. This allows us to make the substitution

$$A = \int_0^1 \frac{2\pi}{3} y^3 \sqrt{y^4 + 1} dy$$

$$= \int_1^2 \frac{\pi}{6} \sqrt{u} du$$

$$= \frac{\pi}{9} u^{3/2} \Big|_1^2$$

$$= \frac{\pi}{9} (2^{3/2} - 1^{3/2})$$

$$= \frac{\pi}{9} (2\sqrt{2} - 1)$$

2 points for evaluating Integral

2 points for u-sub

