

GROWTH IN THE MINIMAL INJECTIVE RESOLUTION OF A LOCAL RING

LARS WINTHER CHRISTENSEN, JANET STRIULI, AND OANA VELICHE

ABSTRACT. Let R be a commutative noetherian local ring with residue field k and assume that it is not Gorenstein. In the minimal injective resolution of R , the injective envelope E of the residue field appears as a summand in every degree starting from the depth of R . The number of copies of E in degree i equals the k -vector space dimension of the cohomology module $\text{Ext}_R^i(k, R)$. These dimensions, known as Bass numbers, form an infinite sequence of invariants of R about which little is known. We prove that it is non-decreasing and grows exponentially if R is Golod, a non-trivial fiber product, or Teter, or if it has radical cube zero.

1. INTRODUCTION

In this paper R is a commutative noetherian local ring with maximal ideal \mathfrak{m} . It is a pervasive theme in local algebra that properties of R can be retrieved from homological invariants of the residue field $k = R/\mathfrak{m}$. A prime example is vanishing of cohomology with coefficients in k . Indeed, R is regular if and only if $\text{Ext}_R^i(k, k) = 0$ for all $i \gg 0$, and R is Gorenstein if and only if $\text{Ext}_R^i(k, R) = 0$ for all $i \gg 0$.

The cohomology groups $\text{Ext}_R^i(k, k)$ behave rigidly: if R is singular (i.e. not regular), then they are all non-zero. In this case, one focuses on their size, captured by the sequence of invariants $\beta_i^R(k) = \text{rank}_k \text{Ext}_R^i(k, k)$, called the *Betti numbers* of k . Through work pioneered by Gulliksen [12], the behavior of these Betti numbers is understood well enough to provide valuable information about singular local rings. The sequence $\{\beta_i^R(k)\}_{i \geq 0}$ is known to be non-decreasing; it is eventually constant if and only if R is a hypersurface, and it has polynomial growth if and only if R is a complete intersection. If R is not a complete intersection, then the Betti numbers are increasing and they grow exponentially.

As shown by Foxby [8], also the cohomology groups $\text{Ext}_R^i(k, R)$ behave rigidly: if R is not Gorenstein, then $\text{Ext}_R^i(k, R)$ is non-zero for all $i \geq \text{depth } R$. Their size is captured by the *Bass numbers* $\mu^i(R) = \text{rank}_k \text{Ext}_R^i(k, R)$, but these invariants are not understood nearly as well as the Betti numbers $\beta_i^R(k)$. It is not even known if the sequence $\{\mu^i(R)\}_{i \geq \text{depth } R}$ is non-decreasing. Existence of a local ring whose Bass numbers grow polynomially is also an open question.

Let M be a finitely generated R -module. For $i \geq 0$ the i^{th} *Betti number* of M is defined as $\beta_i^R(M) = \text{rank}_k \text{Ext}_R^i(M, k)$. Let \hat{R} denote the \mathfrak{m} -adic completion of R .

Date: 26 December 2008.

2000 Mathematics Subject Classification. Primary 13D02; secondary 13D07, 13H10.

Key words and phrases. Bass number, Betti number, minimal free resolution, minimal injective resolution.

This work started while L.W.C. visited the University of Nebraska-Lincoln, partly supported by a grant from the Carlsberg Foundation. J.S. was supported by NSF grant DMS 0201904.

Foxby [9] shows that there is a finitely generated \widehat{R} -module Ω such that one has

$$(1.0.1) \quad \mu^{\dim R+i}(R) = \beta_i^{\widehat{R}}(\Omega) \quad \text{for all } i \geq 0.$$

Thus, the non-zero Bass numbers of R , except the first $c := \dim R - \text{depth } R$, can be realized as the Betti numbers of a module. In particular, the sequence $\{\mu^i(R)\}_{i \geq d}$ has at most exponential growth. If R is Cohen–Macaulay, then $c = 0$ and Ω is the canonical module for \widehat{R} . Jorgensen and Leuschke [15] take this approach to prove that the sequence $\{\mu^i(R)\}_{i \geq \text{depth } R}$ grows exponentially and is eventually increasing for certain families of Cohen–Macaulay rings. Their work was motivated by a question of Huneke about the asymptotic behavior of these numbers; however, they are naturally lead to raise a question about the *initial* behavior [15, 2.6]:

(1.1) **Question.** *Let R be a Cohen–Macaulay local ring of depth d . If the inequality $\mu^{d+1}(R) \leq \mu^d(R)$ holds, is then R Gorenstein?*

There is already a history in local algebra for studying the initial behavior of Bass numbers—in particular, the number $\mu^{\dim R}(R)$ which is called the *type* of R . Bass [7] characterized Gorenstein rings as Cohen–Macaulay rings of type 1. Later, Foxby [9] and Roberts [22] proved that any local ring with $\mu^{\dim R}(R) = 1$ is Gorenstein. This confirmed a conjecture of Vasconcelos [27].

* * *

In this paper we study the initial, local, and asymptotic behavior of Bass numbers for certain families of local rings. While the realization of Bass numbers as Betti numbers (1.0.1) remains important for our work, it provides no information about the initial behavior of the Bass numbers of a ring that is not Cohen–Macaulay. We employ a bag of ad-hoc methods to deal with this situation.

The *embedding dimension* of R , denoted $\text{edim } R$, is the minimal number of generators of the maximal ideal. Question (1.1) is answered affirmatively in [15] when $\text{edim } R$ is $d + 2$, where d denotes the depth of R . This is the first interesting, case in as much as R is a hypersurface if $\text{edim } R \leq d + 1$. We improve the result from [15] by computing the Bass numbers of these rings in closed form; see (2.7). The conclusion is strong: if R is not Gorenstein, then there are (in)equalities

$$\mu^{d+1}(R) = \mu^d(R) + 1 \quad \text{and} \quad \mu^{i+1}(R) \geq 2\mu^i(R) \quad \text{for all } i \geq d + 1.$$

Similar detailed information is obtained for other rings. The Main Theorem below extracts the overall conclusion that applies to several families of rings. For the Cohen–Macaulay rings among them, it answers Question (1.1) affirmatively.

(1.2) **Main Theorem.** *Let $(R, \mathfrak{m}, \mathfrak{k})$ be a local ring of embedding dimension e at least 2 and depth d . Assume that R satisfies one of the following conditions:*

- (1) R is Golod and it has $e - d > 2$ or $\mu^d(R) > 1$.
- (2) R is the fiber product of two local rings (both different from \mathfrak{k}) and not Golod.
- (3) R is not Gorenstein and $\mathfrak{m}^3 = 0$.
- (4) R is Teter; that is, $R \cong Q/\text{Soc } Q$ where Q is artinian and Gorenstein.
- (5) R is artinian with $\text{Soc } R \not\subseteq \mathfrak{m}^2$.

Then the sequence of Bass numbers $\{\mu^i(R)\}_{i \geq d}$ is increasing and has exponential growth; if R satisfies (1), (3), (4), or (5), then the growth is termwise exponential.

Notice that the assumption $\text{edim } R \geq 2$ only excludes hypersurface. A sequence $\{a_i\}_{i \geq 0}$ is said to have *exponential growth* if there exists a real number $A > 1$ such that $a_i \geq A^i$ for all $i \gg 0$, and the growth is said to be *termwise exponential* of rate A if there exists a real number $A > 1$ such that $a_{i+1} \geq Aa_i$ for all $i \gg 0$.

Golod rings and fiber products are, in general, far from being Gorenstein, so one could expect their Bass numbers grow rapidly. Teter rings have been called “almost Gorenstein” [14], but they still differ significantly from Gorenstein rings, and some of them are even Golod. In this respect, (3) is the most surprising part of the Main Theorem, as there is empirical evidence in [1] and other works that rings with $\mathfrak{m}^3 = 0$ are excellent grounds for testing homological questions in local algebra.

Based on the Main Theorem—and the example below, which shows that two consecutive Bass numbers *can* be equal and non-zero—we extend and explicitly state the question that motivated Jorgensen and Leuschke’s work [15]:

(1.3) **Question.** *Let R be a local ring of depth d and assume that it is not Gorenstein. Does the sequence of Bass numbers $\{\mu^i(R)\}_{i \geq d}$ then have exponential growth, and is it non-decreasing? Is it eventually increasing, and if so, from which step?*

Our work towards the Main Theorem started from two explicit computations of Bass numbers. We discovered (1.4) while computing examples with the aid of MACAULAY 2 [11]; it can also be deduced from work of Wiebe [28].

(1.4) **Example.** Let F be a field. The first few Bass numbers of the local ring $F[[x, y]]/(x^2, xy)$ are 1, 2, 2, 4, 6, 10.

This ring is not Cohen–Macaulay, so the example says nothing about Question (1.1); it merely frames it. The ring is the archetype of the exceptional case in part (1) of the Main Theorem. We show in Proposition (2.8) that the Bass numbers of such rings (Golod with $e = d + 2$ and $\mu^d(R) = 1$) are given by the Fibonacci numbers as follows: $\mu^{d+i}(R) = 2F_i$ for all $i \geq 1$. In particular, the sequence $\{\mu^i(R)\}_{i \geq d}$ is non-decreasing with termwise exponential growth, and it increases from the 3rd step.

Another simple example [5, (10.8.2)] provides a textbook illustration of termwise exponential growth. The Main Theorem generalizes it in several directions.

(1.5) **Example.** For local ring (R, \mathfrak{m}) with $\mathfrak{m}^2 = 0$ and embedding dimension $e \geq 2$, the Bass numbers are $\mu^0(R) = e$ and $\mu^i(R) = e^{i-1}(e^2 - 1)$ for all $i \geq 1$.

These rings are in the intersection of the five families in the Main Theorem. Indeed, they are Golod and they trivially satisfy $\mathfrak{m}^3 = 0$, so they belong to (1) and (3). Since they have $\text{Soc } R = \mathfrak{m}$, they belong to (5) and, as will be explained in (3.7), also to (2). Finally, it is a result of Teter [26] that a local ring with $\mathfrak{m}^2 = 0$ is Teter.

* * *

The organization of the paper follows the agenda set by the Main Theorem (1.2). The Appendix has results on local and asymptotic behavior of Betti numbers for modules over artinian rings. These are used in the proofs of parts (3) and (5) of (1.2), which make heavy use of the realization of Bass number as Betti numbers (1.0.1). A curious upshot—an immediate consequence of Lemma (A.1)—is a reformulation of Bass’ characterization of Gorenstein rings:

(1.6) **Characterization.** *If R is a Cohen–Macaulay local ring and $\mu^n(R) = 1$ for some $n \geq 0$, then R is Gorenstein of dimension n .*

Example (1.4) shows that the Cohen–Macaulay hypothesis is necessary in (1.6), but it would be interesting to know if there is a similar reformulation of the result of Foxby [9] and Roberts [22]. The best one can hope for is a positive answer to:

(1.7) **Question.** *Let R be a local ring. If $\mu^n(R) = 1$ for some $n \geq \dim R$, is then R Gorenstein of dimension n ?*

If R is a domain, then an affirmative answer is already contained in [22].

2. GOLOD RINGS

The central result of this section, Theorem (2.5), is part (1) of the Main Theorem. Throughout the section, d denotes the depth of R and e its embedding dimension.

We use the standard notation for Koszul homology: given a sequence \mathbf{y} of elements in the maximal ideal \mathfrak{m} and an R -module M , the i^{th} homology module of the Koszul complex $K(\mathbf{y}) \otimes_R M$ is denoted $H_i(\mathbf{y}; M)$. Moreover, the notation $H_i(\mathbf{y}; R)$ is abbreviated $H_i(\mathbf{y})$; see also [24, IV.A.§1].

(2.1) Golod rings are the local rings for which the Betti numbers of the residue field have extremal growth; see [2, sec. 5]. All hypersurface rings are Golod, and a Golod ring is Gorenstein if and only if it is a hypersurface; see [2, rmk. after prop. 5.2.5]. The *codimension* of R is defined as $\text{codim } R = \text{edim } R - \dim R$. Every ring of codimension at most 1 is Golod; see [2, prop. 5.2.5]. Thus, the ring in Example (1.4) is Golod. So is the ring in Example (1.5), but for a different reason; see [2, prop. 5.3.4.(1)].

Here we use a characterization of Golod rings in terms of Bass numbers. The Bass numbers of R are encoded into a formal power series,

$$I_R(t) = \sum_{i=0}^{\infty} \mu^i(R) t^i,$$

called the *Bass series* of R .

Assume that R is singular, and let \mathbf{x} be a minimal system of generators for \mathfrak{m} . Avramov and Lescot [4, (0.2)] prove that there is a coefficient-wise inequality

$$(2.1.1) \quad I_R(t) \preceq \frac{\sum_{i=0}^{e-1} \text{rank}_{\mathbf{k}} H_{e-i}(\mathbf{x}) t^i - t^{e+1}}{1 - \sum_{i=1}^e \text{rank}_{\mathbf{k}} H_i(\mathbf{x}) t^{i+1}},$$

where equality holds if and only if R is Golod.

A crucial step in the proof of Theorem (2.5) is a reduction of (2.1.1), which comes about because -1 is a common root of the numerator and the denominator. One can deduce this from the work of Avramov, Iyengar, and Miller [3]. In Lemma (2.3) we provide a direct argument; first we split the coefficients $\text{rank}_{\mathbf{k}} H_j(\mathbf{x})$ in (2.1.1):

(2.2) **Lemma.** *Let R be singular, and let $\mathbf{x} = x_1, \dots, x_e$ be a minimal system of generators of \mathfrak{m} such that $\mathbf{x}' = x_1, \dots, x_{e-1}$ generates an \mathfrak{m} -primary ideal. For every integer i the Koszul homology module $H_0(x_e; H_i(\mathbf{x}'))$ is a finite dimensional \mathbf{k} -vector space, and it is non-zero if and only if $i \in \{0, \dots, e - d - 1\}$. Moreover, for every $i \geq 0$ there is an equality*

$$\text{rank}_{\mathbf{k}} H_i(\mathbf{x}) = \text{rank}_{\mathbf{k}} H_0(x_e; H_i(\mathbf{x}')) + \text{rank}_{\mathbf{k}} H_0(x_e; H_{i-1}(\mathbf{x}')).$$

Proof. Since \mathbf{x}' generates an \mathfrak{m} -primary ideal, the homology module $H_i(\mathbf{x}')$ has finite length for every i , and by depth sensitivity and rigidity it is non-zero if and only if $0 \leq i \leq e - d - 1$; see [24, prop. IV.3]. For every i and j the module $H_j(x_e; H_i(\mathbf{x}'))$ is annihilated by \mathfrak{m} and hence it is a finite dimensional \mathbf{k} -vector space. By Nakayama's lemma, $H_0(x_e; H_i(\mathbf{x}'))$ is non-zero if and only if $H_i(\mathbf{x}')$ is so.

For every i , there is a short exact sequence of finite dimensional \mathbf{k} -vector spaces

$$0 \rightarrow H_0(x_e, H_i(\mathbf{x}')) \rightarrow H_i(\mathbf{x}) \rightarrow H_1(x_e, H_{i-1}(\mathbf{x}')) \rightarrow 0;$$

see [24, prop. IV.1]. It yields

$$\text{rank}_{\mathbf{k}} H_i(\mathbf{x}) = \text{rank}_{\mathbf{k}} H_0(x_e; H_i(\mathbf{x}')) + \text{rank}_{\mathbf{k}} H_1(x_e; H_{i-1}(\mathbf{x}')).$$

To finish the proof we need to verify the equality

$$\text{rank}_{\mathbf{k}} H_1(x_e, H_{i-1}(\mathbf{x}')) = \text{rank}_{\mathbf{k}} H_0(x_e, H_{i-1}(\mathbf{x}')).$$

It follows from a length count in the exact sequence

$$0 \rightarrow H_1(x_e, H_{i-1}(\mathbf{x}')) \rightarrow H_{i-1}(\mathbf{x}') \xrightarrow{x_e} H_{i-1}(\mathbf{x}') \rightarrow H_0(x_e, H_{i-1}(\mathbf{x}')) \rightarrow 0. \quad \square$$

(2.3) **Lemma.** *Let R be singular and let $\mathbf{x} = x_1, \dots, x_e$ be a minimal system of generators of \mathfrak{m} such that $\mathbf{x}' = x_1, \dots, x_{e-1}$ generates an \mathfrak{m} -primary ideal. For $i \geq 0$ set $h_i = \text{rank}_{\mathbf{k}} H_0(x_e, H_i(\mathbf{x}'))$. Then there is a coefficient-wise inequality*

$$I_R(t) \preceq \frac{\sum_{i=0}^{e-d-1} h_{e-d-1-i} t^{d+i} - t^e}{1 - \sum_{i=0}^{e-d-1} h_i t^{i+1}},$$

and equality holds if and only if R is Golod.

Proof. Since R is singular, we have $e - d \geq 1$. For $i \geq 0$ set $c_i = \text{rank}_{\mathbf{k}} H_i(\mathbf{x})$, then (2.1.1) takes the form

$$I_R(t) \preceq \frac{\sum_{i=0}^{e-1} c_{e-i} t^i - t^{e+1}}{1 - \sum_{i=1}^e c_i t^{i+1}}.$$

We first verify that -1 is a root in both numerator and denominator. Indeed, by Lemma (2.2) there are equalities $c_{e-i} = h_{e-i} + h_{e-i-1}$; in particular, $h_0 = c_0 = 1$. Now we have

$$\begin{aligned} \sum_{i=0}^{e-1} c_{e-i} (-1)^i - (-1)^{e+1} &= \sum_{i=d+1}^{e-1} (h_{e-i} + h_{e-i-1}) (-1)^i - (-1)^{e+1} \\ &= h_0 (-1)^{e-1} - (-1)^{e+1} = 0 \end{aligned}$$

and

$$1 - \sum_{i=1}^e c_i (-1)^{i+1} = 1 - \sum_{i=1}^{e-d} (h_i + h_{i-1}) (-1)^{i+1} = 1 - h_0 (-1)^2 = 0.$$

Cancellation of the common factor $1 + t$ gives the equality

$$\frac{\sum_{i=0}^{e-1} c_{e-i} t^i - t^{e+1}}{1 - \sum_{i=1}^e c_i t^{i+1}} = \frac{\sum_{i=0}^{e-d-1} h_{e-d-1-i} t^{d+i} - t^e}{1 - \sum_{i=0}^{e-d-1} h_i t^{i+1}},$$

and the statement about Golodness follows from [4, (0.2)]. □

(2.4) **Observation.** Let R be Golod and assume it is not Gorenstein—that is, not a hypersurface—then one has $e - d \geq 2$; see [2, 5.1]. Let h_i for $i \geq 0$ be as defined in Lemma (2.3). As $h_0 = 1$ the Bass series of R takes the form

$$(2.4.1) \quad I_R(t) = \frac{\sum_{i=0}^{e-d-2} h_{e-d-1-i} t^{d+i} + t^{e-1} - t^e}{1 - t - \sum_{i=1}^{e-d-1} h_i t^{i+1}}.$$

Set $\mu^i = \mu^i(R)$ for $i \geq 0$. It is straightforward to deduce the next equalities from (2.4.1); one can also extract them from the proof of [4, (0.2)].

$$(2.4.2) \quad \mu^d = h_{e-d-1},$$

$$(2.4.3) \quad \mu^e = \mu^{e-1} + \sum_{i=0}^{e-d-2} \mu^{d+i} h_{e-d-1-i} - 1, \quad \text{and}$$

$$(2.4.4) \quad \mu^{e+n} = \mu^{e+n-1} + \sum_{i=n}^{n+e-d-2} \mu^{d+i} h_{e-d-1+n-i} \quad \text{for } n \geq 1.$$

The expression for the rate of growth A in the next theorem is inspired by Peeva's proof of [21, prop. 3]; see also [2, thm. 5.3.3.(5)]. In view of (1.0.1) it follows from the latter results that the Bass sequence for a Golod ring R with $e - d \geq 2$ has termwise exponential growth. The next theorem and Proposition (2.8) explains the initial behavior of these Bass sequences.

(2.5) **Theorem.** *Let R be Golod of depth d and embedding dimension e . If one of the inequalities $e - d > 2$ or $\mu^d(R) > 1$ holds, then the sequence $\{\mu^i(R)\}_{i \geq d}$ is increasing, and it has termwise exponential growth of rate*

$$A = \min \left\{ \frac{\mu^{d+1}(R)}{\mu^d(R)}, \frac{\mu^{d+2}(R)}{\mu^{d+1}(R)}, \dots, \frac{\mu^e(R)}{\mu^{e-1}(R)} \right\} > 1.$$

In the exceptional case with $e - d = 2$ and $\mu^d(R) = 1^*$, the Bass numbers of R are given by $\mu^{d+i}(R) = 2F_i$ for $i \geq 1$, where F_i is the i^{th} Fibonacci number. In particular the sequence $\{\mu^i(R)\}_{i \geq d}$ is non-decreasing and it has termwise exponential growth; see Proposition (2.8).

Proof. It follows from the assumptions on R that it is not a hypersurface, so we have $e - d \geq 2$. For $i \geq 0$ set $\mu^i = \mu^i(R)$ and adopt the notation from Lemma (2.3). There is a coefficient-wise inequality

$$\begin{aligned} (1-t)I_R(t) &\succcurlyeq \left(1 - t - \sum_{i=1}^{e-d-1} h_i t^{i+1}\right) I_R(t) \\ &= \sum_{i=0}^{e-d-2} h_{e-d-1-i} t^{d+i} + t^{e-1} - t^e, \end{aligned}$$

where the equality follows from (2.4.1). In particular, there are the following inequalities among the coefficients of $I_R(t)$:

$$\mu^d < \mu^{d+1} < \dots < \mu^{e-1}.$$

Moreover, at least one of the inequalities $\mu^d \geq 2$ or $e - d - 2 \geq 1$ holds, so (2.4.3) yields $\mu^{e-1} < \mu^e$ and, therefore, $A > 1$. By recursion it now follows from (2.4.4) and (2.4.3) that $\mu^{e+n} \geq A\mu^{e+n-1}$ for every $n \geq 1$. \square

* Let F be a field. The ring $R = F[[x, y]]/(x^2, xy)$ is an example. Indeed, it has Krull dimension 1 and depth 0, so it is Golod by [2, prop. 5.2.5], and $\text{Hom}_R(k, R)$ is generated by x , so $\mu^0(R) = 1$.

(2.6) **Corollary.** *If R is Golod of codimension at least 2, then the sequence of Bass numbers $\{\mu^i(R)\}_{i \geq d}$ is increasing and has termwise exponential growth.*

Proof. By assumption there is an inequality $e - d \geq 2$; in particular R is not a hypersurface and hence not Gorenstein. If equality holds, then R is Cohen–Macaulay, and then one has $\mu^d(R) > 1$ by [7, thm. 6.3]. The statement now follows from Theorem (2.5). \square

(2.7) **Remark.** This corollary covers Cohen–Macaulay rings of codimension 2 that are not Gorenstein. Indeed, such rings are Golod by [23]; see also [2, prop. 5.3.4]. In the next proposition we compute the Bass numbers of these rings in closed form.

(2.8) **Proposition.** *Let R be of depth d and embedding dimension $e = d + 2$; set $r = \mu^d(R)$. If R is not a complete intersection, then there is an equality*

$$I_R(t) = t^d \frac{r + t - t^2}{1 - t - rt^2}.$$

That is, the Bass numbers of R are

$$\mu^{d+i}(R) = \begin{cases} 0 & \text{for } i < 0 \\ r & \text{for } i = 0 \\ r + 1 & \text{for } i = 1 \\ r(r + 1) & \text{for } i = 2 \\ 2r(r + 1) & \text{for } i = 3 \\ \mu^{d+i-1} + r\mu^{d+i-2} & \text{for } i \geq 4. \end{cases}$$

Set $\delta = \sqrt{1 + 4r}$; for $i \geq 3$ the expression for $\mu^{d+i}(R)$ in closed form is then

$$\mu^{d+i}(R) = \frac{r+1}{2\delta} \left((2r-1+\delta) \left(\frac{1+\delta}{2} \right)^{i-1} + (1-2r+\delta) \left(\frac{1-\delta}{2} \right)^{i-1} \right).$$

Proof. By [23] R is Golod; see also [2, prop. 5.3.4]. The expression for the Bass series is, therefore, immediate from (2.4.1). For $i \geq 0$ set $\mu^i = \mu^i(R)$. A straightforward computation yields the expressions for μ^{d+1} and μ^{d+2} , and (2.4.4) yields the recurrence relation

$$\mu^{d+i} = \mu^{d+i-1} + r\mu^{d+i-2} \quad \text{for } i \geq 3.$$

The corresponding matrix $\begin{pmatrix} 0 & 1 \\ r & 1 \end{pmatrix}$ is diagonalizable with eigenvalues $\frac{1}{2}(1 \pm \sqrt{1 + 4r})$, and the expression for μ^{d+i} in closed form follows. \square

(2.9) **Remark.** Let R be as in Proposition (2.8). If $r = \mu^d(R)$ is 1, then one has $\mu^{d+1}(R) = 2 = \mu^{d+2}(R)$, and the recurrence relation yields $\mu^{d+i}(R) = 2F_i$ for $i \geq 1$, where F_i is the i^{th} Fibonacci number.

If $r \geq 2$, then the recurrence relation and the equality $\mu^{d+3}(R) = 2\mu^{d+2}(R)$ immediately yield $\mu^{i+1}(R) \geq 2\mu^i(R)$ for every $i \geq d + 3$, and equality holds if and only if $r = 2$. Thus, if $r = 2$, then one has $\mu^{d+i}(R) = 3(2^{i-1})$ for all $i \geq 1$.

3. FIBER PRODUCT RINGS

In this section, S and T are local rings with the same residue field k and both different from k . The fiber product $S \times_k T$ is a local ring with residue field k and embedding dimension $e = \text{edim } S + \text{edim } T$. We denote its depth by d .

We start by observing a few fiber product rings that fail to have increasing Bass numbers, because they are either hypersurfaces or of the type considered in Proposition (2.8). The main result of the section—Theorem (3.4)—is that they are the only (non-trivial) ones. This will establish part (2) of the Main Theorem (1.2).

(3.1) Let M be a finitely generated R -module. Recall that the *Poincaré series* of M is the formal power series

$$P_M^R(t) = \sum_{i=0}^{\infty} \beta_i^R(M) t^i.$$

The Poincaré series of k as an $S \times_k T$ -module was first computed by Kostrikin and Shafarevich [16]:

$$(3.1.1) \quad \frac{1}{P_k^{S \times_k T}(t)} = \frac{1}{P_k^S(t)} + \frac{1}{P_k^T(t)} - 1.$$

Lescot computes the quotient of the Bass series and the Poincaré series of $S \times_k T$ in [17, thm. 3.1]. For later reference we record some details from this work. If S and T are both singular, then one has

$$(3.1.2) \quad \frac{I_{S \times_k T}(t)}{P_k^{S \times_k T}(t)} = t + \frac{I_S(t)}{P_k^S(t)} + \frac{I_T(t)}{P_k^T(t)}.$$

If S is singular and T is regular of dimension n , then the formula is

$$(3.1.3) \quad \frac{I_{S \times_k T}(t)}{P_k^{S \times_k T}(t)} = t + \frac{I_S(t)}{P_k^S(t)} - \frac{t^{n+1}}{(1+t)^n}.$$

If S and T are regular of dimension m and n , then one has

$$(3.1.4) \quad \frac{I_{S \times_k T}(t)}{P_k^{S \times_k T}(t)} = t - \frac{t^{m+1}}{(1+t)^m} - \frac{t^{n+1}}{(1+t)^n}.$$

The *order* of a power series $\sum_{i=0}^{\infty} v_i t^i$ is the number $\min\{i \geq 0 \mid v_i \neq 0\}$. Note that the order of the Bass series $I_R(t)$ is equal to $\text{depth } R$.

(3.2) **Remark.** It follows from (3.1.2)–(3.1.4) that the depth of $S \times_k T$ is at most 1, and that it is 0 if either S or T has depth 0. That is, one has

$$(3.2.1) \quad d = \min\{\text{depth } S, \text{depth } T, 1\}.$$

(3.3) **Observation.** Let S be a 1-dimensional regular ring. If also T is regular of dimension 1, then one has $e = 2$ and $d = 1$ by (3.2.1), so $S \times_k T$ is a hypersurface.

If T is either a 0-dimensional hypersurface or a 2-dimensional regular ring, then one has $e - d = 2$ by (3.2.1). Moreover, $S \times_k T$ is Golod as both S and T are Golod; see [18, thm. 4.1]. Finally, it follows from (3.1.3) and (3.1.4) that $\mu^d(S \times_k T)$ is 1 in either case, and then one has $\mu^{d+1}(S \times_k T) = 2 = \mu^{d+2}(S \times_k T)$ as worked out in Remark (2.9).

(3.4) **Theorem.** *Let S and T be local rings with common residue field k and assume that both S and T are different from k . If the fiber product ring $S \times_k T$ is not one of the three types from (3.3), then the sequence of Bass numbers $\{\mu^i(S \times_k T)\}_{i \geq d}$ is increasing and has exponential growth.*

Notice that part (2) of the Main Theorem, stated in the Introduction, follows from Theorem (3.4), as the rings in Observation (3.3) are Golod.

For the proof of Theorem (3.4) we need some terminology and a technical lemma. A power series $\sum_{i=0}^{\infty} v_i t^i$ of order n is said to have non-negative (or positive) coefficients if $v_i \geq 0$ (or $v_i > 0$) for all $i \geq n$; it has non-decreasing (or increasing) coefficients if $v_{i+1} \geq v_i$ (or $v_{i+1} > v_i$) for all $i \geq n$.

(3.5) **Lemma.** *Let $\sum_{i=0}^{\infty} c_i t^i$ be a formal power series with increasing coefficients and assume $c_0 > 1$. Then the following hold:*

(a) *The coefficients of the power series*

$$V(t) = \sum_{i=0}^{\infty} v_i t^i := \frac{1}{1 - t^2 \sum_{i=0}^{\infty} c_i t^i}$$

grow exponentially and satisfy:

$$v_0 = 1, \quad v_1 = 0, \quad v_2 = c_0, \quad \text{and} \quad v_{i+1} > v_i \quad \text{for all } i \geq 2.$$

(b) *Let $W(t) = \sum_{i=0}^{\infty} w_i t^i$ be a power series of order 0 and assume that the series $(1 - t + t^2)W(t)$ has non-negative coefficients. Then the series $(1 - t)V(t)W(t)$ has order 0 and positive coefficients in each degree except, possibly, in degree 1 where the coefficient is $w_1 - w_0 \geq 0$.*

(3.6) **Remark.** If R has positive embedding dimension, then the Poincaré series $P_k^R(t)$ satisfies the condition on $W(t)$ in Lemma (3.5)(b). Indeed, $P_k^R(t)$ is either a power series with non-decreasing coefficients or the polynomial $(1 + t)^n$ for some $n \geq 1$. In the first case the claim is obvious, and in the second case it follows from the inequality $\binom{n}{i} \leq \binom{n}{i-1} + \binom{n}{i+1}$, which holds for all integers i .

Proof of Lemma (3.5). (a): The equality $(1 - t^2 \sum_{i=0}^{\infty} c_i t^i)(\sum_{i=0}^{\infty} v_i t^i) = 1$ immediately yields $v_0 = 1$ and $v_1 = 0$, and it yields $v_2 - c_0 v_0 = 0$ whence $v_2 = c_0$. For $i \geq 3$ it yields $v_i - c_{i-2} - \sum_{j=0}^{i-3} c_j v_{i-2-j} = 0$, and it follows by recursion that the coefficients v_i are positive. Now the desired inequalities

$$v_{i+1} = c_{i-1} + \sum_{j=0}^{i-2} c_j v_{i-1-j} > c_{i-2} + \sum_{j=0}^{i-3} c_j v_{i-2-j} = v_i \quad \text{for } i \geq 2$$

follow as the sequence $\{c_i\}_{i \geq 0}$ is increasing by assumption. Finally, the expression for v_i yields an inequality $v_{i+2} > c_0 v_i$ for each $i \geq 2$. In particular, we have $v_{2j} > c_0^j$ and $v_{2j-1} > c_0^{j-1} c_1$ for $j \geq 2$. As $c_1 > c_0 > 1$ we now have $v_i > \sqrt{c_0}^i$ for $i \geq 3$, so the sequence $\{v_i\}_{i \geq 0}$ has exponential growth.

(b): The first equality in the computation below holds as $v_1 = 0$.

$$\begin{aligned} (1-t)V(t)W(t) &= (1-t) \left(1 + \sum_{i=2}^{\infty} v_i t^i\right) W(t) \\ &= (1-t)W(t) + t^2 W(t) (1-t) \sum_{i=2}^{\infty} v_i t^{i-2} \\ &= (1-t+t^2)W(t) + t^2 W(t) \left((1-t) \sum_{i=2}^{\infty} v_i t^{i-2} - 1 \right) \end{aligned}$$

By the assumptions on $W(t)$, the first summand in the last line of the display, $(1-t+t^2)W(t)$, has order 0 and non-negative coefficients; the coefficient in degree 1 is $w_1 - w_0$. It follows from the inequality $v_2 > 1$ that the second summand has order 2, and it has positive coefficients as $v_{i+1} > v_i$ for all $i \geq 2$. \square

Proof of Theorem (3.4). Set $R = S \times_k T$. Assume first that R is Golod; then both S and T are Golod by [18, thm. 4.1]. From (3.2.1) and the equality $e = \text{edim } S + \text{edim } T$, it is straightforward to verify that the (in)equalities $e - d \leq 2$ and $\mu^d(R) = 1$ hold exactly when R is one of the three types of rings from (3.3). If $e - d > 2$ or $\mu^d(R) > 1$, then it is proved in Theorem (2.5) that the sequence $\{\mu^i(R)\}_{i \geq d}$ is increasing and has exponential growth.

Assume now that R is not Golod. Without loss of generality we may assume that S is not a hypersurface; cf. [18, thm. 4.1]. Rewrite (3.1.1) as

$$P_k^R(t) = \frac{P_k^S(t) P_k^T(t)}{1 - (P_k^S(t) - 1)(P_k^T(t) - 1)}.$$

Combine it with (3.1.2) to obtain

$$(3.6.1) \quad I_R(t) = \frac{I_S(t) P_k^T(t) + I_T(t) P_k^S(t) + t P_k^S(t) P_k^T(t)}{1 - (P_k^S(t) - 1)(P_k^T(t) - 1)}$$

in case T is singular. And in case T is regular, combine it with (3.1.3) to obtain

$$(3.6.2) \quad I_R(t) = \frac{I_S(t)(1+t)^n - t^{n+1} P_k^S(t) + t(1+t)^n P_k^S(t)}{1 - (P_k^S(t) - 1)(P_k^T(t) - 1)},$$

where $n \geq 1$ is the dimension of T . As S is not a hypersurface, its Poincaré series has increasing coefficients, so the power series $(1-t)(P_k^S(t) - 1)$ has positive coefficients, and hence so has $(1-t)(P_k^S(t) - 1)(P_k^T(t) - 1)$. That is, the series $(P_k^S(t) - 1)(P_k^T(t) - 1)$ of order 2 has increasing coefficients. Moreover, the degree 2 coefficient $(\text{edim } S)(\text{edim } T)$ is greater than 1 as S is not a hypersurface. From Lemma (3.5)(a) it now follows that the series $1/[1 - (P_k^S(t) - 1)(P_k^T(t) - 1)]$ has exponential growth. In (3.6.1) as well as in (3.6.2) the numerator has non-negative coefficients, so in either case the sequence $\{\mu^i(R)\}_{i \geq d}$ has exponential growth.

If T is singular, then the series $(1-t)P_k^T(t)$ has non-negative coefficients, and as above $(1-t)P_k^S(t)$ has positive coefficients. Thus, the numerator in the expression

$$(1-t)I_R(t) = \frac{(1-t)(I_S(t) P_k^T(t) + I_T(t) P_k^S(t) + t P_k^S(t) P_k^T(t))}{1 - (P_k^S(t) - 1)(P_k^T(t) - 1)},$$

which is derived from (3.6.1), is a power series of order $d \leq 1$ with positive coefficients. As the power series $1/[1 - (P_k^S(t) - 1)(P_k^T(t) - 1)]$ has non-negative coefficients

and order 0, it follows that $(1-t)I_R(t)$ has positive coefficients in all degrees $i \geq d$. That is, the sequence $\{\mu^i(R)\}_{i \geq d}$ is increasing.

Finally, assume that T is regular of dimension n . The Poincaré series $(1+t)^n$ and $P_k^S(t)$ satisfy the condition on $W(t)$ in Lemma (3.5)(b); see also Remark (3.6). By (3.6.2) the series $(1-t)I_R(t)$ of order $d \leq 1$ can be expressed as a sum

$$(1-t)I_R(t) = (1-t) \frac{(1+t)^n}{1 - (P_k^S(t) - 1)(P_k^T(t) - 1)} I_S(t) + (1-t) \frac{P_k^S(t)}{1 - (P_k^S(t) - 1)(P_k^T(t) - 1)} (t(1+t)^n - t^{n+1}).$$

The first summand has order depth $S \geq d$, and by Lemma (3.5)(b) it has non-negative coefficients. Similarly, the second summand has order 1 and positive coefficients, also in degree 2 as $\beta_1^S(k) - \beta_0^S(k) = \text{edim } S - 1 > 0$. It follows that the power series $(1-t)I_R(t)$ has positive coefficients in all degrees $i \geq d$, therefore, the sequence $\{\mu^i(R)\}_{i \geq d}$ is increasing. \square

The last result of this section is part (5) of the Main Theorem. It is proved in Remark (4.6) and Proposition (A.3), but we state it here because it is concerned with fiber product rings.

(3.7) Remark. Let R be artinian, and assume that k splits out of the maximal ideal \mathfrak{m} —that is, $\text{Soc } R$ contains a minimal generator of \mathfrak{m} . If R is not a hypersurface, i.e. $\text{edim } R$ is at least 2, then it is a non-trivial fiber product of artinian local rings. Indeed, let x, y_1, \dots, y_m be a minimal set of generators of \mathfrak{m} , such that x is in $\text{Soc } R$, then there is an isomorphism $R \cong R/(x) \times_k R/(y_1, \dots, y_m)$. By Theorem (3.4) the sequence $\{\mu^i(R)\}_{i \geq 0}$ is increasing and has exponential growth.

One can, however, do better for this particular kind of fiber products. Let R be as in (3.7). In case $\mathfrak{m}^2 = 0$ the sequence $\{\mu^i(R)\}_{i \geq 0}$ is increasing and has even termwise exponential; see Example (1.5). In case $\mathfrak{m}^2 \neq 0$ the same conclusion follows from Proposition (A.3) and Remark (4.6). Thus, we have the following:

(3.8) Proposition. *Let R be an artinian local ring of embedding dimension at least 2. If $\text{Soc } R \not\subseteq \mathfrak{m}^2$, then the sequence of Bass numbers $\{\mu^i(R)\}_{i \geq 0}$ is increasing and it has termwise exponential growth.* \square

4. RINGS WITH $\mathfrak{m}^3 = 0$.

Theorem (4.7) at the end of this section establishes part (3) of the Main Theorem from the Introduction. The assumption $\mathfrak{m}^3 = 0$ is only needed in (4.7). Until then, (R, \mathfrak{m}, k) is just assumed to be artinian; the injective envelope of k is denoted $E_R(k)$.

(4.1) Let M be a finitely generated R -module. We write $\ell_R(M)$ for the length of M and $\text{Soc } M$ for its socle. It is straightforward to verify the following:

$$(4.1.1) \quad k \text{ is a direct summand of } M \text{ if and only if } \text{Soc } M \not\subseteq \mathfrak{m}M.$$

For $i \geq 1$ we denote the i^{th} syzygy in a minimal free resolution of M by M_i ; we set $M_0 = M$. Since the differentials in a minimal free resolution are given by matrices with entries in \mathfrak{m} , there are equalities

$$(4.1.2) \quad \text{Soc } M_{i+1} = \text{Soc } R^{\beta_i^R(M)} \quad \text{for all } i \geq 1.$$

(4.2) Recall that one has

$$(4.2.1) \quad \mu^i(R) = \beta_i^R(\mathbf{E}_R(\mathbf{k})) \quad \text{for all } i \geq 0.$$

If R is not Gorenstein, then the equality $\ell_R(\mathbf{E}_R(\mathbf{k})) = \ell(R)$ and Lemma (A.1) yield

$$(4.2.2) \quad \mu^1(R) \geq \mu^0(R).$$

Recall also that one has $\text{rank}_{\mathbf{k}} \text{Soc } R = \mu^0(R)$, and denote this number by r .

The first preparatory result contains a special case of Theorem (4.7), namely the one where the socle rank r exceeds the embedding dimension.

(4.3) **Proposition.** *Set $h = \max\{i \mid \mathfrak{m}^i \neq 0\}$ and assume that R satisfies $h \geq 2$ and $r > \ell_R(R/\mathfrak{m}^h) - 1$. If R is not Gorenstein, then the sequence of Bass numbers $\{\mu^i(R)\}_{i \geq 0}$ is increasing and has termwise exponential growth of rate at least*

$$\frac{r}{\ell_R(R/\mathfrak{m}^h) - 1}.$$

Proof. By Proposition (A.2) we need only prove the inequality $\mu^1(R) > \mu^0(R)$, and to this end it suffices, by (4.2.2), to show $\mu^1(R) \neq r$. Assume, towards a contradiction, that one has $\mu^1(R) = r$. Set $E = \mathbf{E}_R(\mathbf{k})$ and consider the exact sequence

$$0 \rightarrow E_2 \rightarrow R^r \rightarrow R^r \rightarrow E \rightarrow 0,$$

that comes from the minimal free resolution of E . By (4.1.2) and additivity of length, the sequence yields (in)equalities

$$r^2 = \text{rank}_{\mathbf{k}} \text{Soc } R^r = \text{rank}_{\mathbf{k}} \text{Soc } E_2 \leq \ell_R(E_2) = \ell_R(E).$$

By additivity of length, the assumptions on R , and the containment $\mathfrak{m}^h \subseteq \text{Soc } R$ there are (in)equalities

$$\ell(R) = \ell_R(R/\mathfrak{m}^h) + \ell_R(\mathfrak{m}^h) < 1 + r + \text{rank}_{\mathbf{k}} \mathfrak{m}^h \leq 1 + 2r.$$

By the equality $\ell_R(E) = \ell(R)$, the last two displays combine to yield $r^2 \leq 1 + 2r$, which implies $r = 1$ contradicting the assumption that R is not Gorenstein. \square

(4.4) **Lemma.** *If \mathbf{k} is a direct summand of the first syzygy of $\mathbf{E}_R(\mathbf{k})$, then one has $\mu^1(R) > \mu^0(R)$.*

Proof. By (4.2.2) it is enough to rule out the possibility of an equality $\mu^1(R) = r$. Suppose this equality holds. For brevity, set $E = \mathbf{E}_R(\mathbf{k})$. By hypothesis, there is an isomorphism $E_1 \cong \mathbf{k} \oplus N$ for some finitely generated R -module N and, therefore, there is an isomorphism of syzygies $E_2 \cong \mathfrak{m} \oplus N_1$. From the exact sequence $0 \rightarrow E_2 \rightarrow R^r \rightarrow R^r \rightarrow E \rightarrow 0$ one obtains $\ell_R(\mathfrak{m}) + \ell_R(N_1) = \ell_R(E)$ by additivity of length. Since $\ell_R(E) = \ell_R(R) = \ell_R(\mathfrak{m}) + 1$, it follows that N_1 is a \mathbf{k} -vector space of rank 1. Hence there is an exact sequence $0 \rightarrow \mathbf{k} \rightarrow R^{r-1} \rightarrow N \rightarrow 0$, where the surjective homomorphism is given by a matrix with entries in \mathfrak{m} , as N is a submodule of $\mathfrak{m}R^r$. Thus, we have $\text{Soc } R^{r-1} = \mathbf{k}$, and the ensuing equality of ranks $r(r-1) = 1$ is absurd. \square

(4.5) **Lemma.** *Assume that R is not Gorenstein, and set $E = \mathbf{E}_R(\mathbf{k})$. Let n be an integer such that $\text{Soc } E_1$ is contained in $\mathfrak{m}^n E_1$. Then one has $n < \max\{i \mid \mathfrak{m}^i \neq 0\}$ and $\text{Soc } R \subseteq \mathfrak{m}^{n+1}$.*

Proof. Set $h = \max\{i \mid \mathfrak{m}^i \neq 0\}$, then $\mathfrak{m}^h E_1 = 0$ as E_1 is a submodule of $\mathfrak{m}R^r$; and the inequality $n < h$ follows.

Let e_1, \dots, e_r be the standard basis for R^r . Choose a minimal set $\varepsilon_1, \dots, \varepsilon_r$ of generators of E and consider the short exact sequence $0 \rightarrow E_1 \rightarrow R^r \rightarrow E \rightarrow 0$, where the surjection maps e_i to ε_i . Suppose that $\text{Soc } R$ is not contained in \mathfrak{m}^{n+1} and choose an element x in $(\text{Soc } R) \setminus \mathfrak{m}^{n+1}$. Since E is a faithful R -module, there is an index i such that the element $x\varepsilon_i$ is non-zero; it clearly belongs to $\text{Soc } E$. Similarly, by the definition of h , it follows that $\mathfrak{m}^h E$ is a non-zero submodule of $\text{Soc } E$. Since $\text{rank}_{\mathfrak{k}} \text{Soc } E = 1$, there exists an element y in \mathfrak{m}^h and an index j such that $y\varepsilon_j = x\varepsilon_i$. The element $z = xe_i - ye_j$ is in the socle of E_1 , as x and y are in the socle of R . As y is in \mathfrak{m}^{n+1} but x is not, the element z is not in $\mathfrak{m}^n E_1 \subseteq \mathfrak{m}^{n+1} R^r$, which contradicts the assumption that $\text{Soc } E_1$ is contained in $\mathfrak{m}^n E_1$. \square

(4.6) **Remark.** Set $E = E_R(\mathfrak{k})$. If \mathfrak{k} is a direct summand of \mathfrak{m} , that is $\text{Soc } R \not\subseteq \mathfrak{m}^2$, then it follows from Lemma (4.5) that $\text{Soc } E_1 \not\subseteq \mathfrak{m}E_1$. Thus, \mathfrak{k} is a direct summand of E_1 , cf. (4.1.1), and then Lemma (4.4) yields the inequality $\mu^1(R) > \mu^0(R)$.

(4.7) **Theorem.** *Let (R, \mathfrak{m}) be local with $\mathfrak{m}^3 = 0$. If R is not Gorenstein, then the Bass sequence $\{\mu^i(R)\}_{i \geq 0}$ is increasing and has termwise exponential growth.*

Proof. In view of Example (1.5) we can assume that \mathfrak{m}^2 is not 0. Set

$$a = \text{rank}_{\mathfrak{k}} \mathfrak{m}^2, \quad e = \text{rank}_{\mathfrak{k}} \mathfrak{m}/\mathfrak{m}^2, \quad \text{and} \quad r = \text{rank}_{\mathfrak{k}} \text{Soc } R.$$

The embedding dimension e is at least 2, as R is not a hypersurface, and for $e = 2$ the statement is contained in Corollary (2.6), cf. Remark (2.7). In the rest of the proof, we assume $e \geq 3$. Set $E = E_R(\mathfrak{k})$. For $i \geq 0$ set $\mu^i = \mu^i(R)$ and recall from (4.2.1) that $\mu^i = \beta_i^R(E)$. The containment $\mathfrak{m}^2 \subseteq \text{Soc } R$ implies an inequality $a \leq r$.

If $a < r$, then $\text{Soc } R$ is not contained in \mathfrak{m}^2 and Remark (4.6) gives the inequality $\mu^1 > \mu^0$. Then Theorem (A.5)(a) applies to the module E , so the sequence $\{\mu^i\}_{i \geq 0}$ is increasing and it has termwise exponential growth.

In the remainder of the proof assume $a = r$. From the equalities $\ell_R(E) = \ell(R)$ and $1 = \text{rank}_{\mathfrak{k}} \text{Soc } E = \text{rank}_{\mathfrak{k}} \mathfrak{m}^2 E$ one gets $\text{rank}_{\mathfrak{k}} \mathfrak{m}E/\mathfrak{m}^2 E = \ell(R) - r - 1 = e$. By [19, lem. 3.3]—see also (A.4.2)—there is an inequality

$$(1) \quad \mu^1 \geq e\mu^0 - \text{rank}_{\mathfrak{k}} \mathfrak{m}E/\mathfrak{m}^2 E = e(r-1),$$

and equality holds if and only if \mathfrak{k} is not a direct summand of E_1 . Since $e \geq 3$ and $r \geq 2$, an equality $\mu^1 = \mu^0$ would imply $r = e(r-1) \geq r + 2r - 3 \geq r + 1$, which is absurd. Thus, the inequality $\mu^1 > \mu^0$ holds, cf. (4.2.2). Now Theorem (A.5)(a) applies to the module E , so the sequence $\{\mu^i\}_{i \geq 0}$ is increasing with termwise exponential growth except, possibly, when $a = e = r$.

Assume now that all three invariants a , e , and r are equal. Note that (1) yields

$$(2) \quad \mu^1 \geq (r-1)\mu^0.$$

By [19, lem. 3.3]—see also (A.4.3)—there are inequalities

$$(3) \quad \mu^{i+1} \geq r(\mu^i - \mu^{i-1}) \quad \text{for } i \geq 1$$

and equality holds if $i \geq 2$ and \mathfrak{k} is not a summand of either syzygy E_i and E_{i-1} .

Consider the case where the common value of a , e , and r is at least 4. The quantity $A = \frac{1}{2}(r + \sqrt{r^2 - 4r})$ is then a real number greater than or equal to 2. We

claim that the inequality $\mu^{i+1} \geq A\mu^i$ holds for all $i \geq 0$. The proof is by induction on i . The base case $i = 0$ is furnished by (2) and the induction step follows from (3):

$$\mu^{i+1} \geq r(\mu^i - \mu^{i-1}) \geq r(\mu^i - A^{-1}\mu^i) = r(1 - A^{-1})\mu^i = A\mu^i,$$

where the last equality follows as A is a solution to the equation $r(1 - x^{-1}) = x$.

As $e \geq 3$ we are left with only one case to consider, namely $a = e = r = 3$. First we prove that the sequence $\{\mu^i\}_{i \geq 0}$ is increasing and that \mathfrak{k} is a direct summand of one of the first four syzygies of E . If \mathfrak{k} is a direct summand of E_1 or E_2 , then the sequence $\{\mu^i\}_{i \geq 0}$ is increasing by Theorem (A.5)(b). Assume now that \mathfrak{k} is not a direct summand of E_1 , then (1) yields

$$\mu^1 = e(r - 1) = 6.$$

Assume also that \mathfrak{k} is not a direct summand of E_2 . In the computation

$$\text{rank}_{\mathfrak{k}} \mathfrak{m}E_2 = \text{rank}_{\mathfrak{k}} \text{Soc } E_2 = \text{rank}_{\mathfrak{k}} \text{Soc } R^6 = 18$$

the first equality follows from (4.1.1) and the second from (4.1.2). Combined with length computations in the minimal free resolution of E , this gives

$$\mu^2 = \ell_R(E_2) - \ell_R(\mathfrak{m}E_2) = 4\ell_R(R) - 18 = 10.$$

In particular, one has $\mu^2 > \mu^1$. If \mathfrak{k} is a direct summand of E_3 , then the sequence $\{\mu^i\}_{i \geq 0}$ is increasing by (A.5)(b). If \mathfrak{k} is not a direct summand of E_3 , then (3) yields

$$\mu^3 = 3(\mu^2 - \mu^1) = 12.$$

If \mathfrak{k} were not a direct summand of E_4 , then (3) would yield $\mu^4 = 3(\mu^3 - \mu^2) = 6 < \mu^3$ which is impossible by Theorem (A.5)(b). Thus, \mathfrak{k} is a direct summand of E_4 , and the sequence $\{\mu^i\}_{i \geq 0}$ is increasing by (A.5)(b) as one has $\mu^3 > \mu^2$.

Finally, we can conclude that the growth of the series $\{\mu^i\}_{i \geq 0}$ is termwise exponential. Since \mathfrak{k} is a direct summand of a syzygy of E , the radius of convergence of the power series $P_E^R(t) = I_R(t)$ is bounded above by that of $P_{\mathfrak{k}}^R(t)$. The opposite inequality always holds by [19, prop. 1.1], so the two power series have the same radius of convergence ρ . As R is not complete intersection, the sequence $\{\beta_i^R(\mathfrak{k})\}_{i \geq 0}$ has exponential growth, so one has $\rho < 1$. From work of Sun [25, thm. 1.2] it now follows that the sequence $\{\mu^i\}_{i \geq 0}$ has termwise exponential growth. \square

(4.8) **Scholium.** Let R be as in Theorem (4.7). The table below gives a lower bound for the rate A of the termwise exponential growth of the Bass sequence $\{\mu^i(R)\}_{i \geq 0}$ in terms of the invariants $a = \text{rank}_{\mathfrak{k}} \mathfrak{m}^2$, $e = \text{rank}_{\mathfrak{k}} \mathfrak{m}/\mathfrak{m}^2$, and $r = \text{rank}_{\mathfrak{k}} \text{Soc } R$.

a	$a = 0$	$a < r$	$a = r$				
			$e < r$	$e = r$		$e = r + 1$	$e \geq r + 2$
r				$2 \leq r \leq 3$	$4 \leq r$		
A	$e = r$	$r - a + \frac{r - a}{a + e}$	$\frac{r}{e}$	2	$\frac{r + \sqrt{r^2 - 4r}}{2}$	$\frac{r^2 - 1}{r}$	$e - r$

For rings with $a = e = r = 3$, Backelin and Fröberg [6] give closed form expressions for the possible Poincaré series $P_{\mathfrak{k}}^R(t)$, and one can verify directly that the radius of convergence is less than $\frac{1}{2}$. Sun's [25, thm. 1.2] then yields the lower bound 2 for A . In all other cases, the bound follows by inspection of Example (1.4), the proof of Theorem (A.5)(a), the proof of Theorem (4.7), and Peeva's proof of [21, prop. 3].

5. TETER RINGS

Following Huneke and Vraciu [14], we say that R is *Teter* if there exists an artinian Gorenstein local ring Q such that $R \cong Q/\text{Soc } Q$.

(5.1) Let Q be an artinian Gorenstein local ring with residue field k and set $R = Q/\text{Soc } Q$. The maximal ideal of Q is isomorphic to $\text{Hom}_Q(R, Q)$, which is the injective hull of k as an R -module. Thus, R is Gorenstein if and only if $\text{edim } Q = 1$, in which case both Q and R are hypersurfaces.

If Q is not a hypersurface, i.e. $\text{edim } Q \geq 2$, then one has $\text{edim } R = \text{edim } Q$ and works of Avramov and Levin [20, thm. 2.9] and Herzog and Steurich [13, prop. 1] provide the following expression for the Bass series of R :

$$(5.1.1) \quad I_R(t) = \frac{P_k^Q(t) - 1}{t(1 - t^2 P_k^Q(t))}.$$

Notice the equality $\mu^0(R) = \text{edim } R$.

(5.2) For each positive integer e set

$$\Upsilon_i^e = \binom{e}{i}^{-1} \left(\sum_{j=0}^{i+1} \binom{e-1}{j} \right) \quad \text{for } i \in \{0, \dots, e\} \quad \text{and}$$

$$\Upsilon^e = \min\{\Upsilon_i^e \mid 0 \leq i \leq e\}.$$

Notice that $\Upsilon_0^e = e$ and $\Upsilon_e^e = 2^{e-1}$ for every $e \geq 1$. Moreover, one has $\Upsilon_1^2 = 1$, and for $e \geq 3$ there are inequalities

$$\Upsilon_1^e \leq \frac{e}{2}$$

$$1 < 1 + \binom{e}{i}^{-1} \binom{e-1}{i+1} \leq \Upsilon_i^e \quad \text{for } 1 \leq i \leq \lfloor \frac{e}{2} \rfloor \quad \text{and}$$

$$\Upsilon_i^e \leq \Upsilon_{i+1}^e \quad \text{for } \lfloor \frac{e}{2} \rfloor \leq i \leq e-1.$$

In particular, one has

$$(5.2.1) \quad \Upsilon^1 = 1 = \Upsilon^2 \quad \text{and} \quad 1 < \Upsilon^e = \min\{\Upsilon_i^e \mid 1 \leq i \leq \lfloor \frac{e}{2} \rfloor\} \quad \text{for } e \geq 3.$$

Direct computations yield $\Upsilon^e = \Upsilon_{\lfloor \frac{e}{2} \rfloor}^e$ for $e \leq 9$ but $\Upsilon^{10} = \Upsilon_4^{10}$. To find Υ^e for larger values of e it is useful to know that Υ_i^e is an upwards convex function of i ; Roger W. Barnard proved this upon request. Thus, if n is an integer between 1 and $\lfloor \frac{e}{2} \rfloor$ and there are inequalities $\Upsilon_{n-1}^e \geq \Upsilon_n^e \leq \Upsilon_{n+1}^e$, then one has $\Upsilon^e = \Upsilon_n^e$.

(5.3) **Lemma.** *Let R be artinian of embedding dimension $e \geq 1$. For every real number $A < \Upsilon^e$ the formal power series $(1 - At + At^3) P_k^R(t)$ has positive coefficients, and for $A = \Upsilon^e$ it has non-negative coefficients.*

Proof. By [2, 7.1] there are equalities

$$P_k^R(t) = \frac{\prod_{i=1}^{i=\infty} (1 + t^{2i-1})^{\varepsilon_{2i-1}}}{\prod_{i=1}^{i=\infty} (1 - t^{2i})^{\varepsilon_{2i}}} = \frac{(1+t)^{\varepsilon_1}}{1-t^2} \cdot \frac{\prod_{i=2}^{i=\infty} (1 + t^{2i-1})^{\varepsilon_{2i-1}}}{(1-t^2)^{\varepsilon_2-1} \prod_{i=2}^{i=\infty} (1 - t^{2i})^{\varepsilon_{2i}}},$$

and one has $\varepsilon_2 - 1 \geq 0$ as R is not regular; see [2, thm. 7.3.2]. By [2, cor. 7.1.5] there is an equality $\varepsilon_1 = e$, so the factor $F = \frac{(1+t)^{\varepsilon_1}}{1-t^2}$ can be rewritten as

$$F = \frac{(1+t)^{e-1}}{1-t} = (1+t)^{e-1} \sum_{i=0}^{\infty} t^i = \sum_{i=0}^{e-2} \left(\sum_{j=0}^i \binom{e-1}{j} \right) t^i + \sum_{i=e-1}^{\infty} 2^{e-1} t^i.$$

The power series $(1 - At + At^3) P_k^R(t)$ has positive/non-negative coefficients if the series $(1 - At + At^3)F$ has positive/non-negative coefficients. From the computation

$$\begin{aligned} (1 - At + At^3)F &= \sum_{i=0}^{e+1} \left(\sum_{j=0}^i \binom{e-1}{j} - A \left(\binom{e-1}{i-2} + \binom{e-1}{i-1} \right) \right) t^i + \sum_{i=e+2}^{\infty} 2^{e-1} t^i \\ &= 1 + \sum_{i=1}^{e+1} \left(\sum_{j=0}^i \binom{e-1}{j} - A \binom{e}{i-1} \right) t^i + \sum_{i=e+2}^{\infty} 2^{e-1} t^i \\ &= 1 + \sum_{i=0}^e \left(\sum_{j=0}^{i+1} \binom{e-1}{j} - A \binom{e}{i} \right) t^{i+1} + \sum_{i=e+2}^{\infty} 2^{e-1} t^i \end{aligned}$$

it follows that $(1 - At + At^3)F$ has non-negative coefficients if and only if $A \leq \Upsilon^e$ and positive coefficients if $A < \Upsilon^e$. \square

The next result establishes part (4) of the Main Theorem (1.2).

(5.4) **Theorem.** *Let R be Teter of embedding dimension e at least 2.*

- (a) *If $e = 2$, then $\mu^0(R) = 2$ and $\mu^i(R) = 3(2^{i-1})$ for $i \geq 1$.*
- (b) *If $e > 2$, then $\Upsilon^e > 1$ and the formal power series $(1 - \Upsilon^e t) I_R(t)$ has non-negative coefficients.*

In particular, the sequence of Bass numbers $\{\mu^i(R)\}_{i \geq 0}$ is increasing and has term-wise exponential growth.

Proof. If $e = 2$, then R is Golod; see [2, prop. 5.3.4]. Since R is Teter, it follows from (5.1.1) that $\mu^0(R) = e = 2$. Thus, part (a) follows from Remark (2.9).

Assume now that $e > 2$, then $\Upsilon^e > 1$ by (5.2.1). To prove part (b), it is sufficient to show that the series $(1 - \Upsilon^e t)(1 + t I_R(t))$ has non-negative coefficients. To this end, let Q be an artinian Gorenstein ring such that $R \cong Q/\text{Soc } Q$. From (5.1.1) one obtains

$$(1) \quad 1 + t I_R(t) = (1 + t^3 I_R(t)) P_k^Q(t) = (1 - t^2 + t^2(1 + t I_R(t))) P_k^Q(t).$$

Set $1 + t I_R(t) = \sum_{i=0}^{\infty} a_i t^i$ and $P_k^Q(t) = \sum_{i=0}^{\infty} b_i t^i$. Now (1) yields relations

$$\begin{aligned} a_0 &= b_0 = 1, \\ a_1 &= b_1, \\ a_2 &= b_2 - b_0 + b_0 a_0 = b_2, \end{aligned}$$

and for $n \geq 2$:

$$a_n = b_n - b_{n-2} + \sum_{i=0}^{n-2} b_i a_{n-2-i}.$$

The coefficients of the series $(1 - \Upsilon^e t)(1 + tI_R(t))$ may now be expressed as

$$\begin{aligned} a_1 - \Upsilon^e a_0 &= b_1 - \Upsilon^e b_0, \\ a_2 - \Upsilon^e a_1 &= b_2 - \Upsilon^e b_1, \\ a_3 - \Upsilon^e a_2 &= b_3 - \Upsilon^e b_2 + \Upsilon^e b_0 + b_0(a_1 - \Upsilon^e a_0), \end{aligned}$$

and for $n \geq 3$:

$$a_n - \Upsilon^e a_{n-1} = b_n - \Upsilon^e b_{n-1} + \Upsilon^e b_{n-3} + \sum_{i=0}^{n-3} b_i(a_{n-2-i} - \Upsilon^e a_{n-3-i}).$$

Since R is not a hypersurface, the ring Q is not a hypersurface and $\text{edim } Q = e$. By Lemma (5.3) the series $(1 - \Upsilon^e t + \Upsilon^e t^3) P_k^Q(t)$ has non-negative coefficients, which means that $b_n - \Upsilon^e b_{n-1} + \Upsilon^e b_{n-3} \geq 0$ for all $n \geq 1$. It now follows by recursion that $a_n - \Upsilon^e a_{n-1} \geq 0$ for all $n \geq 1$. \square

APPENDIX. FREE RESOLUTIONS OVER ARTINIAN RINGS

Here we collect a few results on the growth of Betti numbers of modules over an artinian local ring $(R, \mathfrak{m}, \mathfrak{k})$. Their main application in this paper is to the module $E_R(\mathfrak{k})$, the injective hull of \mathfrak{k} .

The first result, Lemma (A.1) below, proves the characterization (1.6) of Gorenstein rings. Indeed, let Q be a Cohen–Macaulay local ring of dimension d and let $\mathbf{x} = x_1, \dots, x_d$ be a Q -regular sequence. Then $R = Q/(\mathbf{x})$ is artinian, and one has

$$\mu^{d+i}(Q) = \mu^i(R) = \beta_i^R(E_R(\mathfrak{k})) \quad \text{for all } i \geq 0;$$

see (4.2.1) and [8, prop. 1.2.(c)]. If Q is not Gorenstein, then R is not Gorenstein and Lemma (A.1) applies to $E_R(\mathfrak{k})$ and yields $\mu^{d+i}(Q) \geq 2$ for all $i \geq 0$.

(A.1) **Lemma.** *Let M be a non-free finitely generated R -module. If $\ell(R)$ divides $\ell_R(M)$, then one has*

$$\beta_i^R(M) \geq 2 \quad \text{for all } i \geq 0.$$

Moreover, if $\ell_R(M) = \ell(R)$, then the inequality $\beta_1^R(M) \geq \beta_0^R(M)$ holds.

Proof. Since M is not free, we have $\beta_i^R(M) \geq 1$ for all $i \geq 0$. For every $j \geq 0$, a length computation based on the exact sequence

$$(1) \quad 0 \rightarrow M_{j+1} \rightarrow R^{\beta_j^R(M)} \rightarrow \dots \rightarrow R^{\beta_0^R(M)} \rightarrow M \rightarrow 0$$

shows that $\ell_R(M) \equiv \pm \ell_R(M_{j+1}) \pmod{\ell(R)}$. By assumption, $\ell(R)$ divides $\ell_R(M)$, so it follows that $\ell(R)$ divides $\ell_R(M_{j+1})$; in particular, we have $\ell(R) \leq \ell_R(M_{j+1})$.

Assume that $\beta_j^R(M) = 1$ for some $j \geq 0$. Then the embedding $0 \rightarrow M_{j+1} \rightarrow R$ yields $\ell(R) > \ell_R(M_{j+1})$, which contradicts the inequality obtained above. Therefore, $\beta_i^R(M)$ is at least 2 for all $i \geq 0$.

Now assume that $\ell_R(M) = \ell(R)$. For $j = 1$ the sequence (1) gives

$$\ell_R(M_2) = (\beta_1^R(M) - \beta_0^R(M) + 1) \ell(R).$$

As $\ell_R(M_2) > 0$, this forces the desired inequality. \square

In the following, r denotes the socle rank of R . We set $h = \max\{i \mid \mathfrak{m}^i \neq 0\}$ and adopt the notation from (4.1). For a finitely generated R -module M , the rank of the largest \mathfrak{k} -vector space that is a direct summand of M is called the \mathfrak{k} -rank of M .

The next proposition applies to rings of large socle rank, compared to the length of the ring. It complements a result of Gasharov on Peeva [10, prop. (2.2)] that applies to rings of large embedding dimension.

(A.2) Proposition. *Assume that $h \geq 2$ and $r > \ell_R(R/\mathfrak{m}^h) - 1$. For every finitely generated non-free R -module M , the sequence $\{\beta_i^R(M)\}_{i \geq 1}$ is increasing, and for $i \geq 2$ the next inequality holds*

$$\beta_{i+1}^R(M) \geq \frac{r}{\ell_R(R/\mathfrak{m}^h) - 1} \beta_i^R(M).$$

Proof. Note that the assumption $h \geq 2$ yields $\ell_R(R/\mathfrak{m}^h) \geq 2$, thus the quotient $r/(\ell_R(R/\mathfrak{m}^h) - 1)$ is well-defined. For $i \geq 1$ set $b_i = \beta_i^R(M)$ and let s_i be the \mathfrak{k} -rank of M_i . Write the syzygy M_{i+1} as a direct sum $M_{i+1} \cong \mathfrak{k}^{s_{i+1}} \oplus N^{(i+1)}$, where \mathfrak{k} is not a direct summand of $N^{(i+1)}$. The isomorphism $\text{Soc } R^{b_i} \cong \mathfrak{k}^{s_{i+1}} \oplus \text{Soc } N^{(i+1)}$, see (4.1.2), explains the second equality below.

$$\begin{aligned} rb_i = \text{rank}_{\mathfrak{k}} \text{Soc } R^{b_i} &= s_{i+1} + \text{rank}_{\mathfrak{k}} \text{Soc } N^{(i+1)} \\ &\leq s_{i+1} + \ell_R(\mathfrak{m}N^{(i+1)}) \\ &= s_{i+1} + \ell_R(N^{(i+1)}) - \beta_0^R(N^{(i+1)}) \\ &= s_{i+1} + \ell_R(N^{(i+1)}) - (b_{i+1} - s_{i+1}) \\ &\leq s_{i+1} + (b_{i+1} - s_{i+1}) \ell(R/\mathfrak{m}^h) - (b_{i+1} - s_{i+1}) \\ &= (\ell(R/\mathfrak{m}^h) - 1)b_{i+1} + (2 - \ell(R/\mathfrak{m}^h))s_{i+1} \\ &\leq (\ell(R/\mathfrak{m}^h) - 1)b_{i+1} \end{aligned}$$

The first inequality uses the containment $\text{Soc } N^{(i+1)} \subseteq \mathfrak{m}N^{(i+1)}$, which holds as \mathfrak{k} is not a direct summand of $N^{(i+1)}$; see (4.1.1). The second inequality follows as $N^{(i+1)}$, being a summand of a syzygy, is an R/\mathfrak{m}^h module generated by $b_{i+1} - s_{i+1}$ elements. The last inequality holds as we have $\ell_R(R/\mathfrak{m}^h) \geq 2$. \square

Applied to the module $E_R(\mathfrak{k})$, the next result establishes the termwise exponential growth of the sequence $\{\mu^i(R)\}_{i \geq 0}$ stated in Proposition (3.8), see (4.2.1). If there is more than one minimal generator of the maximal ideal in the socle of R , then Proposition (A.3) gives a higher rate of growth than Peeva's [21, lem. 6].

(A.3) Proposition. *Assume that R satisfies $\mathfrak{m}^2 \neq 0$, $\text{Soc } R \not\subseteq \mathfrak{m}^2$ and $\text{edim } R \geq 2$. For every finitely generated non-free R -module M , the sequence $\{\beta_i^R(M)\}_{i \geq 1}$ is increasing, and for $i \geq 2$ the next inequality holds*

$$\beta_{i+1}^R(M) \geq (r - \text{rank}_{\mathfrak{k}}(\mathfrak{m}^2 \cap \text{Soc } R)) \left(1 + \frac{1}{\ell(R) - 1}\right) \beta_i^R(M).$$

Proof. For every $i \geq 0$ set $b_i = \beta_i^R(M)$. By assumption there is a minimal generator of \mathfrak{m} in $\text{Soc } R$. As $\text{Soc } R^{b_1}$ is contained in M_2 , this generator gives b_1 elements in $M_2 \setminus \mathfrak{m}M_2$ that are linearly independent modulo $\mathfrak{m}M_2$. It follows from the assumptions on R that $\text{rank}_{\mathfrak{k}} \text{Soc } R$ is at least 2, so there are elements in M_2 which are not in the span of these b_1 elements. This proves the inequality $b_2 > b_1$.

Set $A = r - \text{rank}_{\mathfrak{k}}(\mathfrak{m}^2 \cap \text{Soc } R)$. By assumption A is at least 1, and since $\mathfrak{m}^2 \neq 0$ there is a minimal generator of \mathfrak{m} outside the socle, so $e \geq A + 1$. For $i \geq 2$ write M_i as a direct sum $M_i \cong \mathfrak{k}^{s_i} \oplus N^{(i)}$, where \mathfrak{k} is not a summand of $N^{(i)}$. By (4.1.2)

we have $\text{Soc } M_i = \text{Soc } R^{b_{i-1}}$. A \mathbf{k} -vector space of rank Ab_{i-1} is a direct summand of $\mathfrak{m}R^{b_{i-1}}$ and hence in $\text{Soc } M_i \setminus \mathfrak{m}M_i$. In particular, there is an inequality

$$(1) \quad s_i \geq Ab_{i-1}.$$

There is an isomorphism $M_{i+1} \cong \mathfrak{m}^{s_i} \oplus (N^{(i)})_1$, so M_{i+1} decomposes as a direct sum $\mathfrak{m}^{s_i} \oplus \mathbf{k}^s \oplus N$, where \mathbf{k} is not a summand of N . Notice the equality $As_i + s = s_{i+1}$; it explains the second equality in the computation below.

$$\begin{aligned} b_{i+1} &= es_i + s + \beta_0^R(N) \geq (A+1)s_i + s \\ &= s_{i+1} + s_i \\ &\geq Ab_i + Ab_{i-1} \\ &\geq Ab_i \left(1 + \frac{1}{\ell(R) - 1} \right) \end{aligned}$$

The penultimate inequality follows from (1) and the last one from [2, lem. 4.2.7]. \square

(A.4) **Rings with radical cube zero.** Assume $\mathfrak{m}^3 = 0 \neq \mathfrak{m}^2$ and fix the notation:

$$(A.4.1) \quad a = \text{rank}_{\mathbf{k}} \mathfrak{m}^2, \quad e = \text{rank}_{\mathbf{k}} \mathfrak{m}/\mathfrak{m}^2, \quad \text{and} \quad r = \text{rank}_{\mathbf{k}} \text{Soc } R.$$

Notice that the inclusion $\mathfrak{m}^2 \subseteq \text{Soc } R$ yields the inequality $a \leq r$.

Let M be a finitely generated R -module, and set $b_i = \beta_i^R(M)$ for $i \geq 0$. We recall a few facts from Lescot's [19, §3]. There is an inequality

$$(A.4.2) \quad b_1 \geq eb_0 - \text{rank}_{\mathbf{k}}(\mathfrak{m}M/\mathfrak{m}^2M).$$

If \mathbf{k} is not a direct summand of M_1 , then equality holds; the converse is true if $\text{Soc } R = \mathfrak{m}^2$.

For every $i \geq 2$ there is an inequality

$$(A.4.3) \quad b_i \geq eb_{i-1} - ab_{i-2}.$$

If \mathbf{k} is not a summand of M_1 and not a summand of M_2 , then the equality

$$(A.4.4) \quad b_2 = eb_1 - ab_0$$

holds if and only if $\mathfrak{m}^2M = 0$.

If $i \geq 3$, and \mathbf{k} is not a summand of M_i and not a summand of M_{i-1} , then

$$(A.4.5) \quad b_i = eb_{i-1} - ab_{i-2}.$$

If $\text{Soc } R = \mathfrak{m}^2$ and $\mathfrak{m}^2M = 0$, then

$$(A.4.6) \quad eb_1 \geq rb_0 + s(e-1),$$

where s is the \mathbf{k} -rank of M_1 .

The next result strengthens parts of Lescot's [19, thm. B].

(A.5) **Theorem.** *Let R be a local ring with $\mathfrak{m}^3 = 0 \neq \mathfrak{m}^2$ that is not Gorenstein. Let M be a finitely generated R -module that satisfies $\beta_1^R(M) > \beta_0^R(M)$. With the notation from (A.4.1) the following statements hold*

- (a) *If $a \neq e$ or $a \neq r$, then the sequence $\{\beta_i^R(M)\}_{i \geq 0}$ is increasing and has termwise exponential growth.*
- (b) *If $a = e = r$, then the sequence $\{\beta_i^R(M)\}_{i \geq 0}$ is non-decreasing with strict inequalities $\beta_i^R(M) > \beta_{i-1}^R(M)$ for all but, possibly, one index i . Moreover, if there is an equality $\beta_j^R(M) = \beta_{j-1}^R(M)$, then j is at least 2, and \mathbf{k} is a direct summand of M_{j+1} and not a direct summand of M_i when $0 \leq i \leq j$.*

Proof. For $i \geq 0$ set $b_i = \beta_i^R(M)$.

(a): First assume $a \neq r$, that is, $\mathfrak{m}^2 \neq \text{Soc } R$. By [19, prop. 3.9] and the assumption on M , the sequence $\{b_i\}_{i \geq 0}$ is increasing, and by Proposition (A.3) it has termwise exponential growth of rate $(r-a)(1+1/(a+e))$.

Next, assume $a = r$. There are two cases to consider:

Case $e < a$: Since $\ell_R(R/\mathfrak{m}^2) - 1 = e$, it follows from Proposition (A.2) and the assumption on M that the sequence $\{b_i\}_{i \geq 0}$ is increasing and has termwise exponential growth of rate a/e .

Case $e > a$: For every $i \geq 2$, the inequality (A.4.3) yields

$$b_i - b_{i-1} \geq (e-1)b_{i-1} - ab_{i-2} \geq a(b_{i-1} - b_{i-2}).$$

By recursion, based on the assumption $b_1 > b_0$, it follows that the sequence $\{b_i\}_{i \geq 0}$ is increasing. If $e = a + 1$, then the sequence has termwise exponential growth by [21, prop. 3]. If $e > a + 1$, then the same conclusion follows as (A.4.3) yields

$$b_i > (e-a)b_{i-1} \quad \text{for all } i \geq 2.$$

(b): Assume that $a = e = r$. First notice that if $i \geq 3$ and \mathfrak{k} is not a summand of M_i and not a summand of M_{i-1} , then (A.4.5) yields $b_{i-1} > b_{i-2}$, as $b_i > 0$ by assumption. It follows that the sequence $\{b_i\}_{i \geq 0}$ is increasing if for every $i \geq 2$ the residue field \mathfrak{k} is not a direct summand of M_i .

Next, assume that \mathfrak{k} is a summand of a syzygy of M , and let j be the least integer such that $M_{j+1} = \mathfrak{k} \oplus N$ for some R -module N . The sequence $\{b_i\}_{i \geq j+1}$ is then increasing. Indeed, the sequence $\{\beta_i^R(\mathfrak{k})\}_{i \geq 0}$ is increasing, as R is not a hypersurface. If $j \geq 0$, then the sequence $\{\beta_i^R(N)\}_{i \geq 0}$ is non-decreasing by (A.4.6), as $\mathfrak{m}^2 N = 0$. If $j = -1$, i.e. \mathfrak{k} is a summand of M , then the sequence $\{\beta_i^R(N)\}_{i \geq 1}$ is non-decreasing by (A.4.6), and $b_1 > b_0$ by assumption. If $j = -1$ or $j = 0$, it is thus immediate that the sequence $\{b_i\}_{i \geq 0}$ is increasing.

We can now assume that j is at least 1. In the next chain of inequalities, the right-most and left-most ones are already known; the inequalities in-between follow by application of (A.4.6) to the syzygies M_1, \dots, M_j .

$$\dots > b_{j+1} > b_j \geq \dots \geq b_1 > b_0$$

If $j = 1$, it follows that the sequence $\{b_i\}_{i \geq 0}$ is increasing. If $j \geq 2$, then $b_{i-1} > b_{i-2}$ for i with $j \geq i \geq 2$ by (A.4.5) and the assumption on M . In total, this gives inequalities $\dots > b_{j+1} > b_j \geq b_{j-1} > \dots > b_1 > b_0$. \square

ACKNOWLEDGMENTS

We thank Luchezar L. Avramov and Sean Sather-Wagstaff for extensive comments on an earlier version of the paper. Thanks are also due to Christopher Monico and Roger W. Barnard for discussions related to the material in Section 5.

REFERENCES

1. David J. Anick, *A counterexample to a conjecture of Serre*, Ann. of Math. (2) **115** (1982), no. 1, 1–33. MR644015
2. Luchezar L. Avramov, *Infinite free resolutions*, Six lectures on commutative algebra (Bellaterra, 1996), Progr. Math., vol. 166, Birkhäuser, Basel, 1998, pp. 1–118. MR1648664
3. Luchezar L. Avramov, Srikanth Iyengar, and Claudia Miller, *Homology over local homomorphisms*, Amer. J. Math. **128** (2006), no. 1, 23–90. MR2197067
4. Luchezar L. Avramov and Jack Lescot, *Bass numbers and Golod rings*, Math. Scand. **51** (1982), no. 2, 199–211 (1983). MR690525

5. Luchezar L. Avramov and Oana Veliche, *Stable cohomology over local rings*, Adv. Math. **213** (2007), no. 1, 93–139. MR2331239
6. Jürgen Backelin and Ralf Fröberg, *Poincaré series of short Artinian rings*, J. Algebra **96** (1985), no. 2, 495–498. MR810542
7. Hyman Bass, *Injective dimension in Noetherian rings*, Trans. Amer. Math. Soc. **102** (1962), 18–29. MR0138644
8. Hans-Bjørn Foxby, *On the μ^i in a minimal injective resolution*, Math. Scand. **29** (1971), 175–186 (1972). MR0309919
9. ———, *On the μ^i in a minimal injective resolution II*, Math. Scand. **41** (1977), no. 1, 19–44. MR0476801
10. Vesselin N. Gasharov and Irena V. Peeva, *Boundedness versus periodicity over commutative local rings*, Trans. Amer. Math. Soc. **320** (1990), no. 2, 569–580. MR967311
11. Daniel R. Grayson and Michael E. Stillman, *Macaulay 2, a software system for research in algebraic geometry*, available from <http://www.math.uiuc.edu/Macaulay2/>.
12. Tor H. Gulliksen, *On the deviations of a local ring*, Math. Scand. **47** (1980), no. 1, 5–20. MR600076
13. Jürgen Herzog and Manfred Steurich, *Two applications of change of rings theorems for Poincaré series*, Proc. Amer. Math. Soc. **73** (1979), no. 2, 163–168. MR516457
14. Craig Huneke and Adela Vraciu, *Rings that are almost Gorenstein*, Pacific J. Math. **225** (2006), no. 1, 85–102. MR2233726
15. David A. Jorgensen and Graham J. Leuschke, *On the growth of the Betti sequence of the canonical module*, Math. Z. **256** (2007), no. 3, 647–659. MR2299575
16. Aleksei I. Kostrikin and Igor R. Šafarevič, *Groups of homologies of nilpotent algebras*, Dokl. Akad. Nauk SSSR (N.S.) **115** (1957), 1066–1069. MR0092776
17. Jack Lescot, *La série de Bass d'un produit fibré d'anneaux locaux*, C. R. Acad. Sci. Paris Sér. I Math. **293** (1981), no. 12, 569–571. MR647683
18. ———, *La série de Bass d'un produit fibré d'anneaux locaux*, Paul Dubreil and Marie-Paule Malliavin algebra seminar, 35th year (Paris, 1982), Lecture Notes in Math., vol. 1029, Springer, Berlin, 1983, pp. 218–239. MR732477
19. ———, *Asymptotic properties of Betti numbers of modules over certain rings*, J. Pure Appl. Algebra **38** (1985), no. 2-3, 287–298. MR814184
20. Gerson L. Levin and Luchezar L. Avramov, *Factoring out the socle of a Gorenstein ring*, J. Algebra **55** (1978), no. 1, 74–83. MR515760
21. Irena Peeva, *Exponential growth of Betti numbers*, J. Pure Appl. Algebra **126** (1998), no. 1-3, 317–323. MR1600558
22. Paul Roberts, *Rings of type 1 are Gorenstein*, Bull. London Math. Soc. **15** (1983), no. 1, 48–50. MR686348
23. Günter Scheja, *Über die Bettizahlen lokaler Ringe*, Math. Ann. **155** (1964), 155–172. MR0162819
24. Jean-Pierre Serre, *Local algebra*, Springer Monographs in Mathematics, Springer-Verlag, Berlin, 2000, Translated from the French by CheeWhye Chin and revised by the author. MR1771925
25. Li-Chuan Sun, *Growth of Betti numbers of modules over local rings of small embedding codimension or small linkage number*, J. Pure Appl. Algebra **96** (1994), no. 1, 57–71. MR1297441
26. William Teter, *Rings which are a factor of a Gorenstein ring by its socle*, Invent. Math. **23** (1974), 153–162. MR0366911
27. Wolmer V. Vasconcelos, *Divisor theory in module categories*, North-Holland Publishing Co., Amsterdam, 1974, North-Holland Mathematics Studies, No. 14, Notas de Matemática No. 53. [Notes on Mathematics, No. 53]. MR0498530
28. Hartmut Wiebe, *Über homologische Invarianten lokaler Ringe*, Math. Ann. **179** (1969), 257–274. MR0255531

L. W. CHRISTENSEN, DEPARTMENT OF MATHEMATICS AND STATISTICS, TEXAS TECH UNIVERSITY, LUBBOCK, TEXAS 79409, U.S.A.

E-mail address: `lars.w.christensen@ttu.edu`

URL: `http://www.math.ttu.edu/~lchrste`

J. STRIULI, DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, FAIRFIELD UNIVERSITY, FAIRFIELD, CONNECTICUT 06824, U.S.A.

E-mail address: `jstriuli@fairfield.mail.edu`

O. VELICHE, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF UTAH, SALT LAKE CITY, UTAH 84112, U.S.A.

E-mail address: `oveliche@math.utah.edu`

URL: `http://www.math.utah.edu/~oveliche`