

## Inverse Trig Functions

Math 1220 (Spring 2003)

Recall the trig functions and their derivatives:

$$\sin(x), D_x(\sin(x)) = \cos(x)$$

$$\cos(x), D_x(\cos(x)) = -\sin(x)$$

$$\tan(x) = \frac{\sin(x)}{\cos(x)}, D_x(\tan(x)) = \sec^2(x)$$

$$\cot(x) = \frac{\cos(x)}{\sin(x)}, D_x(\cot(x)) = -\csc^2(x)$$

$$\sec(x) = \frac{1}{\cos(x)}, D_x(\sec(x)) = \sec(x)\tan(x)$$

$$\csc(x) = \frac{1}{\sin(x)}, D_x(\csc(x)) = -\csc(x)\cot(x)$$

As we discussed earlier, we can now find inverse functions:

$$\sin^{-1}(y), \text{ Domain: } (-1, 1) \text{ Range: } \left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$$

$$\cos^{-1}(y), \text{ Domain: } (-1, 1) \text{ Range: } (0, \pi)$$

$$\tan^{-1}(y), \text{ Domain: } (-\infty, \infty) \text{ Range: } \left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$$

$$\sec^{-1}(y), \text{ Domain: } (-\infty, -1) \cup (1, \infty) \text{ Range: } \left(0, \frac{\pi}{2}\right) \cup \left(\frac{\pi}{2}, \pi\right)$$

Recall that we get the domain and range of the inverse function  $f^{-1}(y)$  by restricting to domains where  $f(x)$  is increasing (or decreasing) so that the inverse function exists, and then swapping them. For example,  $\sin(x)$  is increasing on  $(-\frac{\pi}{2}, \frac{\pi}{2})$ ,  $\cos(x)$  is decreasing on  $(0, \pi)$ , and  $\tan(x)$  is increasing on  $(-\frac{\pi}{2}, \frac{\pi}{2})$ .  $\sec(x)$  is a little different. It is increasing and positive on  $(0, \frac{\pi}{2})$  and increasing and negative on  $(\frac{\pi}{2}, \pi)$ , so we can keep both these intervals.

Recall the magical formula for the derivative of the inverse function:

$$(f^{-1})'(y) = \frac{1}{f'(x)} \text{ whenever } y = f(x), \text{ or equivalently } f^{-1}(y) = x$$

**Examples:** (i) Find  $\sin^{-1}(\frac{1}{2})$  and  $(\sin^{-1})'(\frac{1}{2})$ .

$\sin^{-1}(\frac{1}{2}) = \frac{\pi}{6}$  because  $\sin(\frac{\pi}{6}) = \frac{1}{2}$  and  $\frac{\pi}{6}$  is in the range

$$(\sin^{-1})'(\frac{1}{2}) = \frac{1}{\sin'(\sin^{-1}(\frac{1}{2}))} = \frac{1}{\cos(\frac{\pi}{6})} = \frac{2}{\sqrt{3}}$$

(ii) Find  $\tan^{-1}(-1)$  and  $(\tan^{-1})'(-1)$ .

$$\tan^{-1}(-1) = -\frac{\pi}{4}$$

$$(\tan^{-1})'(-1) = \frac{1}{\sec^2(-\frac{\pi}{4})} = \cos^2(-\frac{\pi}{4}) = \frac{1}{2}$$

(iii) Find  $\sec^{-1}(-2)$  and  $(\sec^{-1})'(-2)$

$\sec^{-1}(-2) = \frac{2\pi}{3}$  because  $\cos(\frac{2\pi}{3}) = -\frac{1}{2}$  and  $\frac{2\pi}{3}$  is in the range

$$(\sec^{-1})'(-2) = \frac{1}{\sec(\frac{2\pi}{3}) \tan(\frac{2\pi}{3})} = (-\frac{1}{2})(-\frac{1}{\sqrt{3}}) = \frac{1}{2\sqrt{3}}$$

OK. Now we want formulas for the derivatives of the inverse trig functions. We will be considerably helped by the following geometric results:

$$\sin(\cos^{-1}(x)) = \sqrt{1-x^2}$$

$$\cos(\sin^{-1}(x)) = \sqrt{1-x^2}$$

$$\sec(\tan^{-1}(x)) = \sqrt{1+x^2}$$

$$\tan(\sec^{-1}(x)) = \begin{cases} \sqrt{x^2-1} & \text{if } x > 1 \\ -\sqrt{x^2-1} & \text{if } x < -1 \end{cases}$$

(see the triangles on page 354 of the text for a visual proof)

Notice that by using these results, you can often get exact answers for  $\sin(\cos^{-1}(x))$  even when you cannot get exact answers for  $\cos^{-1}(x)$ .

**Examples:** (i)

$$\sin(\cos^{-1}(\frac{3}{5})) = \sqrt{1 - \frac{9}{25}} = \sqrt{\frac{16}{25}} = \frac{4}{5}$$

(ii)

$$\sec(\tan^{-1}(4)) = \sqrt{1+16} = \sqrt{17}$$

**Now for the Formulas:**

$$(\sin^{-1})'(y) = \frac{1}{\sin'(x)} = \frac{1}{\cos(\sin^{-1}(y))} = \frac{1}{\sqrt{1-y^2}}$$

$$(\cos^{-1})'(y) = \frac{1}{\cos'(x)} = \frac{1}{-\sin(\cos^{-1}(y))} = -\frac{1}{\sqrt{1-y^2}}$$

$$(\tan^{-1})'(y) = \frac{1}{\tan'(x)} = \frac{1}{\sec^2(\tan^{-1}(y))} = \frac{1}{1+y^2}$$

$$\begin{aligned} (\sec^{-1})'(y) &= \frac{1}{\sec'(x)} = \frac{1}{\sec(\sec^{-1}(y)) \tan(\sec^{-1}(y))} \\ &= \begin{cases} \frac{1}{y\sqrt{y^2-1}} & \text{if } y > 1 \\ -\frac{1}{y\sqrt{y^2-1}} & \text{if } y < -1 \end{cases} = \frac{1}{|y|\sqrt{y^2-1}} \end{aligned}$$

**Examples:** Differentiate using the chain rule:

$$D_x(\sin^{-1}(e^x)) = D_x(\sin^{-1}(u)) = \frac{1}{\sqrt{1-u^2}} \cdot u' = \frac{e^x}{\sqrt{1-e^{2x}}}$$

(using the substitution  $u = e^x$ )

$$D_x(\tan^{-1}(x^a)) = D_x(\tan^{-1}(u)) = \frac{u'}{1+u^2} = \frac{ax^{a-1}}{1+x^{2a}}$$

(using the substitution  $u = x^a$ )

$$D_x(\sec^{-1}(a^x)) = \frac{\ln(a)a^x}{a^x\sqrt{a^{2x}-1}} = \frac{\ln(a)}{\sqrt{a^{2x}-1}}$$

**More Examples:** Some surprising improper integrals:

$$\int_0^{\frac{1}{2}} \frac{1}{\sqrt{1-x^2}} dx = \sin^{-1}(x) \Big|_0^{\frac{1}{2}} = \frac{\pi}{6} - 0 = \frac{\pi}{6}$$

$$\int_1^{\sqrt{3}} \frac{1}{1+x^2} dx = \tan^{-1}(x) \Big|_1^{\sqrt{3}} = \frac{\pi}{3} - \frac{\pi}{4} = \frac{\pi}{12}$$

**Some Improper Integrals:**

$$\int \frac{x}{1+x^4} dx =$$

Hmmm, looks like the derivative of  $\tan^{-1}$ . Let's try  $u = x^2, du = 2x dx$

$$\int \frac{x dx}{1+x^4} = \int \frac{\frac{1}{2} du}{1+u^2} = \frac{1}{2} \tan^{-1}(u) = \frac{1}{2} \tan^{-1}(x^2)$$

**A Word Problem:** Suppose you are standing 4 miles from a rocket that takes off with a constant acceleration of 32 feet/sec<sup>2</sup> (so the crew feels 2Gs). What is the acceleration in the angle you see between the rocket and the horizon at time  $t$ ?

The height of the rocket is given by the formula:

$$h(t) = 32 \frac{t^2}{2} = 16t^2 \text{ (integrating the acceleration twice)}$$

so the angle you see (as a function of time  $t$ ) is:

$$\tan^{-1} \left( \frac{16t^2}{4} \right) = \tan^{-1} (4t^2)$$

so to get the *acceleration* in this angle, we need to differentiate twice(!)

$$(\tan^{-1})' (4t^2) = \frac{8t}{1+16t^4}$$

$$(\tan^{-1})'' (4t^2) = D_t \left( \frac{8t}{1+16t^4} \right) = \frac{8(1+16t^4) - 8t(64t^3)}{(1+16t^4)^2} = \frac{8(1-48t^3)}{(1+16t^4)^2}$$

measured in radians/sec<sup>2</sup>.