## DYLAN ZWICK'S MATH 1010 CLASS

## 9.1 EXPONENTIAL FUNCTIONS

Simplify the expression.

**9.1.1:** 
$$3^x \cdot 3^{x+2}$$
 **9.1.4:**  $\frac{3^{2x+3}}{3^{x+1}}$ 

**9.1.3:** 
$$\frac{e^{x+2}}{e^x}$$
 **9.1.5:**  $3(e^x)^{-2}$ 

Evaluate the function as indicated. Use a calculator ONLY IF it is necessary or more efficient.

**9.1.17:** 
$$f(x) = 3^x, (a)x = -2, (b)x = 0, (c)x = 1.$$

**9.1.21:** 
$$f(t) = 500(\frac{1}{2})^t, (a)t = 0, (b)t = 1, (c)t = \pi.$$

**9.1.26:** 
$$P(t) = \frac{10,000}{(1.01)^{12t}}, (a)t = 2, (b)t = 10, (c)t = 20$$

Sketch the graph of the function. Identify the horizontal asymptote.

**9.1.31:** 
$$f(x) = 3^x$$
 **9.1.35:**  $g(x) = 3^x - 2$ 

**9.1.32:** 
$$h(x) = \frac{1}{2}(3^x)$$
 **9.1.40:**  $f(t) = 2^{t^2}$ 

**9.1.33:**  $f(x) = 3^{-x}$ 

**9.1.69:** Radioactive Decay After t years, 16 grams of a radioactive element with a half-life of 30 years decays to a mass y(in grams) given by  $y = 16(\frac{1}{2})^{t/30}, t \ge 0$ . How much of the initial mass remains after 80 years?

**9.1.70:** Radioactive Substance In July of 1999, an individual bought several leaded containers from a metals recycler and found two of them labeled" radioactive". An investigation showed that the containers, originally obtained from Ohio State University, apparently had been used to store iodine-131 starting in January of 1999. Because iodine-131 has a half life of only 8 days, no elevated radiation levels were detected. Suppose 6 grams of iodine-131 is stored in January. The mass y ( in grams) that remains after t days is given by  $y = 6(\frac{1}{2})^{t/8}, t \ge 0$ . How much of the substance is left in July, after 180 days have passed.

**9.1.71:** Compound Interest A sum of \$5000 is invested at an annual interest rate of 6%, compounded monthly. Find the balance in the account after 5 years.

**9.1.72:** Compound Interest A sum of \$2000 is invested at an annual interest rate of 8%, compounded quarterly. Find the balance in the account after 10 years.

Compound Interest Complete the table to determine the balance A for P dollars invested at rate r for t years, compounded n times per year.

**9.1.73:** P = \$100, r = 7%, t = 15 years.

n	1	4	12	365	Continuous compounding
A					

**9.1.75:** P = \$2000, r = 9.5%, t = 10 years.

n	1	4	12	365	Continuous compounding
A					

Compound Interest Complete the table to determine the principal P that will yield a balance of A dollars when invested at rate r for t year, compounded n times per year.

**9.1.77:** A = \$5000, r = 7%, t = 10 years

					· · · · · · · · · · · · · · · · · · ·
n	1	4	12	365	Continuous compounding
P					

9.	<b>9.1.79:</b> $A = \$1,000,000, r = 10.5\%, t = 40$ ye									
	n	1	4	12	365	Continuous compounding				
	P									

**9.1.82:** Population Growth The populations P (in millions) of the United States from 1980 to 2006 can be approximated by the exponential function  $P(t) = 226(1.0110)^t$ , where t is the time in years, with t = 0 corresponding to 1980. Use the model to estimate the populations in years (a) 2010 and (b) 2020.

### 9.2 Composite and Inverse Functions

Find the compositions.

**9.2.1:** f(x) = 2x + 3, g(x) = x - 6

(a) 
$$(f \circ g)(x)$$
 (c)  $(f \circ g)(4)$ 

(b) 
$$(g \circ f)(x)$$
 (d)  $(g \circ f)(7)$ 

**9.2.2:** 
$$f(x) = x - 5, g(x) = 3x + 2$$

(a) 
$$(f \circ g)(x)$$
 (c)  $(f \circ g)(3)$ 

(b) 
$$(g \circ f)(x)$$
 (d)  $(g \circ f)(3)$ 

**9.2.3:** 
$$f(x) = x^2 + 3, g(x) = x + 2$$
  
(a)  $(f \circ g)(x)$  (c)  $(f \circ g)(2)$ 

(b) 
$$(g \circ f)(x)$$
 (d)  $(g \circ f)(-3)$ 

**9.2.5:** 
$$f(x) = |x - 3|, g(x) = 3x$$
  
(a)  $(f \circ g)(x)$  (c)  $(f \circ g)(1)$ 

(b) 
$$(g \circ f)(x)$$
 (d)  $(g \circ f)(2)$ 

**9.2.8:** 
$$f(x) = \sqrt{x+6}, g(x) = 2x-3$$
  
(a)  $(f \circ g)(x)$  (c)  $(f \circ g)(3)$ 

(b) 
$$(g \circ f)(x)$$
 (d)  $(g \circ f)(-2)$ 

**9.2.10:** 
$$f(x) = \frac{4}{x^2 - 4}, g(x) = \frac{1}{x}$$
  
(a)  $(f \circ g)(x)$  (c)  $(f \circ g)(-2)$ 

(b) 
$$(g \circ f)(x)$$
 (d)  $(g \circ f)(1)$ 

**9.2.11:** Use the functions f and g to find the indicated values.  $f = \{(-2,3), (-1,1), (0,0), (1,-1), (2,-3)\}$   $g = \{(-3,1), (-1,-2), (0,2), (2,2), (3,1))\}$ (a) f(1) (c)  $(g \circ f)(1)$ 

(b) g(-1)

**9.2.15:** Use the functions f and g to find the indicated values.  $f = \{(0, 1), (1, 2), (2, 5), (3, 10), (4, 17))\}$   $g = \{(5, 4), (10, 1), (2, 3), (17, 0), (1, 2)\}$ (a) f(2) (c)  $(g \circ f)(1)$ 

(b) g(10)

Find the compositions.  $(a)f\circ g$  and  $(b)g\circ f.$  Then find the domain of each composition.

**9.2.19:** 
$$f(x) = 3x + 4, g(x) = x - 7$$

**9.2.21:** 
$$f(x) = \sqrt{x+2}, g(x) = x-4$$

**9.2.23:** 
$$f(x) = x^2 + 3, g(x) = \sqrt{x-1}$$

**9.2.26:** 
$$f(x) = \frac{x}{x-4}, g(x) = \sqrt{x}$$

Use the Horizontal Line test to determine if the function is one-to-one and so has an inverse function.

**9.2.35:**  $f(x) = x^2 - 2$ 



**9.2.37:**  $f(x) = x^2, x \ge 0$ 



**9.2.39:**  $g(x) = \sqrt{25 - x^2}$ 



Verify algebraically that the functions f and g are inverse functions of each other.

**9.2.41:**  $f(x) = -6x, g(x) = -\frac{1}{6}x$ 

**9.2.43:** 
$$f(x) = 1 - 2x, g(x) = \frac{1}{2}(1 - x)$$

**9.2.45:** 
$$f(x) = \sqrt[3]{x+1}, g(x) = x^3 - 1$$

Find the inverse function of f. Verify that  $f(f^{-1}(x))$  and  $f^{-1}(f(x))$  are equal to the identity function.

**9.2.49:** 
$$f(x) = 5x$$
 **9.2.55:**  $f(x) = 5 - x$ 

**9.2.52:** 
$$f(x) = \frac{1}{3}x$$
 **9.2.59:**  $f(x) = \sqrt[3]{x}$ 

Find the inverse function (if it exists).

**9.2.64:** 
$$g(t) = 6t + 1$$
 **9.2.67:**  $g(x) = x^2 + 4$ 

**9.2.70:** 
$$h(x) = \sqrt{x+5}$$

**9.2.105:** Ripples You are standing on a bridge over a calm pond and drop a pebble, causing ripples of concentric circles in the water. The radius (in feet) of the outermost ripple is given by r(t) = 0.6t, where t is time in seconds after the pebble hits the water. The area of the circle is given by the function  $A(r) = \pi r^2$ . Find an equation for the composition A(r(t)). What are the input and output of this composite function? What is the area of the circle after 3 seconds. 9.3 EXPONENTIAL AND LOGARITHMIC FUNCTIONS Write the logarithmic equation in exponential form.

**9.3.2:** 
$$\log_{11} 121$$
 **9.3.8:**  $\log_{64} 4$ 

**9.3.3:** 
$$\log_2 \frac{1}{32}$$
 **9.3.10:**  $\log_{16} 8$ 

Write the exponential equation in logarithmic form.

**9.3.13:** 
$$6^2 = 36$$
  
**9.3.19:**  $25^{-1/2} = \frac{1}{5}$   
**9.3.15:**  $5^{-3} = \frac{1}{125}$   
**9.3.22:**  $6^1 = 6$ 

Evaluate the logarithmic without using a calculator. (If not possible, state the reason)

<b>9.3.25:</b> log <sub>2</sub> 8	<b>9.3.35:</b> $\log_2(-3)$
<b>9.3.29:</b> $\log_2 \frac{1}{16}$	<b>9.3.37:</b> log <sub>4</sub> 1
<b>9.3.32:</b> $\log_6 \frac{1}{216}$	<b>9.3.38:</b> log <sub>3</sub> 1
	<b>9.3.41:</b> log <sub>9</sub> 3

Sketch the graph of f, Then use the graph of f to sketch the graph of g.

**9.3.57:** 
$$f(x) = 3^x, g(x) = \log_3 x$$

Identify the transformation of the graph of  $f(x) = \log_2 x$ . Then sketch the graph of h.

**9.3.61:**  $h(x) = 3 + \log_2 x$  **9.3.64:**  $h(x) = \log_2(x+5)$ 

Sketch the graph of the function. Identify the vertical asymptote. 9.3.67:  $f(x) = \log_5 x$ 

**9.3.73:**  $g(x) = \log_2(x-3)$ 

**9.3.99:**  $f(x) = -\ln x$ 

**9.3.103:**  $f(x) = 3 + \ln x$ 

Solving Problems:

**9.3.125:** American Elk The antler spread a (in inches) and shoulder height h (in inches) of an adult male American Elk are related by the model

 $h = 116 \log_{10}(a + 40) - 176.$ 

Approximate to one decimal place the shoulder height of a male American Elk with an antler spread of 55 inches.

**9.3.127:** Compound Interest The time t in years for an investment to double in value when compounded continuously at interest rate r is given by  $t = \frac{\ln 2}{r}$ . Complete the table, which shows the "doubling times" for several interest rates.

r	0.07	0.08	0.09	0.10	0.11	0.12
t						