

Math 2280 - Lecture 23

Dylan Zwick

Fall 2013

In our last lecture we dealt with solutions to the system:

$$\mathbf{x}' = A\mathbf{x}$$

where A is an $n \times n$ matrix with n distinct eigenvalues. As promised, today we will deal with the question of what happens if we have less than n distinct eigenvalues, which is what happens if any of the roots of the characteristic polynomial are repeated.

This lecture corresponds with section 5.4 of the textbook, and the assigned problems from that section are:

Section 5.4 - 1, 8, 15, 25, 33

The Case of an Order 2 Root

Let's start with the case of an order 2 root.¹ So, our eigenvalue equation has a repeated root, λ , of multiplicity 2.

There are two ways this can go. The first possibility is that we have two distinct (linearly independent) eigenvectors associated with the eigenvalue λ . In this case, all is good, and we just use these two eigenvectors to create two distinct solutions.

¹Admittedly, not one of Sherlock Holmes's more popular mysteries.

Example - Find a general solution to the system:

$$\mathbf{x}' = \begin{pmatrix} 9 & 4 & 0 \\ -6 & -1 & 0 \\ 6 & 4 & 3 \end{pmatrix} \mathbf{x}$$

More room for the example problem

Well, that was no problem. Unfortunately, as we will see momentarily, it isn't always the case that we can find two linearly independent eigenvectors for the same eigenvalue.

Example - Calculate the eigenvalues and eigenvectors for the matrix:

$$\mathbf{A} = \begin{pmatrix} 1 & -3 \\ 3 & 7 \end{pmatrix}$$

In this situation we call this eigenvalue *defective*, and the *defect* of this eigenvalue is the difference between the multiplicity of the root and the number of linearly independent eigenvectors. In the example above the defect is of order 1.

Now, how does this relate to systems of differential equations, and how do we deal with it if a defective eigenvalue shows up?² Well, suppose we have the same matrix \mathbf{A} as above, and we're given the differential equation:

$$\mathbf{x}' = \mathbf{A}\mathbf{x}.$$

For this differential equation we know how to find one solution:

$$\begin{pmatrix} 1 \\ -1 \end{pmatrix} e^{4t}$$

but for a complete set of solutions we'll need another linearly independent solution. How do we get this second solution?

Based on experience we may think that, if $\mathbf{v}_1 e^{\lambda t}$ is a solution, then we can get a second solution of the form $\mathbf{v}_2 t e^{\lambda t}$. Let's try out this solution:

$$\mathbf{v}_2 e^{\lambda t} + \lambda \mathbf{v}_2 t e^{\lambda t} = \mathbf{A} \mathbf{v}_2 t e^{\lambda t}.$$

Unfortunately for this to be true it must be true when $t = 0$, which would imply that $\mathbf{v}_2 = \mathbf{0}$, which wouldn't be a linearly independent solution. So, this approach doesn't work.³

But wait, there's hope. We don't have to abandon this method entirely.⁴ Suppose instead we modify it slightly and replace $\mathbf{v}_2 t$ with $\mathbf{v}_1 t + \mathbf{v}_2$, where \mathbf{v}_1 is the one eigenvector that we were able to find. Well, if we plug this into our differential equation we get the relations:

²No, run and hide is not an acceptable answer.

³Nuts!

⁴Hooray!

$$\mathbf{v}_1 e^{\lambda t} + \lambda \mathbf{v}_1 t e^{\lambda t} + \lambda \mathbf{v}_2 e^{\lambda t} = \mathbf{A} \mathbf{v}_1 t e^{\lambda t} + \mathbf{A} \mathbf{v}_2 e^{\lambda t}.$$

If we equate the $e^{\lambda t}$ terms and the $t e^{\lambda t}$ terms we get the two equalities:

$$(\mathbf{A} - \lambda \mathbf{I}) \mathbf{v}_1 = 0$$

and

$$(\mathbf{A} - \lambda \mathbf{I}) \mathbf{v}_2 = \mathbf{v}_1.$$

Let's think about this. What this means is that, if we have a defective eigenvalue with defect 1, we can find two linearly independent solutions by simply finding a solution to the equation

$$(\mathbf{A} - \lambda \mathbf{I})^2 \mathbf{v}_2 = \mathbf{0}$$

such that

$$(\mathbf{A} - \lambda \mathbf{I}) \mathbf{v}_2 \neq \mathbf{0}.$$

If we can find such a vector \mathbf{v}_2 , then we can construct two linearly independent solutions:

$$\mathbf{x}_1(t) = \mathbf{v}_1 e^{\lambda t}$$

and

$$\mathbf{x}_2(t) = (\mathbf{v}_1 t + \mathbf{v}_2) e^{\lambda t}$$

where \mathbf{v}_1 is the non-zero vector given by $(\mathbf{A} - \lambda \mathbf{I}) \mathbf{v}_2$.

For our particular situation we have:

$$(\mathbf{A} - 4\mathbf{I})^2 = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}$$

So, *any* non-zero vector could potentially work as our vector \mathbf{v}_2 . If we try

$$\mathbf{v}_2 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

then we get:

$$\begin{pmatrix} -3 & -3 \\ 3 & 3 \end{pmatrix} \begin{pmatrix} 1 \\ 0 \end{pmatrix} = \begin{pmatrix} -3 \\ 3 \end{pmatrix}$$

So, our linearly independent solutions are:

$$\mathbf{x}_1(t) = \mathbf{v}_1 e^{4t} = \begin{pmatrix} -3 \\ 3 \end{pmatrix} e^{4t},$$

and

$$\mathbf{x}_2(t) = (\mathbf{v}_1 t + \mathbf{v}_2) e^{4t} = \begin{pmatrix} -3t + 1 \\ 3t \end{pmatrix} e^{4t}.$$

with corresponding general solution:

$$\mathbf{x}(t) = c_1 \mathbf{x}_1(t) + c_2 \mathbf{x}_2(t).$$

We note that our eigenvector \mathbf{v}_1 is not our original eigenvector, but is a multiple of it. That's fine.

The General Case

The vector \mathbf{v}_2 above is an example of something called a *generalized eigenvector*. If λ is an eigenvalue of the matrix \mathbf{A} , then a *rank r generalized eigenvector* associated with λ is a vector \mathbf{v} such that:

$$(\mathbf{A} - \lambda \mathbf{I})^r \mathbf{v} = \mathbf{0}$$

but

$$(\mathbf{A} - \lambda \mathbf{I})^{r-1} \mathbf{v} \neq \mathbf{0}.$$

We note that a rank 1 generalized eigenvector is just our standard eigenvector, where we treat a matrix raised to the power 0 as the identity matrix.

We define a *length k chain of generalized eigenvectors based on the eigenvector v_1* as a set $\{\mathbf{v}_1, \dots, \mathbf{v}_k\}$ of k generalized eigenvectors such that

$$\begin{aligned}(\mathbf{A} - \lambda\mathbf{I})\mathbf{v}_k &= \mathbf{v}_{k-1} \\(\mathbf{A} - \lambda\mathbf{I})\mathbf{v}_{k-1} &= \mathbf{v}_{k-2} \\&\vdots \\(\mathbf{A} - \lambda\mathbf{I})\mathbf{v}_2 &= \mathbf{v}_1.\end{aligned}$$

Now, a fundamental theorem from linear algebra states⁵ that every $n \times n$ matrix \mathbf{A} has n linearly independent generalized eigenvectors. These n generalized eigenvectors may be arranged in chains, with the sum of the lengths of the chains associated with a given eigenvalue λ equal to the multiplicity of λ (a.k.a. the order of the root in the characteristic equation). However, the structure of these chains can be quite complicated.

The idea behind how we build our solutions is that we calculate the defect d of our eigenvalue λ and we figure out a solution to the equation:

$$(\mathbf{A} - \lambda\mathbf{I})^{d+1}\mathbf{u} = \mathbf{0}.$$

We then successively multiply by the matrix $(\mathbf{A} - \lambda\mathbf{I})$ until the zero vector is obtained. The sequence gives us a chain of generalized eigenvectors, and from these we build up solutions as follows (assuming there are k generalized eigenvectors in the chain):

$$\begin{aligned}\mathbf{x}_1(t) &= \mathbf{v}_1 e^{\lambda t} \\ \mathbf{x}_2(t) &= (\mathbf{v}_1 t + \mathbf{v}_2) e^{\lambda t} \\ \mathbf{x}_3(t) &= \left(\frac{1}{2} \mathbf{v}_1 t^2 + \mathbf{v}_2 t + \mathbf{v}_3 \right) e^{\lambda t}\end{aligned}$$

⁵This is mathspeak for “A theorem that’s true but we’re not going to prove. So just trust me.”

$$\mathbf{x}_k(t) = \left(\mathbf{v}_1 \frac{t^{k-1}}{(k-1)!} + \cdots + \mathbf{v}_{k-2} \frac{t^2}{2!} + \mathbf{v}_{k-1} t + \mathbf{v}_k \right) e^{\lambda t}$$

We then amalgamate all these chains of generalized eigenvectors, and these gives us our complete set of linearly independent solutions.⁶ This always works. We note that in all the examples we're doing we're assuming all our eigenvalues are real, but that assumption isn't necessary. This method works just fine if we have complex eigenvalues, as long as we allow for complex eigenvectors as well.

Example - Find the general solution of the system:

$$\mathbf{x}' = \begin{pmatrix} 0 & 1 & 2 \\ -5 & -3 & -7 \\ 1 & 0 & 0 \end{pmatrix} \mathbf{x}.$$

⁶In practice we'll only be dealing with smaller (2x2, 3x3, maybe a 4x4) systems, so things won't get too awful.

Room for the example problem.

Even MORE room for the example problem.

Notes on Homework Problems

Problem 5.4.1 is a 2×2 system with a repeated eigenvalue. Shouldn't be too hard. As in section 5.2, the problem also asks you to use a computer system or graphing calculator to construct a direction field and typical solution curves for the system. A sketch is fine.

Problems 5.4.8 and 5.4.16 deal with larger systems than problem 5.4.1, but the approach is the same. They're larger, so they're a bit harder, but on the upside you don't have to draw any direction fields!

For problem 5.4.25 you've got to find some chains. You should take some powers of the coefficient matrix. You'll find it's nilpotent, and that should help you a lot in generating these chains!

Problem 5.4.33 investigates what you do when you've got a defective complex root. Please note that there's a typo in the textbook! The equation $\mathbf{v}_2 = [9 \ 0 \ 1 \ i]^T$ should be $\mathbf{v}_2 = [0 \ 0 \ 1 \ i]^T$.