

# Math 2280 - Lecture 34

Dylan Zwick

Spring 2013

Last time we learned how to solve linear ODEs of the form:

$$A(x)y'' + B(x)y' + C(x)y = 0,$$

around ordinary points using power series. Today, we'll learn how to solve them around a specific type of singular point called a *regular singular point*.

Today's lecture corresponds with section 8.3 of the textbook. The assigned problems are:

Section 8.3 - 1, 8, 15, 18, 24

## Regular Singular Points

Last time we looked at how to solve linear ODEs of the form:

$$A(x)y'' + B(x)y' + C(x)y = 0.$$

The first thing we do is rewrite the ODE as:

$$y'' + P(x)y' + Q(x)y = 0,$$

where, of course,

$$P(x) = \frac{B(x)}{A(x)}, \text{ and } Q(x) = \frac{C(x)}{A(x)}.$$

If  $P(x)$  and  $Q(x)$  are analytic around the point  $a$  then we know there are two linearly independent solutions of the form:

$$y(x) = \sum_{n=0}^{\infty} c_n (x - a)^n$$

where the radii of convergence are at least as great as the distance in the complex plane from  $a$  to the nearest singular point of either  $P(x)$  or  $Q(x)$ .

## Ordinary, Regular, and Irregular Points

We first state without proof that either  $P(x)$  and  $Q(x)$  are analytic at  $x = a$  or approach  $\pm\infty$  as  $x \rightarrow a$ .

Now, of course, we must ask what we do if either  $P(x)$  or  $Q(x)$  is not analytic at  $a$ ? It turns out we have methods for dealing with this as long as they're not analytic in the "right way". We'll get into what that means in just a moment.

We'll restrict ourselves to dealing with the case  $a = 0$ , but we note that by just shifting our coordinates this restriction incurs no loss of generality.

Alright. So, we divide singular points into two types: regular singular points, and irregular singular points. A regular singular point is a singular point where, if we rewrite:

$$y'' + P(x)y' + Q(x)y = 0$$

as

$$y'' + \frac{p(x)}{x}y' + \frac{q(x)}{x^2}y = 0$$

the functions  $p(x)$  and  $q(x)$  are analytic. This is the situation upon which we'll focus. We will not discuss how to solve ODEs around irregular singular points, as that is a much more difficult and advanced topic.

*Example* - Determine whether  $x = 0$  is an ordinary point, a regular singular point, or an irregular singular point of the ODE:

$$x^2y'' + (6 \sin x)y' + 6y = 0$$

We again state a fact without proof. If the limits:

$$\lim_{x \rightarrow 0} p(x) \text{ and } \lim_{x \rightarrow 0} q(x)$$

exist, are finite, and are not 0 then  $x = 0$  is a regular singular point. If both limits are 0 then  $x = 0$  may be a regular singular point or an ordinary point. If either limit fails to exist or is  $\pm\infty$  then  $x = 0$  is an irregular singular point. This gives us a useful way for testing if a singular point is regular.

## The Method of Frobenius

Now we'll figure out how to actually solve these ODEs around regular singular points. We start by examining the simplest such ODE:

$$x^2y'' + p_0xy' + q_0y = 0$$

where  $p_0, q_0$  are both constants. This ODE is solved by  $y = x^r$ , where  $r$  satisfies the quadratic:

$$r(r - 1) + p_0r + q_0 = 0.$$

Using this as our starting point, in general we assume our solution has the form:

$$y(x) = x^r \sum_{n=0}^{\infty} c_n x^n.^1$$

This is called a Frobenius series. We want to figure out what this constant  $r$  needs to be. So, assume that we have a solution in this form. In this case we have:

$$y(x) = x^r \sum_{n=0}^{\infty} c_n x^n = \sum_{n=0}^{\infty} c_n x^{n+r},$$

$$y'(x) = \sum_{n=0}^{\infty} c_n (n+r) x^{n+r-1},$$

$$y''(x) = \sum_{n=0}^{\infty} c_n (n+r)(n+r-1) x^{n+r-2}.$$

We substitute these into:

$$x^2y'' + xp(x)y' + q(x)y = 0,$$

where  $p(x)$  and  $q(x)$  are analytic around  $x = 0$ , and so have a power series representation of the form:

---

<sup>1</sup>This is *NOT* a power series if  $r \notin \mathbb{Z}^+$ .

$$p(x) = p_0 + p_1x + p_2x^2 + \dots$$

$$q(x) = q_0 + q_1x + q_2x^2 + \dots$$

Plugging all this stuff in we get:

$$[r(r-1)c_0x^r + (r+1)rc_1x^{r+1} + \dots] + [p_0x + p_1x^2 + \dots] \cdot [rc_0x^{r-1} + (r+1)c_1x^r + \dots] + [q_0 + q_1x + \dots] \cdot [c_0x^r + c_1x^{r+1} + \dots] = 0.$$

If we examine the  $x^r$  term we get, assuming (as we of course always can and should) that  $c_0 \neq 0$ , we get the relation:

$$r(r-1) + p_0r + q_0 = 0.$$

This is called the indicial equation of the ODE, and it must, according to the identity principle, be satisfied for our solution to work. This is, of course, only a necessary condition, and we certainly haven't proven it's sufficient. That's where the next theorem comes in:

**Theorem** - Suppose that  $x = 0$  is a regular singular point of the ODE:

$$x^2y'' + xp(x)y' + q(x)y = 0.$$

Let  $\rho > 0$  denote the minimum of the radii of convergence of the power series:

$$p(x) = \sum_{n=0}^{\infty} p_nx^n \text{ and } q(x) = \sum_{n=0}^{\infty} q_nx^n.$$

Let  $r_1$  and  $r_2$  be the real roots (we'll always be assuming our roots are real), of the indicial equation with  $r_1 \geq r_2$ . Then

1. For  $x > 0$ , there exists a solution to our ODE of the form:

$$y_1(x) = x^{r_1} \sum_{n=0}^{\infty} a_nx^n, \quad a_0 \neq 0,$$

corresponding to the larger root  $r_1$ .

2. If  $r_1 - r_2$  is neither zero nor a positive integer, then there exists a second linearly independent solution for  $x > 0$  of the form:

$$y_2(x) = x^{r_2} \sum_{n=0}^{\infty} b_n x^n, b_0 \neq 0,$$

corresponding to the smaller root  $r_2$ .

The radii of convergence of  $y_1$  and  $y_2$  are at least  $\rho$ . We determine the coefficients by plugging our series into:

$$x^2 y'' + xp(x)y' + q(x)y = 0.$$

*Example* - Use the method of Frobenius to solve the ODE:

$$2x^2 y'' + 3xy' - (x^2 + 1)y = 0$$

around the regular singular point  $x = 0$ .

More room for the example problem, if you need it.

Even MORE room for the example problem, if you need it.