

STABLE COMBINATIONS OF SPECIAL UNIPOTENT REPRESENTATIONS

DAN M. BARBASCH AND PETER E. TRAPA

To Gregg Zuckerman, with respect and admiration

1. INTRODUCTION

In [Ar1]–[Ar2], Arthur outlined a set of conjectures describing the automorphic spectrum of semisimple Lie group over a local field. He suggested that the set of automorphic representations is arranged into (possibly overlapping) packets satisfying a number of properties. In particular, each packet was predicted to give rise to a canonical linear combination of its elements whose character was stable in the sense of Langlands and Shelstad [La], [LaSh].

In the real case, Arthur’s predictions are made precise, refined, and in many cases established in [BV3] and, most completely, in [ABV]. Many of Arthur’s conjectures can be reduced to the case of a certain (precisely defined) set of special unipotent representations. This set is a union of Arthur packets, and since each Arthur packet gives rise to a stable virtual representation, one thus obtains a collection of stable linear combinations of special unipotent representations. One is naturally led to ask if their span exhausts the space of stable virtual special unipotent representations. Simple examples show this is too naive. For example, in the complex case (where stability is empty) Arthur packets are typically not singletons. So the question becomes: can one give a canonical basis of stable virtual special unipotent representations which accounts for these “extra” stable sums?

Under certain natural hypotheses we give a positive answer in terms of the geometry of “special pieces” of nilpotent cone of the Langlands dual Lie algebra. Recall (from [Sp]) that if \mathcal{O}' is a nilpotent adjoint orbit for a complex reductive Lie algebra, there is a unique special orbit \mathcal{O} of smallest dimension which contains \mathcal{O}' in its closure. The collection of all \mathcal{O}' for which \mathcal{O} is this unique orbit is called the special piece of nilpotent cone parametrized by \mathcal{O} . We denote it $\mathcal{SP}(\mathcal{O})$. The special pieces form a partition of the set of nilpotent adjoint orbits indexed by the special orbits.

In order to formulate our main results, like Theorem 2.12 below, a number of technicalities must be treated with care. A significant complication, as in [ABV], is that one cannot work with a single real form individually, but instead must work with an inner class of them simultaneously. (We begin recalling the relevant details in Section 2.) In spite of these technicalities, some consequences of our results are easy to state and have nothing to do with real groups. For example, suppose \mathfrak{g} is a complex semisimple Lie algebra with adjoint group G . Fix a Cartan subalgebra \mathfrak{h} and a system of positive roots $\Delta^+ = \Delta^+(\mathfrak{h}, \mathfrak{g})$, and write $\mathfrak{n} = \bigoplus_{\alpha \in \Delta^+} \mathfrak{g}_\alpha$. Write W for the Weyl group of \mathfrak{h} in \mathfrak{g} . For $w \in W$, write $\mathfrak{n}^w = \bigoplus_{\alpha \in \Delta^+} \mathfrak{g}_{w\alpha}$. Then for each $w \in W$ there is always a dense nilpotent adjoint orbit $\mathcal{O}(w)$ contained in

$$G \cdot (\mathfrak{n} \cap \mathfrak{n}^w).$$

Sometimes the map $w \mapsto \mathcal{O}(w)$ is called the Springer-Steinberg map (since a closely related variant was studied in [St]). A natural question is if there is a canonical section of this map. That is, given a nilpotent adjoint orbit \mathcal{O} , can one canonically define a Weyl group element $w \in W$ such that \mathcal{O} is dense in $G \cdot (\mathfrak{n} \cap \mathfrak{n}^w)$?

For example, suppose \mathcal{O} is even in the sense all of the labels on the associated weighted Dynkin diagram are even. Let $\mathfrak{l} = \mathfrak{l}(\mathcal{O})$ denote the subalgebra of \mathfrak{g} corresponding to the roots labeled zero. If

The first author was partially supported by NSF grant DMS-0300172. The second author was partially supported by NSF grants DMS-0554118 and DMS-0968060.

w_l denotes the long element of the Weyl group of \mathfrak{l} , $W(\mathfrak{l}) \subset W$, then indeed \mathcal{O} is dense in $G \cdot (\mathfrak{n} \cap \mathfrak{n}^{w_l})$. As a consequence of Corollary 4.12 below (applied to the diagonal symmetric subgroup in $G \times G$), we have the following generalization.

Theorem 1.1. *Suppose \mathcal{O} is an adjoint orbit for \mathfrak{g} . Let W^l denote the set of maximal length coset representatives of $W(\mathfrak{l}) \backslash W/W(\mathfrak{l})$ where $\mathfrak{l} = \mathfrak{l}(\mathcal{O})$ corresponds to the nodes labeled zero in the weighted Dynkin diagram for \mathcal{O} . Let $d(\mathcal{O})$ denote the Spaltenstein dual of \mathcal{O} (e.g. [BV3, Appendix B] or [CM, Section 6.3]). Assume that both \mathcal{O} and $d(\mathcal{O})$ are even, and fix an adjoint orbit \mathcal{O}' in $SP(\mathcal{O})$.*

- (a) *There exists a unique element $w(\mathcal{O}') \in W^l$ such that \mathcal{O}' is dense in $G \cdot (\mathfrak{n} \cap \mathfrak{n}^{w(\mathcal{O}')})$. (For example, if $\mathcal{O}' = \mathcal{O}$, then $w(\mathcal{O}') = w_l$, the longest element in the identity coset.)*
- (b) *Let π' denote the Springer representation associated to the trivial local system on \mathcal{O}' , and let sgn denote the sign representation of $W(\mathfrak{l})$. Then*

$$\dim \text{Hom}_{W(\mathfrak{l})}(\text{sgn}, \pi') = 1.$$

Under the conditions of the theorem, the map $\mathcal{O}' \mapsto w(\mathcal{O}')$ in part (a) is thus a natural section of the Springer-Steinberg map. (The equivalence of statements (a) and (b) goes back to Borho-MacPherson. A more general statement is given in Proposition 4.8 below.) It would be interesting to investigate how to relax the evenness hypotheses in the theorem.

The paper is organized as follows. After recalling the machinery of [ABV] in Section 2, we state our main result in Theorem 2.12. We prove it in the final two sections. Examples 2.14, 2.17, and 2.18 illustrate many of the main ideas.

Acknowledgements. We thank Jeffrey Adams for drawing our attention to the problem considered in this paper. In particular, using the software package `atlas` he computed a basis for the space of stable virtual special unipotent representations in many exceptional examples; see www.liegroups.org/tables/unipotent. These examples led us to the formulation of Theorem 2.12.

Finally, it is a pleasure to dedicate this paper to Gregg Zuckerman. His revolutionary ideas, particularly the construction of cohomological induction and his approach to the character theory of real reductive groups (and its relation with tensoring with finite-dimensional representations), are the foundations on which the results in this paper are built.

2. STATEMENT AND EXAMPLES OF THE MAIN RESULTS

Let G be a connected reductive complex algebraic. We begin by fixing a weak extend group G^Γ for G as in [ABV, Definition 2.13]. This means that there is an exact sequence of real Lie groups

$$1 \longrightarrow G \longrightarrow G^\Gamma \longrightarrow \Gamma := \text{Gal}(\mathbb{C}/\mathbb{R}) \longrightarrow 1$$

and each $\delta \in G^\Gamma - G$ acts by conjugation as an antiholomorphic automorphism of G . If $\delta \in G^\Gamma - G$ is such that $\delta^2 \in Z(G)$ — that is if δ is a strong real form for G^Γ in the language of [ABV] — then conjugation by δ defines an antiholomorphic involution of G . In this case, we write $G(\mathbb{R}, \delta)$ for the corresponding fixed points, a real form of G . It follows from [ABV, Proposition 2.14] that the set of real forms which arise in this way constitute exactly one inner class of real forms, and moreover every such inner class arises in this way. In particular, by fixing G^Γ we have fixed an inner class of real forms of G .

Recall (again from [ABV, Definition 2.13]) that a representation of a strong real form for G^Γ is a pair (π, δ) where δ is a strong real form of G^Γ and π is an admissible representation of $G(\mathbb{R}, \delta)$. Two representations (π, δ) and (π', δ') are said to be equivalent if there is an element $g \in G$ such that $\delta' = g\delta g^{-1}$ and π' is infinitesimal equivalent to $\pi \circ \text{Ad}(g^{-1})$. Write $\Pi(G/\mathbb{R})$ for the set of infinitesimal equivalence classes of irreducible representations of strong real forms for G^Γ .

Fix a maximal ideal I in the center of the enveloping algebra $U(\mathfrak{g})$ of the Lie algebra \mathfrak{g} of G . Choose a Cartan subalgebra $\mathfrak{h} \in \mathfrak{g}$ and write W for the Weyl group of \mathfrak{h} in \mathfrak{g} . According to the Harish-Chandra isomorphism we may attach an element $\lambda \in \mathfrak{h}^*/W$ to I . Let $\Pi^\lambda(G/\mathbb{R})$ denote the subset of $\Pi(G/\mathbb{R})$ consisting of those representations whose associated Harish-Chandra modules are annihilated by I . Write $\mathbb{Z}\Pi^\lambda(G/\mathbb{R})$ for the (finite rank) free \mathbb{Z} module with basis indexed by $\Pi^\lambda(G/\mathbb{R})$.

We next introduce various objects on the dual side. Let G^\vee denote the Langlands dual group corresponding to G , and write \mathfrak{g}^\vee for its Lie algebra. Let $\tilde{G}_{\text{alg}}^\vee$ denote the algebraic universal cover of G^\vee (e.g. [ABV, Definition 1.18]). For later use, recall that the construction of the dual group specifies a Cartan subalgebra \mathfrak{h}^\vee of \mathfrak{g}^\vee which is canonically isomorphic to \mathfrak{h}^* .

Definition 1.8 and Lemma 1.9 of [ABV] introduce a smooth complex algebraic variety $X = X(G^\Gamma)$ attached to the extended group fixed above, and provide an action of $\tilde{G}_{\text{alg}}^\vee$ on X which factors to an action of G^\vee . (To be more precise, [ABV, Definition 1.8] explains how to define X from an L -group, and the discussion around [ABV, Proposition 4.14] explains how to build an L -group from a fixed inner class of real forms, in particular the class specified by our fixed weak extended group G^Γ .) The variety X is a disjoint union of smooth (possibly empty) finite-dimensional varieties X^λ indexed by $\lambda \in \mathfrak{h}^*/W$. The action of G^\vee (and $\tilde{G}_{\text{alg}}^\vee$) on X preserves each X^λ . The orbits for both actions on X^λ are the same and are finite in number. We do not recall the general structure of X^λ here, but instead describe certain special cases in detail below.

Let $\mathcal{P}(X^\lambda, \tilde{G}_{\text{alg}}^\vee)$ denote the category of $\tilde{G}_{\text{alg}}^\vee$ -equivariant perverse sheaves on X^λ , and write $\mathbb{Z}\mathcal{P}(X^\lambda, \tilde{G}_{\text{alg}}^\vee)$ for its integral Grothendieck group. Let $T_{G^\vee}^*(X^\lambda)$ denote the conormal variety for the action of G^\vee on X^λ , namely the subvariety of $T^*(X^\lambda)$ consisting of the unions of the various conormal bundles $T_Q^*(X^\lambda)$ to G^\vee orbits Q on X^λ . (Recall that the orbits of G^\vee and $\tilde{G}_{\text{alg}}^\vee$ are the same.) The characteristic cycle functor gives a map

$$\text{CC} : \mathbb{Z}\mathcal{P}(X^\lambda, \tilde{G}_{\text{alg}}^\vee) \longrightarrow H_{\text{top}}(T_{G^\vee}^*(X^\lambda), \mathbb{Z}) \simeq \bigoplus_{Q \in G^\vee \backslash X^\lambda} \mathbb{Z} \left[\overline{T_Q^*(X^\lambda)} \right].$$

The right-hand side is the top-dimensional integral Borel-Moore homology group of $T_{G^\vee}^*(X^\lambda)$ which, as indicated, is isomorphic to the direct sum of the \mathbb{Z} span of the fundamental classes of closures of the individual conormal bundles.

The ABV interpretation of the Local Langlands Conjecture, summarized in [ABV, Corollary 1.26], provides a \mathbb{Z} -module isomorphism

$$\Phi : \mathbb{Z}\Pi^\lambda(G/\mathbb{R}) \simeq \left(\mathbb{Z}\mathcal{P}(X^\lambda, \tilde{G}_{\text{alg}}^\vee) \right)^*$$

for each $\lambda \in \mathfrak{h}^*/W$; here and elsewhere $(\cdot)^*$ applied to a \mathbb{Z} -module denotes $\text{Hom}_{\mathbb{Z}}(\cdot, \mathbb{Z})$. The isomorphism Φ depends on more data than just the weak extended group G^Γ fixed above. It requires fixing a (strong) extended group (G^Γ, \mathcal{W}) as in [ABV, Definition 1.12] and a strong real form [ABV, Definition 1.13]. We define

$$(2.1) \quad \mathbb{Z}_{\text{st}}\Pi^\lambda(G/\mathbb{R}) := \Phi^{-1} \left(\mathbb{Z}\mathcal{P}(X^\lambda, \tilde{G}_{\text{alg}}^\vee) / \ker(\text{CC}) \right)^*.$$

This is a space of integral linear combinations of irreducible representations of G^Γ , that is virtual representations. (This space depends only on the weak extended group G^Γ .)

For the purpose of this paper, we may take (2.1) as the definition of the subspace of stable virtual characters in $\mathbb{Z}\Pi^\lambda(G/\mathbb{R})$. The equivalence with Langlands' original formulation of stability is given in [ABV, Chapter 18].

The main aim of this paper is to define a canonical basis of $\mathbb{Z}_{\text{st}}\Pi^\lambda(G/\mathbb{R})$ (in certain special cases) indexed by rational forms of special pieces of the nilpotent cone of \mathfrak{g}^\vee . We now specify the special

cases of interest. Begin by fixing a nilpotent adjoint orbit \mathcal{O}^\vee for \mathfrak{g}^\vee . Choose a Jacobson-Morozov triple $\{e^\vee, f^\vee, h^\vee\}$ with $h^\vee \in \mathfrak{h}^\vee$ (\mathfrak{h}^\vee as defined above). Set

$$(2.2) \quad \lambda(\mathcal{O}^\vee) = \frac{1}{2}h^\vee \in \mathfrak{h}^\vee \simeq \mathfrak{h}^*.$$

Define

$$(2.3) \quad \mathfrak{l}^\vee(\mathcal{O}^\vee) = \text{the centralizer in } \mathfrak{g}^\vee \text{ of } \lambda(\mathcal{O}^\vee);$$

equivalently $\mathfrak{l}^\vee(\mathcal{O}^\vee)$ is the sum of the zero eigenspaces of $\text{ad}(h^\vee)$. Set

$$(2.4) \quad \mathfrak{p}^\vee(\mathcal{O}^\vee) = \text{the sum of the non-negative eigenspaces of } \text{ad}(h^\vee).$$

Let $I(\mathcal{O}^\vee)$ denote the maximal ideal in the center of $U(\mathfrak{g})$ corresponding to $\lambda(\mathcal{O}^\vee)$ under the Harish-Chandra isomorphism. According to a result of Dixmier [Di], there is a unique maximal primitive ideal $J(\mathcal{O}^\vee)$ in $U(\mathfrak{g})$ containing $I(\mathcal{O}^\vee)$. We say a representation (δ, π) of G^Γ is *special unipotent attached to \mathcal{O}^\vee* if the Harish-Chandra module of π is annihilated by $J(\mathcal{O}^\vee)$. We write

$$\Pi(\mathcal{O}^\vee) \subset \Pi^{\lambda(\mathcal{O}^\vee)}(G/\mathbb{R})$$

for the subset of irreducible special unipotent representations of G^Γ attached to \mathcal{O}^\vee , write $\mathbb{Z}\Pi(\mathcal{O}^\vee)$ for their span, and define

$$\mathbb{Z}_{\text{st}}\Pi(\mathcal{O}^\vee) := \mathbb{Z}\Pi(\mathcal{O}^\vee) \cap \mathbb{Z}_{\text{st}}\Pi^\lambda(G/\mathbb{R}).$$

It is this space for which we will find a canonical basis under certain natural hypotheses.

To state our main results, we need more detailed information about the structure of the G^\vee action on X^λ assuming λ is integral. Let

$$(2.5) \quad Y^\vee = \text{the variety of parabolic subalgebras of } \mathfrak{g}^\vee \text{ conjugate to } \mathfrak{p}^\vee(\mathcal{O}^\vee)$$

with notation as in (2.4). Proposition 6.16 of [ABV] provides a collection of symmetric subgroups $K_1^\vee, \dots, K_k^\vee$ of G^\vee . Each K_i^\vee acts on Y^\vee with finitely many orbits. Furthermore, [ABV, Proposition 7.14] implies the existence of an isomorphism

$$(2.6) \quad \mathcal{P}(X^\lambda, G^\vee) \simeq \mathcal{P}(Y^\vee, K_1^\vee) \oplus \dots \oplus \mathcal{P}(Y^\vee, K_k^\vee),$$

where $\mathcal{P}(Y^\vee, K_i^\vee)$ once again denote the category of K_i^\vee equivariant perverse sheaves on Y^\vee . Moreover, if we let CC_i denote the characteristic cycle functor for $\mathcal{P}(Y^\vee, K_i^\vee)$, then the isomorphism in (2.6) descends to an isomorphism

$$(2.7) \quad \mathcal{P}(X^\lambda, G^\vee)/\ker(\text{CC}) \simeq \mathcal{P}(Y^\vee, K_1^\vee)/\ker(\text{CC}_1) \oplus \dots \oplus \mathcal{P}(Y^\vee, K_k^\vee)/\ker(\text{CC}_k).$$

General properties of the characteristic cycle construction imply that it is insensitive to central extensions of the group acting in the sense that

$$\mathcal{P}(X^\lambda, \tilde{G}_{\text{alg}}^\vee)/\ker(\text{CC}) \simeq \mathcal{P}(X^\lambda, G^\vee)/\ker(\text{CC});$$

see [Ch, Proposition 2.6.2], for example. Thus (2.7) in fact gives

$$(2.8) \quad \mathcal{P}(X^\lambda, \tilde{G}_{\text{alg}}^\vee)/\ker(\text{CC}) \simeq \mathcal{P}(Y^\vee, K_1^\vee)/\ker(\text{CC}_1) \oplus \dots \oplus \mathcal{P}(Y^\vee, K_k^\vee)/\ker(\text{CC}_k).$$

As a matter of notation, we let \mathfrak{k}_i^\vee denote the Lie algebra of K_i^\vee and write

$$(2.9) \quad \mathfrak{g}^\vee = \mathfrak{k}_i^\vee \oplus \mathfrak{s}_i^\vee$$

for the corresponding Cartan decomposition. According to [KR], if \mathcal{O}^\vee is any nilpotent adjoint orbit in \mathfrak{g}^\vee , then each K_i^\vee acts with finitely many orbits on $\mathcal{O}^\vee \cap \mathfrak{s}_i^\vee$,

$$\#(K_i^\vee \backslash (\mathcal{O}^\vee \cap \mathfrak{s}_i^\vee)) < \infty.$$

Recall that an orbit \mathcal{O}^\vee for \mathfrak{g}^\vee is said to be even, if the eigenvalues of $\text{ad}(h^\vee)$ acting on \mathfrak{g}^\vee are all even integers; equivalently if $\lambda(\mathcal{O}^\vee)$ is integral. Assume this is the case and fix an orbit

$$\mathcal{O}_K^\vee \in \bigcup_{i=1}^k K_i^\vee \backslash (\mathcal{O}^\vee \cap \mathfrak{s}_i^\vee).$$

Chapter 27 of [ABV] defines an Arthur packet parametrized by \mathcal{O}_K^\vee ,

$$A(\mathcal{O}_K^\vee) \subset \Pi(\mathcal{O}^\vee).$$

The union of the various Arthur packets (over all possible orbits \mathcal{O}_K^\vee) exhausts $\Pi(\mathcal{O}^\vee)$ (but the union is not in general disjoint). Moreover, for each \mathcal{O}_K^\vee , the discussion around [ABV, (1.34c)] defines a stable integral linear combination of elements of $\Pi(\mathcal{O}_K^\vee)$,

$$(2.10) \quad \pi(\mathcal{O}_K^\vee) \in \mathbb{Z}_{\text{st}}\Pi(\mathcal{O}^\vee).$$

These virtual representations are all linearly independent, so in particular one has

$$(2.11) \quad \dim_{\mathbb{Z}} \mathbb{Z}_{\text{st}}\Pi(\mathcal{O}^\vee) \geq \sum_i \#(K_i^\vee \backslash (\mathcal{O}^\vee \cap \mathfrak{s}_i^\vee)).$$

Our main result finds other interesting stable representations attached to K^\vee orbits on the special piece parametrized by \mathcal{O}^\vee and (in favorable instances) proves they are a basis of $\mathbb{Z}_{\text{st}}(\mathcal{O}^\vee)$.

Theorem 2.12. *Let G be a connected reductive algebraic group with dual group G^\vee . Fix a weak extended group G^Γ for G (in particular, an inner class of real forms for G). Fix an even nilpotent adjoint orbit \mathcal{O}^\vee for \mathfrak{g}^\vee . Assume further that the Spaltenstein dual $\mathcal{O} := d(\mathcal{O}^\vee)$, a nilpotent adjoint orbit for \mathfrak{g} (cf. [BV3, Appendix B]), is also even.*

Recall the Cartan decompositions of (2.9) and the corresponding symmetric subgroups K_i^\vee introduced above. Write $\mathcal{SP}(\mathcal{O}^\vee)$ for the special piece of the nilpotent cone of \mathfrak{g}^\vee containing \mathcal{O}^\vee . Then

$$(2.13) \quad \dim_{\mathbb{Z}} \mathbb{Z}_{\text{st}}\Pi(\mathcal{O}^\vee) = \sum_i \#(K_i^\vee \backslash (\mathcal{SP}(\mathcal{O}^\vee) \cap \mathfrak{s}_i^\vee));$$

cf. (2.11). In fact, for each element \mathcal{O}_K^\vee on the right-hand side, equation (4.13) below defines an element $\pi(\mathcal{O}_K^\vee) \in \mathbb{Z}_{\text{st}}\Pi(\mathcal{O}^\vee)$ so that

$$\left\{ \pi(\mathcal{O}_K^\vee) \mid \mathcal{O}_K^\vee \in \bigcup_i K_i^\vee \backslash (\mathcal{SP}(\mathcal{O}^\vee) \cap \mathfrak{s}_i^\vee) \right\}$$

is a basis of $\mathbb{Z}_{\text{st}}\Pi(\mathcal{O}^\vee)$. (When $G^\vee \cdot \mathcal{O}_K^\vee = \mathcal{O}^\vee$, $\pi(\mathcal{O}_K^\vee)$ coincides with the stable virtual representation in (2.10).)

Example 2.14. Suppose $G = \text{Sp}(4, \mathbb{C})$ and G^Γ gives rise to the inner class of G containing the split form. There are four equivalence classes of strong real forms for G^Γ , $\{\delta_s, \delta_{2,0}, \delta_{1,1}, \delta_{0,2}\}$. The labeling is arranged so that $G(\mathbb{R}, \delta_s) = \text{Sp}(4, \mathbb{R})$ and $G(\mathbb{R}, \delta_{p,q}) = \text{Sp}(p, q)$.

Let \mathcal{O}^\vee denote the (even) nilpotent orbit for $G^\vee = \text{SO}(5, \mathbb{C})$ whose Jordan type is given by the partition 311. Then $d(\mathcal{O}^\vee)$ is the orbit for G with Jordan type corresponding to the partition 22, which is also even, so Theorem 2.12 applies.

In [ABV, Example 27.14], the elements of $\Pi(\mathcal{O}^\vee)$ are enumerated. Among them are eight representations of $\text{Sp}(4, \mathbb{R})$ and one of $\text{Sp}(1, 1)$. The representations of $\text{Sp}(4, \mathbb{R})$ are the three irreducible constituents of $\text{Ind}_{GL(2, \mathbb{R})}^{\text{Sp}(4, \mathbb{R})}(\det)$; the three irreducible constituents of $\text{Ind}_{GL(2, \mathbb{R})}^{\text{Sp}(4, \mathbb{R})}(|\det|)$; and the two irreducible constituents of $\text{Ind}_{GL(1, \mathbb{R}) \times \text{Sp}(2, \mathbb{R})}^{\text{Sp}(4, \mathbb{R})}(\text{sgn}(\det) \otimes 1)$. These eight representations are distinguished by their lowest $\text{U}(2)$ types which in the respective cases are $(2, 0)$, $(0, 2)$, and $(0, 0)$; $(1, 1)$, $(-1, -1)$, and $(1, -1)$; and $(1, 0)$ and $(0, -1)$. Write $\pi_s(m, n)$ for the corresponding special unipotent representation of $\text{Sp}(4, \mathbb{R})$ with lowest $\text{U}(2)$ type (m, n) . Meanwhile the unique special unipotent representation of $\text{Sp}(1, 1)$ attached to \mathcal{O}^\vee is the irreducible spherical representation with infinitesimal character $\lambda(\mathcal{O}^\vee)$ which we denote by $\pi_{(1,1)}(0)$.

The symmetric subgroups K_i^\vee above in this case are

$$K_i^\vee = \text{S}(\text{O}(5-i) \times \text{O}(i)) \text{ for } i = 0, 1, 2.$$

In terms of the signed tableau parametrization (e.g. [CM, Chapter 9]), $\bigcup_i K_i^\vee \backslash (\mathcal{O}^\vee \cap \mathfrak{s}^\vee)$ consists of three elements

$$\begin{array}{ccc} + & - & + \\ + & & \\ + & & \end{array}, \begin{array}{ccc} - & + & - \\ + & & \\ + & & \end{array}, \begin{array}{ccc} + & - & + \\ + & & \\ - & & \end{array};$$

the first arises for $i = 1$, the second and third for $i = 2$. This means there are three Arthur packets in $\Pi(\mathcal{O}^\vee)$. They are listed in [ABV, (27.17)]. They give rise, respectively, to the following three stable virtual representations in $\mathbb{Z}_{\text{st}}(\mathcal{O}^\vee)$,

$$\begin{aligned} & \pi_s(1, 0) + \pi_s(0, -), \\ & \pi_s(0, 0) + \pi_s(1, -1), \\ & \pi_s(1, 1) + \pi_s(-1, -1) + \pi_s(2, 2) + \pi_s(-2, -2) + \pi_{(1,1)}(0). \end{aligned}$$

Meanwhile there is another orbit $\mathcal{O}^{\vee'}$ (besides \mathcal{O}^\vee) in $\mathcal{SP}(\mathcal{O}^\vee)$, namely the orbit with Jordan type corresponding to the partition 221. This time $\bigcup_i K_i^\vee \backslash (\mathcal{O}^{\vee'} \cap \mathfrak{s}^\vee)$ consists of one element

$$(2.15) \quad \begin{array}{cc} + & - \\ - & + \\ + & \end{array}$$

arising for K_2^\vee . Theorem 2.12 thus implies $\dim_{\mathbb{Z}} \mathbb{Z}_{\text{st}}(\mathcal{O}^\vee) = 3 + 1$, and gives an additional stable virtual representation parametrized by the orbit in (2.15). This extra stable sum is

$$(2.16) \quad \pi_s(1, 1) + \pi_s(-1, -1) - \pi_s(2, 2) - \pi_s(-2, -2).$$

Example 2.17. Let G be of type F4, and let \mathcal{O}^\vee be the orbit labeled $F4(a3)$ in the Bala-Carter classification (e.g. [CM, Section 8.4]). If we orient the Dynkin diagram of F4 so that the first two roots are long, the weighted Dynkin diagram for \mathcal{O}^\vee is 0200. In particular, the orbit is even. In fact \mathcal{O}^\vee is equal to its own Spaltenstein dual, and thus Theorem 2.12 applies.

The special piece $\mathcal{SP}(\mathcal{O}^\vee)$ consists of four other orbits besides $F4(a3)$. In the Bala-Carter classification, they are labeled $C3(a1)$, $\widetilde{A2} + A1$, $B2$, and $A2 + \widetilde{A1}$. The respective weighted Dynkin diagrams are 1010, 0101, 2001, and 0010.

There is a unique inner class of real forms for G ; it contains the split, rank one, and compact forms. (In fact it is easy to see (from the singularity of the infinitesimal character $\lambda(\mathcal{O}^\vee)$ that $\Pi(\mathcal{O}^\vee)$ can consist of representations only of the split form.) The only symmetric subgroup K^\vee appearing above in this case is the quotient of $\text{Sp}(6, \mathbb{C}) \times \text{SL}(2, \mathbb{C})$ by the diagonal copy of a central $\mathbb{Z}/2$. From the tables in [CM, Section 9.5], we deduce that

$$\#K^\vee \backslash (F4(a3) \cap \mathfrak{s}^\vee) = 3$$

and so there are three Arthur packets in $\Pi(\mathcal{O}^\vee)$. Meanwhile we have

$$\begin{aligned} \#K^\vee \backslash (C3(a1) \cap \mathfrak{s}^\vee) &= 2 \\ \#K^\vee \backslash ((\widetilde{A2} + A1) \cap \mathfrak{s}^\vee) &= 1 \\ \#K^\vee \backslash (B2 \cap \mathfrak{s}^\vee) &= 2 \\ \#K^\vee \backslash ((A2 + \widetilde{A1}) \cap \mathfrak{s}^\vee) &= 1. \end{aligned}$$

Thus Theorem 2.12 says

$$\dim_{\mathbb{Z}} \mathbb{Z}_{\text{st}} \Pi(\mathcal{O}^\vee) = 3 + 2 + 1 + 2 + 1.$$

The definition in (4.13) gives a canonical basis for the space. To write the basis down explicitly requires computing characteristic cycles of irreducible objects in $\mathcal{P}(K^\vee, Y^\vee)$. We have not performed the calculations required to do this.

Example 2.18. Let G be of type E8, and let \mathcal{O}^\vee denote the orbit labeled $E8(a7)$ in the Bala-Carter classification. It is even and self-dual, and thus Theorem 2.12 applies. The special piece it parametrizes consists of the additional orbits $E7(a5)$, $E6(a3) + A1$, $D6(a2)$, $D5(a1) + A2$, $A5 + A1$, and $A4 + A3$.

There is a unique inner class of real forms for G , and (arguing as in the previous example), $\Pi(\mathcal{O}^\vee)$ can consist of representations only of the split form. The only symmetric subgroup K^\vee appearing above in this case is a quotient of $\text{Spin}(16, \mathbb{C})$ by a central $\mathbb{Z}/2$ (but K^\vee is not isomorphic to $\text{SO}(16, \mathbb{C})$). Again using the tables in [CM, Section 9.5], we deduce that

$$\#K^\vee \backslash (E8(a7) \cap \mathfrak{s}^\vee) = 3$$

and so there are three Arthur packets in $\Pi(\mathcal{O}^\vee)$. Meanwhile we have

$$\begin{aligned} \#K^\vee \backslash (E7(a5) \cap \mathfrak{s}^\vee) &= 2 \\ \#K^\vee \backslash ((E6(a3) + A1) \cap \mathfrak{s}^\vee) &= 2 \\ \#K^\vee \backslash (D6(a2) \cap \mathfrak{s}^\vee) &= 2 \\ \#K^\vee \backslash ((A5 + A1) \cap \mathfrak{s}^\vee) &= 1 \\ \#K^\vee \backslash ((D5(a1) + A2) \cap \mathfrak{s}^\vee) &= 1 \\ \#K^\vee \backslash ((A4 + A3) \cap \mathfrak{s}^\vee) &= 1. \end{aligned}$$

Thus Theorem 2.12 implies

$$\dim_{\mathbb{Z}} \mathbb{Z}_{\text{st}} \Pi(\mathcal{O}^\vee) = 3 + 2 + 2 + 2 + 1 + 1 + 1.$$

3. PROOF OF EQUALITY IN (2.13)

Our main technique allows us to compute the numbers in (2.13) in terms of certain Weyl group calculations. The full Weyl group does not act at singular infinitesimal character, and so we must instead translate to regular infinitesimal character and work there.

Retain the setting of Theorem 2.12. Temporarily choose a system of simple roots for \mathfrak{h} in \mathfrak{g} and a representative λ_\circ of $\lambda(\mathcal{O}^\vee)$ which is dominant. Let μ be the highest weight of a finite-dimensional representation of G such that $\nu_\circ := \lambda_\circ + \mu \in \mathfrak{h}^*$ is dominant and regular. Let ν denote the image of ν_\circ in \mathfrak{h}^*/W . As above, we can consider the set $\Pi^\nu(G/\mathbb{R})$ and its \mathbb{Z} span $\mathbb{Z}\Pi^\nu(G/\mathbb{R})$. This space identifies with an appropriate Grothendieck group of representations at regular integral infinitesimal character which admits a coherent continuation action of W .

Recall the symmetric subgroups $K_1^\vee, \dots, K_k^\vee$ of the previous section. Let X^\vee denote the full flag variety for \mathfrak{g}^\vee . There is an action of W on each Grothendieck group $\mathbb{Z}\mathcal{P}(X^\vee, K_i^\vee)$. (One way to see this is to use the Riemann-Hilbert correspondence to identify $\mathcal{P}(X^\vee, K_i^\vee)$ with a category of K_i^\vee equivariant holonomic \mathcal{D} modules on X^\vee . In turn, by localization, this category is a equivalent to a category of \mathfrak{g}^\vee modules which admits a coherent continuation action of W .)

Meanwhile Corollary 1.26 and Proposition 7.14 of [ABV] give an isomorphism (depending on the extended group (G^Γ, \mathcal{W}))

$$\Psi : \mathbb{Z}\Pi^\nu(G/\mathbb{R}) \longrightarrow \bigoplus_i \mathbb{Z}\mathcal{P}(X^\vee, K_i^\vee)^*$$

which intertwines the W action on both sides. Once again we have characteristic cycle functors

$$\text{CC}_i : \mathbb{Z}\mathcal{P}(X^\vee, K_i^\vee) \longrightarrow \text{H}_{\text{top}} \left(T_{K_i^\vee}^*(X^\vee), \mathbb{Z} \right) \simeq \bigoplus_{Q \in K_i^\vee \backslash X^\vee} \left[\overline{T_{K_i^\vee}^*(X^\vee)} \right].$$

We have remarked that the domain of CC_i carries an action of W . The range does as well, and according to results of Tanisaki [Ta], each CC_i is W -equivariant. Thus

$$\mathbb{Z}\mathcal{P}(X^\vee, K_i^\vee) / \ker(\text{CC}_i) \simeq \text{H}_{\text{top}} \left(T_{K_i^\vee}^*(X^\vee), \mathbb{Z} \right)$$

as representations of W . Once again we define

$$\mathbb{Z}_{\text{st}}\Pi^\nu(G/\mathbb{R}) := \Psi^{-1} \left(\bigoplus_i \mathbb{Z}\mathcal{P}(X^\vee, K_i^\vee)/\ker(\text{CC}_i) \right)^\star.$$

and we have an isomorphism

$$(3.1) \quad \mathbb{Z}_{\text{st}}\Pi^\nu(G/\mathbb{R}) \simeq \bigoplus_i \text{H}_{\text{top}} \left(T_{K_i^\vee}^*(X^\vee), \mathbb{Z} \right)^\star$$

of representations of W .

For our counting argument, we need to specify a particular left cell representation. Let $\Gamma^\vee(\mathcal{O}^\vee)$ denote the centralizer in \mathfrak{g}^\vee of $\lambda(\mathcal{O}^\vee) \in \mathfrak{h}^\vee$, and let $w(\mathcal{O}^\vee)$ denote the long element of the Weyl group of $\Gamma^\vee(\mathcal{O}^\vee)$ viewed as an element of $W(\mathfrak{h}^\vee, \mathfrak{g}^\vee) = W$. Let $V(\mathcal{O}^\vee)$ denote the representation of W afforded by the integral linear combinations of elements of the Kazhdan-Lusztig left cell containing $w(\mathcal{O}^\vee)$.

Proposition 3.2 ([BV3, Section 5]). *Retain the setting above. In particular, assume \mathcal{O}^\vee is even. We have*

$$\dim_{\mathbb{Z}} \mathbb{Z}\Pi(\mathcal{O}^\vee) = \dim \text{Hom}_W(V(\mathcal{O}^\vee) \otimes \text{sgn}, \mathbb{Z}\Pi^\nu(G/\mathbb{R})).$$

and

$$\dim_{\mathbb{Z}} \mathbb{Z}_{\text{st}}\Pi(\mathcal{O}^\vee) = \dim \text{Hom}_W(V(\mathcal{O}^\vee) \otimes \text{sgn}, \mathbb{Z}\Pi_{\text{st}}^\nu(G/\mathbb{R})).$$

The following result brings the role of special pieces into play. To state it, we need to introduce some notation for the Springer correspondence. Fix any nilpotent adjoint orbit \mathcal{O} for \mathfrak{g} and a representative x of \mathcal{O} . Let $A_G(\xi)$ denote the component group of the centralizer of x in G . We let $\text{Sp}(x)$ denote the Springer representation of $W \times A_G(x)$ on the top homology of the Springer fiber over x (normalized so that $\text{Sp}(x)$ is the sign representation of W if x is zero). As usual, we set

$$\text{Sp}(x)^{A_G(x)} = \text{Hom}_{A_G(x)}(\mathbb{1}, \text{Sp}(\xi)).$$

This is a representation of W .

Proposition 3.3. *Suppose \mathcal{O}^\vee is an even nilpotent adjoint orbit for \mathfrak{g}^\vee . Let \mathcal{O} denote special nilpotent orbit for \mathfrak{g} obtained as the Spaltenstein dual of \mathcal{O}^\vee . Enumerate representative for the adjoint orbits in the special piece parametrized by \mathcal{O} as x_1, \dots, x_l . Then*

$$V(\mathcal{O}^\vee) \otimes \text{sgn} \simeq \bigoplus_i \text{Sp}(x_i)^{A_G(x_i)}.$$

Proof. This follows by combining [BV3, Proposition 5.28] and [Lu2, Theorem 0.4]. \square

The proposition involves special pieces on the group side, while the statement of Theorem 2.12 involves special pieces on the dual side. If we make the additional hypothesis that \mathcal{O} is even, then we can match up the two sides.

Proposition 3.4. *Suppose \mathcal{O}^\vee is an even nilpotent adjoint orbit for \mathfrak{g}^\vee . Let \mathcal{O} denote its Spaltenstein dual, and further assume that \mathcal{O} is even. Enumerate representative in the special piece parametrized by \mathcal{O}^\vee as $x_1^\vee, \dots, x_l^\vee$. Then*

$$V(\mathcal{O}^\vee) \simeq \bigoplus_i \text{Sp}(x_i^\vee)^{A_{G^\vee}(x_i^\vee)}.$$

That is, $V(\mathcal{O}^\vee)$ is the sum over the orbits in $\mathcal{SP}(\mathcal{O}^\vee)$ of the Weyl group representations attached to the trivial local system on them.

Proof. This follows from the previous proposition and Lusztig's classification of cells [Lu1] (the relevant details of which are recalled in [BV3, Theorem 4.7d]). \square

Proposition 3.2 gives a way to compute the dimension of the left-hand side of (2.13) in terms of Weyl group representations. We need a way to do the same for the right-hand side.

Proposition 3.5. *Suppose K^\vee is a symmetric subgroup of G^\vee and write $\mathfrak{g}^\vee = \mathfrak{k}^\vee \oplus \mathfrak{s}^\vee$ for the corresponding Cartan decomposition. Let $x_1^\vee, x_2^\vee, \dots$, denote representatives of the nilpotent K^\vee orbits on \mathfrak{s}^\vee . Let $A_{K^\vee}(x_i^\vee)$ denote the component group of the centralizer in K^\vee of x_i^\vee . (Since this group maps to $A_{G^\vee}(x_i^\vee)$, it makes sense to consider invariants $\mathrm{Sp}(x_i^\vee)^{A_{K^\vee}(x_i^\vee)}$ in $\mathrm{Sp}(x_i^\vee)$ of the image of $A_{K^\vee}(x_i^\vee)$ in $A_{G^\vee}(x_i^\vee)$.) As W representations, we have*

$$H_{\mathrm{top}}(T_{K^\vee}^*(X^\vee), \mathbb{Z}) \simeq \sum_i \mathrm{Sp}(x_i^\vee)^{A_{K^\vee}(x_i^\vee)}.$$

In particular, each representation of W attached to the trivial local system on a complex nilpotent orbit appears with multiplicity equal to the number of K^\vee orbits on its intersection with \mathfrak{s}^\vee .

Proof. This follows from [Ro, Theorem 3.3]. (Rossmann works with the conormal variety of orbits of real forms of G^\vee on X^\vee . To translate to the conormal variety of orbits for a symmetric subgroup, one can use [MUV], for example.) \square

Proof of equality in (2.13). From Proposition 3.2 and (3.1), we have

$$\begin{aligned} \dim_{\mathbb{Z}} \mathbb{Z}_{\mathrm{st}} \Pi(\mathcal{O}^\vee) &= \dim \mathrm{Hom}_W (V(\mathcal{O}^\vee) \otimes \mathrm{sgn}, \mathbb{Z}_{\mathrm{st}} \Pi^\vee(G/\mathbb{R})) \\ &= \dim \mathrm{Hom}_W \left(V(\mathcal{O}^\vee) \otimes \mathrm{sgn}, \bigoplus_i H_{\mathrm{top}}(T_{K_i^\vee}^* \mathcal{B}, \mathbb{Z})^* \right) \\ &= \dim \mathrm{Hom}_W \left(V(\mathcal{O}^\vee), \bigoplus_i H_{\mathrm{top}}(T_{K_i^\vee}^* \mathcal{B}, \mathbb{Z}) \right). \end{aligned}$$

The concluding sentences of Propositions 3.4 and 3.5 show that the right-hand side equals

$$\sum_i \#(K_i^\vee \setminus (\mathcal{SP}(\mathcal{O}^\vee) \cap \mathfrak{s}_i^\vee))$$

as claimed. \square

4. PROOF OF THEOREM 2.12

In this section we prove the last assertion of Theorem 2.12. According to (2.1),

$$(4.1) \quad \mathbb{Z}_{\mathrm{st}} \Pi(\mathcal{O}^\vee) \subset \mathbb{Z}_{\mathrm{st}} \Pi^{\lambda(\mathcal{O}^\vee)}(G/\mathbb{R}) \simeq \bigoplus_i \left[H_{\mathrm{top}}(T_{K_i^\vee}^*(Y^\vee), \mathbb{Z}) \right]^*$$

Our main task is to determine which linear functionals on

$$(4.2) \quad H_{\mathrm{top}}(T_{K_i^\vee}^*(Y^\vee), \mathbb{Z}) = \bigoplus_{Q \in K_i^\vee \setminus Y^\vee} \left[\overline{T_Q^*(Y^\vee)} \right]$$

correspond to elements of $\mathbb{Z}_{\mathrm{st}} \Pi(\mathcal{O}^\vee)$ in (4.1). This is the content of part (2) of the next proposition. To formulate it, we recall the G^\vee equivariant moment map μ from $T^*(Y^\vee)$ to $(\mathfrak{g}^\vee)^*$. We use an invariant form to identify \mathfrak{g}^\vee and $(\mathfrak{g}^\vee)^*$, and view the image of the moment map in \mathfrak{g}^\vee itself.

Proposition 4.3. *Retain the setting of Theorem 2.12. For each orbit Q of some K_i^\vee on Y^\vee , define*

$$m_Q \in \left[H_{\mathrm{top}}(T_{K_i^\vee}^*(Y^\vee), \mathbb{Z}) \right]^*$$

as the multiplicity of the fundamental class corresponding to the closure of the conormal bundle to Q (c.f. (4.2)). Recall the isomorphism

$$(4.4) \quad \mathbb{Z}_{\mathrm{st}} \Pi^{\lambda(\mathcal{O}^\vee)}(G/\mathbb{R}) \simeq \bigoplus_i \left[H_{\mathrm{top}}(T_{K_i^\vee}^*(Y^\vee), \mathbb{Z}) \right]^*$$

and write $\pi(Q)$ for the element of $\mathbb{Z}_{\mathrm{st}} \Pi^{\lambda(\mathcal{O}^\vee)}(G/\mathbb{R})$ corresponding to m_Q .

(1) *The set*

$$(4.5) \quad \{\pi(Q) \mid Q \in K_i^\vee \setminus Y^\vee \text{ for some } i\}$$

is a basis for

$$\mathbb{Z}_{\text{st}}\Pi^{\lambda(\mathcal{O}^\vee)}(G/\mathbb{R}).$$

(2) *The set*

$$(4.6) \quad \{\pi(Q) \mid \mu(T_Q^*(Y^\vee)) \cap \mathcal{SP}(\mathcal{O}^\vee) \text{ is nonempty}\}$$

is a basis for the subspace

$$\mathbb{Z}_{\text{st}}\Pi(\mathcal{O}^\vee) \subset \mathbb{Z}_{\text{st}}\Pi^{\lambda(\mathcal{O}^\vee)}(G/\mathbb{R}).$$

Proof. Since

$$\bigcup_i \{m_Q \mid Q \in K_i^\vee \setminus Y^\vee\}$$

is obviously a basis for the left-hand side of (4.4) (in light of (4.2)), statement (1) of the proposition is clear.

For the second statement, we begin by proving

$$\{\pi(Q) \mid \mu(T_Q^*(Y^\vee)) \cap \mathcal{SP}(\mathcal{O}^\vee) \text{ is nonempty}\}$$

are linearly independent elements of $\mathbb{Z}_{\text{st}}\Pi(\mathcal{O}^\vee)$. We need some additional notation. For an object \mathcal{S} in $\mathcal{P}(Y^\vee, K_i^\vee)$, write

$$\text{CC}_i(\mathcal{S}) = a_1[\overline{T_{Q_1}^*(Y^\vee)}] + \cdots + a_r[\overline{T_{Q_r}^*(Y^\vee)}],$$

and define

$$\text{AV}_{\mathbb{C}}(\mathcal{S}) = G^\vee \cdot \bigcup_i \mu(\overline{T_{Q_i}^*(Y^\vee)}) \subset \overline{\mathcal{O}^\vee}.$$

According to the irreducibility theorem of [BB1], if \mathcal{S} is irreducible, then $\text{AV}_{\mathbb{C}}(\mathcal{S})$ is the closure of a single adjoint orbit. The results of [BV1, BV2] show that the orbit must be special.

Next recall the isomorphism

$$\Phi : \mathbb{Z}_{\text{st}}\Pi^{\lambda(\mathcal{O}^\vee)}(G/\mathbb{R}) \longrightarrow \bigoplus_i (\mathbb{Z}\mathcal{P}(Y^\vee, K_i^\vee)/\ker(\text{CC}_i))^*$$

obtained from (2.1) and (2.8). It follows from [ABV, Theorem 27.12] that $\pi \in \mathbb{Z}_{\text{st}}\Pi^{\lambda(\mathcal{O}^\vee)}(G/\mathbb{R})$ belongs to the subspace $\mathbb{Z}_{\text{st}}\Pi(\mathcal{O}^\vee)$ if and only if there is an irreducible object \mathcal{S} in some $\mathcal{P}(Y^\vee, K_i^\vee)$ with $\text{AV}_{\mathbb{C}}(\mathcal{S}) = \overline{\mathcal{O}^\vee}$ such that $\Phi(\pi)(\mathcal{S}) \neq 0$. As a consequence, suppose Q is an orbit of K_i^\vee on Y^\vee such that

$$\mu(\overline{T_Q^*(Y^\vee)}) \cap \mathcal{SP}(\mathcal{O}^\vee) \neq \emptyset.$$

Set

$$m'_Q = m_Q \circ \text{CC}_i \in (\mathbb{Z}\mathcal{P}(Y^\vee, K_i^\vee)/\ker(\text{CC}_i))^*.$$

There exists at least one irreducible object \mathcal{S} in $\mathcal{P}(Y^\vee, K_i^\vee)$ whose support is the closure of Q . (For example, let \mathcal{S} be the DGM extension of the trivial local system on Q . The fundamental class of the closure of $T_Q^*(Y^\vee)$ appears with multiplicity one in $\text{CC}_i(\mathcal{S})$.) Since $\mu(\overline{T_Q^*(Y^\vee)}) \cap \mathcal{SP}(\mathcal{O}^\vee)$ is nonempty by hypothesis, and since $\text{AV}_{\mathbb{C}}(\mathcal{S})$ must be the closure of a special orbit, it follows that $\text{AV}_{\mathbb{C}}(\mathcal{S}) = \overline{\mathcal{O}^\vee}$. Therefore, by the discussion above, the element $\pi(Q) \in \mathbb{Z}_{\text{st}}\Pi^{\lambda(\mathcal{O}^\vee)}(G/\mathbb{R})$ corresponding to m_Q is nonzero and belongs to $\mathbb{Z}_{\text{st}}\Pi(\mathcal{O}^\vee)$. In other words,

$$\{\pi(Q) \mid \mu(T_Q^*(Y^\vee)) \cap \mathcal{SP}(\mathcal{O}^\vee) \text{ is nonempty}\} \subset \mathbb{Z}_{\text{st}}\Pi(\mathcal{O}^\vee).$$

Because the m_Q are clearly linearly independent, so are the elements $\pi(Q)$ on the left-hand side above.

It remains to show the elements in (4.6) are indeed a basis of $\mathbb{Z}_{\text{st}}\Pi(\mathcal{O}^\vee)$. Because of the linear independence just established, it suffices to check

$$(4.7) \quad \sum_i \# \{Q \in K_i^\vee \backslash Y^\vee \mid \mu(T_Q^*(Y^\vee)) \cap \mathcal{SP}(\mathcal{O}^\vee)\} \geq \dim_{\mathbb{Z}} \mathbb{Z}_{\text{st}}\Pi(\mathcal{O}^\vee).$$

The following general result will be the main tool we use for counting the left-hand side.

Proposition 4.8. *As above, assume \mathcal{O}^\vee is even (but do not necessarily assume the Spaltenstein dual of \mathcal{O}^\vee is even). With notation as in (2.2)–(2.5), fix a symmetric subgroup K^\vee of G^\vee and write $\mathfrak{g}^\vee = \mathfrak{k}^\vee \oplus \mathfrak{s}^\vee$ for the corresponding Cartan decomposition. Fix a nilpotent K^\vee orbit \mathcal{O}_K^\vee on \mathfrak{s}^\vee . Let $c(\mathcal{O}_K^\vee)$ denote the number of K^\vee orbits Q on Y^\vee such that $\mu(T_Q^*Y^\vee)$ meets \mathcal{O}_K^\vee in a dense open set. Then, for $x^\vee \in \mathcal{O}_K^\vee$,*

$$c(\mathcal{O}_K^\vee) = \dim \text{Hom}_{W(\iota^\vee(\mathcal{O}^\vee))}(\text{sgn}, \text{Sp}(x)^{A_{K^\vee}(x^\vee)})$$

with notation for the Springer correspondence in Proposition 3.5.

Proof. This is a general result (and doesn't have anything to do with the dual group). It follows from Rossmann's theory applied to the partial flag setting. See [CNT, Section 2] for a proof. \square

Proposition 4.9. *In the setting of Proposition 4.8, assume further that the Spaltenstein dual of \mathcal{O}^\vee is even, and*

$$G^\vee \cdot \mathcal{O}_K^\vee \subset \mathcal{SP}(\mathcal{O}^\vee).$$

Then numbers $c(\mathcal{O}_{K^\vee}^\vee)$ appearing in Proposition 4.8 are all nonzero. More precisely,

$$\dim \text{Hom}_{W(\iota^\vee(\mathcal{O}^\vee))}(\text{sgn}, \text{Sp}(x^\vee)^{A_{G^\vee}(x^\vee)}) = 1,$$

and since $A_{K^\vee}(\xi) \rightarrow A_{G^\vee}(\xi)$,

$$(4.10) \quad \dim \text{Hom}_{W(\iota^\vee(\mathcal{O}^\vee))}(\text{sgn}, \text{Sp}(x^\vee)^{A_{K^\vee}(x^\vee)}) \geq 1.$$

Proof. Section 5 of [BV3] shows that

$$\dim \text{Hom}_{W(\iota^\vee(\mathcal{O}^\vee))}(\text{sgn}, U) = 1$$

for an irreducible representation U in the left cell representation $V(\mathcal{O}^\vee)$. So the current proposition follows from Proposition 3.4. \square

We now return to (4.7). By Proposition 4.9,

$$(4.11) \quad \sum_i \# \{Q \in K_i^\vee \backslash Y^\vee \mid \mu(T_Q^*(Y^\vee)) \cap \mathcal{SP}(\mathcal{O}^\vee)\} \geq \sum_i \# (K_i^\vee \backslash (\mathcal{SP}(\mathcal{O}^\vee) \cap \mathfrak{s}_i^\vee)).$$

By (2.13) (proved in the previous section), the right hand side equals $\dim_{\mathbb{Z}} \mathbb{Z}_{\text{st}}\Pi(\mathcal{O}^\vee)$. This proves (4.7), and hence completes the proof of Proposition 4.3. \square

Corollary 4.12. *In the setting of Proposition 4.9, there exists a unique orbit $Q = Q(\mathcal{O}_K^\vee)$ of K^\vee on Y^\vee such that*

$$\mu(\overline{T_Q^*(Y^\vee)}) = \overline{\mathcal{O}_K^\vee}.$$

Proof. The proof of Proposition 4.3 shows that the inequality in (4.11) must be an equality. Hence the inequality in (4.10) is an equality. Hence the number $c(\mathcal{O}_K^\vee)$ in Proposition 4.8 must be 1. This proves the corollary. \square

Proof of Theorem 2.12. In the setting of Theorem 2.12, given $\mathcal{O}_K^\vee \in K_i^\vee \backslash (\mathcal{SP}(\mathcal{O}^\vee) \cap \mathfrak{s}_i^\vee)$ define $Q(\mathcal{O}_K^\vee)$ as in Corollary 4.12. In the notation of Proposition 4.3 set

$$(4.13) \quad \pi(\mathcal{O}_K^\vee) := \pi(Q(\mathcal{O}_K^\vee)).$$

Then by Proposition 4.3(2),

$$\left\{ \pi(\mathcal{O}_K^\vee) \mid \mathcal{O}_K^\vee \in \bigcup_i K_i^\vee \backslash (\mathcal{SP}(\mathcal{O}^\vee) \cap \mathfrak{s}_i^\vee) \right\}$$

is a basis for $\mathbb{Z}_{\text{st}}\Pi(\mathcal{O}^\vee)$. □

REFERENCES

- [ABV] J. Adams, D. Barbasch, and D. A. Vogan, Jr., *The Langlands Classification and Irreducible Characters for Real Reductive Groups*, Progress in Math, Birkhäuser (Boston), **104** (1992).
- [Ar1] J. Arthur, “On some problems suggested by the trace formula,” in *Proceedings of the Special Year in Harmonic Analysis, University of Maryland*, R. Herb, R. Lipsman, and J. Rosenberg, eds., Lecture Notes in Mathematics **1024** (1983), Springer-Verlag (Berlin-Heidelberg-New York).
- [Ar2] J. Arthur, “Unipotent automorphic representations: conjectures,” in *Orbites Unipotentes et Représentations II. Groups p-adiques et Réels, Astérisque*, **171–172** (1989), 13–71.
- [BV1] D. Barbasch and D. A. Vogan, Jr., Primitive ideals and orbital integrals in complex classical groups, *Math. Ann.*, **259** (1982), 350–382.
- [BV2] D. Barbasch and D. A. Vogan, Jr., Primitive ideals and orbital integrals in complex exceptional groups, *J. Algebra*, **80** (1983), 350–382.
- [BV3] D. Barbasch and D. A. Vogan, Jr., Unipotent representations of complex semisimple Lie groups, *Annals of Math. (2)*, **121** (1985), no. 1, 41–110.
- [BB1] W. Borho, J.-L. Brylinski, Differential operators on homogeneous spaces I. Irreducibility of the associated variety for annihilators of induced modules, *Invent. Math.*, **69** (1982), no. 3, 437–476.
- [BB3] W. Borho, J.-L. Brylinski, Differential operators on homogeneous spaces III. Characteristic varieties of Harish-Chandra modules and of primitive ideals, *Invent. Math.*, **80** (1985), no. 1, 1–68.
- [Ch] J.-T. Chang, Asymptotics and characteristic cycles for representations of complex groups, *Compositio Math.*, **88** (1993), no. 3, 265–283.
- [CNT] D. Ciobotaru, K. Nishiyama, P. E. Trapa, Orbits of symmetric subgroups on partial flag varieties, to appear in *Representation Theory, Complex Analysis, and Integral Geometry Vol. 2*, (Eds. B. Kroetz, O. Offen and E. Sayag), Progress in Math, Birkhäuser (Boston).
- [CM] D. Collingwood and W. M. McGovern, *Nilpotent orbits in semisimple Lie algebras*, Van Nostrand Reinhold (New York), 1993.
- [Di] J. Dixmier, Idéaux primitifs dans l’algèbre enveloppante d’une de Lie semisimple complexe, *C. R. Acad. Sci. Paris (A)*, **271** (1970), 134–136.
- [KR] B. Kostant, S. Rallis, On representations associated with symmetric spaces, *Bull. Amer. Math. Soc.*, **75** (1969), 884–888.
- [La] R. P. Langlands, Stable conjugacy: definitions and lemmas, *Canad. J. Math.*, **31** (1979), 700–725.
- [LaSh] R. P. Langlands and D. Shelstad, On the definition of transfer factors, *Math. Ann.*, **278** (1987), 219–271.
- [Lu1] G. Lusztig, *Characters of Reductive Groups over a Finite Field*, Annals of Mathematics Studies, **107** (1987), Princeton University Press (Princeton, NJ).
- [Lu2] G. Lusztig, Notes on unipotent classes, *Asian Journal of Math.*, **1** (1997), no. 1, 194–207.
- [MUV] I. Mirković, T. Uzawa, K. Vilonen, The Matsuki correspondence for sheaves, *Inventiones Math.*, **109** (1992), no. 1, 231–245.
- [Ro] W. Rossmann, Nilpotent orbital integrals and characters, in Operator algebras, unitary representations, enveloping algebras and invariant theory; in honor of Jacques Dixmier, Progress in Math. **92** (1990), Birkhäuser.
- [Sp] N. Spaltenstein, Classes unipotentes et sous-groupes de Borel, Lecture Notes in Math **946** (1982), Springer Verlag (Berlin-Heidelberg-New York).
- [St] R. Steinberg, On the desingularisation of the unipotent variety, *Inventiones Math.*, **36** (1976), pp. 209–224.
- [Ta] T. Tanisaki, Hodge modules, equivariant K-theory and Hecke algebras, *Publ. RIMS*, **23** (1987).

DEPARTMENT OF MATHEMATICS, CORNELL UNIVERSITY, ITHACA, NY 14853, USA
E-mail address: `dmb14@cornell.edu`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF UTAH, SALT LAKE CITY, UT 84112, USA
E-mail address: `ptrapa@math.utah.edu`