To appear in *Differential and Integral Equations* Preprint (1999)

Self-intersections for the surface diffusion and the volume preserving mean curvature flow

Uwe F. Mayer and Gieri Simonett

Abstract

We prove that the surface diffusion flow and the volume preserving mean curvature flow can drive embedded hypersurfaces to self-intersections.

Key words. Surface diffusion, mean curvature, loss of embeddedness, immersed manifold.

AMS subject classifications. 35R35, 35K55, 58G11, 58F39.

1 Introduction

In this paper we consider two geometric evolution laws: the surface diffusion flow and the volume preserving mean curvature flow. We prove that embedded hypersurfaces can be driven to a self-intersection in finite time. This situation is in strict contrast to the behavior of hypersurfaces under the mean curvature flow, where the maximum principle prevents self-intersections.

The surface diffusion flow is a geometric evolution law in which the normal velocity of a moving hypersurface equals the Laplace-Beltrami of the mean curvature. More precisely we assume in the following that Γ_0 is a closed embedded hypersurface in $\mathbb{R}^n$. Then the surface diffusion flow is governed by the law

$$V(t) = \Delta_{\Gamma(t)} H_{\Gamma(t)}\,, \qquad \Gamma(0) = \Gamma_0\,. \tag{1.1}$$

Here $\Gamma = \{\Gamma(t)\,;\, t \geq 0\}$ is a family of smooth immersed orientable hypersurfaces, $V(t)$ denotes the velocity of Γ in the normal direction at time t, while $\Delta_{\Gamma(t)}$ and $H_{\Gamma(t)}$ stand for the Laplace-Beltrami operator and the mean curvature of $\Gamma(t)$, respectively. The volume preserving mean curvature flow is governed by the law

$$V(t) = -(H(t) - \overline{H}(t))\,, \qquad \Gamma(0) = \Gamma_0\,, \tag{1.2}$$

where $\overline{H}(t) := |\Gamma(t)|^{-1} \int_{\Gamma(t)} H(t)\, d\sigma$ denotes the average of the mean curvature.

The evolution laws (1.1) and (1.2) do not depend on the local choice of the orientation. However, if $\Gamma(t)$ is embedded and encloses a region $\Omega(t)$ we always choose the

outer normal, so that $V(t)$ is positive if $\Omega(t)$ grows, and so that $H_{\Gamma(t)}$ is positive if $\Gamma(t)$ is convex with respect to $\Omega(t)$.

The surface diffusion flow (1.1) was first proposed by Mullins [20] to model the dynamics for the motion of the surface of a crystal when all mass transport is by curvature driven diffusion along the surface. It has also been examined in a more general mathematical and physical context by Davì and Gurtin [10], and by Cahn and Taylor [7]. The surface diffusion flow has recently attracted attention by various researchers, see [3, 4, 6, 8, 11, 12, 13, 16, 19, 21]. We refer to [5, 14, 15, 17] for work related to the volume preserving mean curvature flow.

The surface diffusion flow and the volume preserving mean curvature flow evolve hypersurfaces in such a way that the surface area decreases. Moreover, if Γ is embedded then both flows preserve the volume of the region $\Omega(t)$ enclosed by $\Gamma(t)$, see for instance [12, 14]. The results herein show that both flows can force $\Gamma(t)$ to lose embeddedness in order to decrease surface area.

Theorem 1. *Let $0<\beta<1$. There exist a closed embedded hypersurface $\Sigma_0 \in C^{2+\beta}$, a constant $T_0 > 0$, numbers $t_0, t_1 \in (0, T_0]$ with $t_0 < t_1$, and a $C^{2+\beta}$–neighborhood U_0 of Σ_0 such that the surface diffusion flow (1.1) has a unique classical solution $\Gamma = \{\Gamma(t)\,;\, t \in [0, T_0]\}$ for all $\Gamma_0 \in U_0$, and such that $\Gamma(t)$ ceases to be embedded for every $t \in (t_0, t_1)$ and every $\Gamma_0 \in U_0$. Each hypersurface $\Gamma(t)$ is of class C^∞ for $t \in (0, T_0]$ and smooth in $t \in (0, T_0)$.*

It was conjectured in [11] and later proved in [16] that the surface diffusion flow can drive a dumbbell curve of an appropriate shape to a self-intersection. Theorem 1 extends this result considerably: we can handle nonsymmetric hypersurfaces in any dimension, whereas the method of [16] seems restricted to (symmetric) curves. It should be noted that the neighborhood U_0 of Theorem 1 also contains C^∞-hypersurfaces that will be driven to a self-intersection in finite time. Our approach relies on results and techniques in [12].

Theorem 2. *Let $0<\beta<1$. There exist a closed embedded hypersurface $\Sigma_0 \in C^{1+\beta}$, a constant $T_0 > 0$, numbers $t_0, t_1 \in (0, T_0]$ with $t_0 < t_1$, and a $C^{1+\beta}$–neighborhood U_0 of Σ_0 such that the volume preserving mean curvature flow (1.2) has a unique classical solution $\Gamma = \{\Gamma(t)\,;\, t \in [0, T_0]\}$ for all $\Gamma_0 \in U_0$, and such that $\Gamma(t)$ ceases to be embedded for every $t \in (t_0, t_1)$ and every $\Gamma_0 \in U_0$. Each hypersurface $\Gamma(t)$ is of class C^∞ for $t \in (0, T_0]$ and smooth in $t \in (0, T_0)$.*

To the best of our knowledge, Theorem 2 provides the first rigorous proof for the occurrence of self-intersections for the volume preserving mean curvature flow. In particular, we give a proof for an example proposed by Gage [15] who considered a curve similar to our Fig. 3.

2 The surface diffusion flow

In this section we prove Theorem 1. We first introduce some notations. Given an open set $U \subset \mathbb{R}^n$, let $h^s(U)$ denote the little Hölder spaces of order $s > 0$, that is, the closure of $BUC^\infty(U)$ in $BUC^s(U)$, the latter space being the Banach space of all bounded and uniformly Hölder continuous functions of order s. If Σ is a (sufficiently) smooth submanifold of $\mathbb{R}^n$ then the spaces $h^s(\Sigma)$ are defined by means of a smooth atlas for Σ. It is known that $BUC^t(\Sigma)$ is continuously embedded in $h^s(\Sigma)$ whenever $t > s$. In the following, we assume that Σ is a smooth compact closed immersed oriented hypersurface in $\mathbb{R}^n$. Let ν be the unit normal field on Σ commensurable with the chosen orientation. Then we can find $a > 0$ and an open covering $\{U_l \,;\, l = 1, \ldots, m\}$ of Σ such that

$$X_l : U_l \times (-a, a) \to \mathbb{R}^n \,, \qquad X_l(s, r) := s + r\nu(s)$$

is a smooth diffeomorphism onto its image $\mathcal{R}_l := \mathrm{im}(X_l)$, that is,

$$X_l \in \mathit{Diff}^\infty\big(U_l \times (-a, a), \mathcal{R}_l\big) \,, \qquad 1 \le l \le m \,.$$

This can be done by selecting the open sets $U_l \subset \Sigma$ in such a way that they are embedded in $\mathbb{R}^n$ instead of only immersed, and then taking $a > 0$ sufficiently small so that each of the U_l has a tubular neighborhood of radius a. It follows that $\mathcal{R} := \cup \mathcal{R}_l$ consists of those points in $\mathbb{R}^n$ with distance less than a to Σ. Let $\beta \in (0, 1)$ be fixed. Then we choose numbers α, $\beta_0 \in (0, 1)$ with $\alpha < \beta_0 < \beta$. Let

$$W := \{\rho \in h^{2+\beta_0}(\Sigma) \,;\, \|\rho\|_\infty < a\} \,. \tag{2.1}$$

Given any $\rho \in W$ we obtain a compact oriented immersed manifold Γ_ρ of class $h^{2+\beta_0}$ by means of the following construction:

$$\Gamma_\rho := \bigcup_{l=1}^m \mathrm{Im}\left(X_l : U_l \to \mathbb{R}^n \,,\; [s \mapsto X_l(s, \rho(s))]\right) .$$

Thus Γ_ρ is a graph in normal direction over Σ and ρ is the signed distance between Σ and Γ_ρ. On the other hand, every compact immersed oriented manifold Γ that is a smooth graph over Σ and that is contained in $\mathcal{R}$ can be obtained in this way. For convenience we introduce the mapping

$$\theta_\rho \,:\, \Sigma \to \Gamma_\rho, \qquad \theta_\rho(s) := X_l(s, \rho(s)) \text{ for } s \in U_l \,, \quad \rho \in W \,.$$

It follows that θ_ρ is a well-defined global $(2+\beta_0)$–diffeomorphism from Σ onto Γ_ρ. The surface diffusion flow (1.1) can now be expressed as an evolution equation for the distance function ρ over the fixed reference manifold Σ,

$$\partial_t \rho = G(\rho) \,, \qquad \rho(0) = \rho_0 \,. \tag{2.2}$$

Here $G(\rho) := L_\rho \theta_\rho^*(\Delta_{\Gamma_\rho} H_{\Gamma_\rho})$ for $\rho \in h^{4+\alpha}(\Sigma) \cap W$, while Δ_{Γ_ρ} and H_{Γ_ρ} are the Laplace-Beltrami operator and the mean curvature of Γ_ρ, respectively, and $L(\rho)$ is a factor that comes in by calculating the normal velocity in terms of ρ, see [12] for more details. We are now ready to state the following existence result for solutions of (2.2).

Proposition 2.1.

(a) *Let $\sigma \in W$ be given. Then there exist a positive constant $T_0 > 0$ and a neighborhood $W_0 \subset W$ of σ in $h^{2+\beta_0}(\Sigma)$ such that (2.2) has a unique solution*

$$\rho(\cdot, \rho_0) \in C([0, T_0], W) \cap C^\infty((0, T_0) \times \Sigma) \text{ for every } \rho_0 \in W_0 .$$

(b) *The map $[(t, \rho_0) \mapsto \rho(t, \rho_0)]$ defines a smooth local semiflow on W_0.*

(c) $\rho(\cdot, \rho_0) \in C([0, T_0], h^{4+\alpha}(\Sigma)) \cap C^1([0, T_0], h^\alpha(\Sigma))$ *for every* $\rho_0 \in h^{4+\alpha}(\Sigma) \cap W_0$.

Proof. (a) and (b) follow from [12, Theorem 2.2]. Moreover, [12, Lemma 2.1] shows that the mapping $[\rho \mapsto G(\rho)] : h^{4+\alpha}(\Sigma) \cap W \to h^\alpha(\Sigma)$ is smooth and that the derivative is given by $G'(\rho) = P(\rho) + B(\rho)$, where

$$P(\rho) \in L(h^{4+\alpha}(\Sigma), h^\alpha(\Sigma)), \quad B(\rho) \in L(h^{2+\alpha}(\Sigma), h^\alpha(\Sigma)), \quad \rho \in h^{4+\alpha}(\Sigma) \cap W .$$

In the following we fix $\rho \in h^{4+\alpha}(\Sigma) \cap W$. Lemma 2.1 in [12] also shows that $P(\rho)$ generates a strongly continuous analytic semigroup on $h^\alpha(\Sigma)$. A well-known perturbation result, see [1, Theorem I.1.3.1], then implies $G'(\rho) \in L(h^{4+\alpha}(\Sigma), h^\alpha(\Sigma))$ also generates a strongly continuous analytic semigroup on $h^\alpha(\Sigma)$. It is known (see [1, Vol II], for instance) that the little Hölder spaces are stable under the continuous interpolation method [1, 2, 9, 18]. Therefore, the spaces $(h^{4+\alpha}(\Sigma), h^\alpha(\Sigma))$ form a pair of maximal regularity for $G'(\rho)$, see [1, Theorem III.3.4.1] or [2, 9, 18]. Part (c) follows now from maximal regularity results, for instance [2, Theorem 2.7]. $\square$

In order to provide a proof of Theorem 1 we now choose Σ to be any smooth compact closed immersed orientable hypersurface in $\mathbb{R}^n$ such that its image contains the flat $(n-1)$-dimensional disk $U := \{(s, 0) \in \mathbb{R}^{n-1} \times \mathbb{R} \,;\, |s| \leq 1\}$ twice, and with opposite orientations. To be precise, let $i : \Sigma \to \mathbb{R}^n$ be the immersion under consideration, then we ask that

$$i^{-1}(U) = U^+ \cup U^-$$

with $U^+ \cap U^- = \emptyset$ and both U^+ and U^- are flat $(n-1)$-dimensional disks of radius 1. Additionally we ask that $\Sigma \setminus (U^+ \cup U^-)$ is embedded in $\mathbb{R}^n$. Identifying U^+ for the moment with its image U we ask that the normal on U^+ points upwards, that is, $\nu(\cdot)|_{U^+} = e_n$, the n^{th} basis vector of $\mathbb{R}^n$. It follows that $\nu(\cdot)|_{U^-} = -e_n$.



Fig. 1 This is a possible choice of Σ, cut in halves.

Let W be as in (2.1) and let $\sigma \in h^{4+\alpha} \cap W$ have the following local symmetry: $\sigma(-s) = \sigma(s)$ for every $s \in U^{\pm}$. This implies $\partial_j \sigma(0) = 0$ for $1 \leq j \leq n-1$. Observe that $\theta_\sigma(s) = (s, \pm\sigma(s))$ (these are coordinates in $\mathbb{R}^n$) for $s \in U^{\pm}$ and that $\theta_\sigma : U^{\pm} \to \theta_\sigma(U^{\pm})$ is an $h^{4+\alpha}$–diffeomorphism. It is straightforward to compute

$$G(\sigma)|_{U^{\pm}} := L(\rho)\theta_\sigma^* \left(\Delta_{\Gamma_\sigma} H_{\Gamma_\sigma}\right)|_{U^{\pm}}$$

in local coordinates, yielding

$$\begin{aligned}(n-1)G(\sigma)|_{U^{\pm}}(0) &= -\Delta_{n-1}^2 \sigma(0) + \sum_{j,k=1}^{n-1} (\partial_j \partial_k \sigma(0))^2 \Delta_{n-1}\sigma(0) \\ &\quad +2 \sum_{j,k,l=1}^{n-1} \partial_j \partial_k \sigma(0) \partial_j \partial_l \sigma(0) \partial_k \partial_l \sigma(0)\,,\end{aligned}$$

where Δ_{n-1} is the Laplacian in Euclidean coordinates of $\mathbb{R}^{n-1}$ (see [12, Section 2] for more details). We will now specify one more property of σ. We choose $r > 0$ small and we require that $\sigma(s) = |s|^4$ for $s \in U_r^{\pm} = \{s \in U^{\pm}\,;\, |s| < r\}$; if r is small enough then this is compatible with $\sigma \in h^{4+\alpha}(\Sigma) \cap W$. We conclude that

$$G(\sigma)|_{U^{\pm}}(0) = -24 < 0\,. \tag{2.3}$$

It follows from Proposition 2.1 that the evolution equation (2.2) with initial value $\rho(0) = \sigma$ has a unique solution

$$\rho(\cdot, \sigma) \in C([0, T_0], h^{4+\alpha}(\Sigma)) \cap C^1([0, T_0], h^{\alpha}(\Sigma))\,. \tag{2.4}$$

Next we consider the restriction $\rho^{\pm}(t, \sigma)$ on $U^{\pm}$ of the function $\rho(t, \sigma)$, that is, $\rho^{\pm}(t, \sigma) := \rho(t, \sigma)|_{U^{\pm}}$ for $0 \leq t \leq T_0$, and we set $d^{\pm}(t) := \rho^{\pm}(t, \sigma)(0)$, to track the position of the center. It follows from (2.4) that $d^{\pm} \in C^1([0, T_0])$. Moreover, using the local character of G, we conclude that $d^{\pm}$ satisfies the equation

$$(d^{\pm})'(t) = G(\rho(t, \sigma))|_{U^{\pm}}(0) \quad \text{for} \quad 0 \leq t \leq T_0\,, \qquad d^{\pm}(0) = 0\,. \tag{2.5}$$

Equations (2.3)–(2.5) and the mean value theorem yield

$$d^{\pm}(t) = -Mt + \Big(\int_0^1 \big((d^{\pm})'(\tau t) - (d^{\pm})'(0) \big)\, d\tau \Big) t \,, \tag{2.6}$$

where $M := 24$. It follows from (2.6) that there exists a positive constant $\mu > 0$ and an interval $(t_0, t_1) \subset (0, T_0]$ such that $\rho^{\pm}(t, \sigma)(0) = d^{\pm}(t) \leq -\mu$ for $t \in (t_0, t_1)$. By Proposition 2.1(b) we can find a function $\sigma_0 \in W_0$ such that $\Sigma_0 := \Gamma_{\sigma_0}$ is embedded and such that $\Gamma(t) := \Gamma_{\rho(t,\sigma_0)}$ is immersed for at least $t \in (t_0, t_1)$. By employing Proposition 2.1(b) once more we conclude there is a neighborhood $W(\sigma_0) \subset W_0$ of σ_0 in $h^{2+\beta_0}(\Sigma)$ such that Γ_{ρ_0} is still embedded, whereas $\Gamma_{\rho(t,\rho_0)}$ is immersed for $t \in (t_0, t_1)$ and all $\rho_0 \in W(\sigma_0)$. We note that $C^{2+\beta}(\Sigma)$ is contained in $h^{2+\beta_0}(\Sigma)$ with continuous injection $j : C^{2+\beta}(\Sigma) \to h^{2+\beta_0}(\Sigma)$. Hence $U_0 := j^{-1}(W(\sigma_0))$ is an open $C^{2+\beta}$–neighborhood of σ_0 and Theorem 1 follows by choosing $\Sigma_0 := \Gamma_{\sigma_0}$ and $\Gamma_0 := \Gamma_{\rho_0}$ for $\rho_0 \in U_0$. $\square$



Fig. 2 This is half of Γ_0, a surface that loses embeddedness and becomes immersed. The gap might have to be much smaller than depicted.

Remark 2.2. The following is the essence of the construction: Γ_σ is an immersed hypersurface such that its image contains two opposing fourth-order paraboloids touching only at the vertex. The symmetry of Γ_σ is irrelevant. Locally we can compute the initial velocity of Γ_σ, and it is such as to create an overlapping of the paraboloids. A continuity argument then guarantees the same behavior for nearby embedded hypersurfaces, which do exist by construction of Γ_σ. We have chosen a fourth-order paraboloid in order to facilitate the computation of $G(\sigma)|_{U^{\pm}}$. Any other configuration that produces the same sign as in (2.3) will work as well.

3 The volume preserving mean curvature flow

As in the previous section Σ denotes a smooth compact closed immersed orientable hypersurface in $\mathbb{R}^n$ and we define $W := \{\rho \in h^{1+\beta_0}(\Sigma) \,;\, \|\rho\|_\infty < a\}$ for $a > 0$ appropriate. The volume preserving mean curvature flow (1.2) in $\mathcal{R}$ is equivalent to the following evolution equation for the distance function ρ:

$$\partial_t \rho = G(\rho), \qquad \rho(0) = \rho_0 \,, \tag{3.1}$$

where $\rho_0 \in W$ is chosen such that $\Gamma_0 = \Gamma_{\rho_0}$, and where

$$G(\rho) := L(\rho)\Big(\overline{H}_{\Gamma_\rho} - \theta_\rho^* H_{\Gamma_\rho}\Big)\,, \qquad \rho \in h^{2+\alpha}(\Sigma) \cap W\,. \tag{3.2}$$

Here H_{Γ_ρ} is the mean curvature of Γ_ρ and $L(\rho)$ comes from calculating the normal velocity in coordinates of Σ, see [14] for more details. We have the following existence result for solutions of (3.1).

Proposition 3.1.

(a) *Let $\sigma \in W$ be given. Then there exists a positive constant $T_0 > 0$ and a neighborhood $W_0 \subset W$ of σ in $h^{1+\beta_0}(\Sigma)$ such that* (3.1) *has a unique solution*

$$\rho(\cdot,\rho_0) \in C([0,T_0],W) \cap C^\infty((0,T_0)\times\Sigma) \text{ for every } \rho_0 \in W_0\,.$$

(b) *The map $[(t,\rho_0) \mapsto \rho(t,\rho_0)]$ defines a smooth local semiflow on W_0.*

(c) $\rho(\cdot,\rho_0) \in C([0,T_0],h^{2+\alpha}(\Sigma)) \cap C^1([0,T_0],h^\alpha(\Sigma))$ *for every* $\rho_0 \in h^{2+\alpha}(\Sigma)\cap W_0$.

Proof. Part (a) and part (b) follow from the results in [14, Section 2]. To be more precise, in [14] only embedded surfaces are considered. However, a careful analysis of the proof shows that the existence, uniqueness, and semiflow results remain valid for immersed hypersurfaces, provided one defines the mappings X and Φ_ρ of [14, Section 2] by their local analogs as in [12, Section 2]. Part (c) can be established by similar arguments as in the proof of Proposition 2.1. □

We proceed to prove Theorem 2. Our first goal is to construct a suitable reference manifold Σ. We take a positively oriented immersed curve in $\mathbb{R}^2$, such as the one in Fig. 3. The immersed image contains a line segment twice, with opposite orientations, and the image without this line segment is an embedded curve.



Fig. 3 Rotation of this curve in $\mathbb{R}^n$ yields the hypersurface Σ.

For the two-dimensional case this curve will be Σ, while for the higher dimensional case we rotate the curve to generate a hypersurface, as outlined below. Let $[s \mapsto (x(s), y(s))] : [0, L] \to \mathbb{R}^2$ be a parameterization by arc length of the curve and let $S^{n-2} \subset \mathbb{R}^{n-1}$ be the standard $(n-2)$-dimensional unit sphere. Then we set

$$\Sigma = \{(x(s), y(s)\omega) \; ; \; s \in [0, L], \omega \in S^{n-2}\} \, .$$

Let κ_1 denote the scalar curvature of the curve $[s \mapsto (x(s), y(s))]$. Then a standard computation yields the mean curvature of Σ as

$$H = \frac{1}{n-1}\Big(\kappa_1 - (n-2)\frac{x'}{y}\Big) \, . \tag{3.3}$$

Furthermore, the symmetry of Σ can be used to compute the average of H as

$$\overline{H} = \frac{|S^{n-2}|}{|\Gamma|} \int_0^L H(s) y(s) \, ds \, .$$

Using equation (3.3) and setting $d = \min\{y(s)\}$ one derives

$$(n-1)\int_0^L H(s)y(s) \, ds = d \int_0^L \kappa_1(s) \, ds + \int_0^L \kappa_1(s)(y(s) - d) \, ds - (n-2)\int_0^L x'(s) \, ds \, .$$

The theorem of the turning tangents implies that $\int \kappa_1(s) ds = 2\pi$, and hence it is clear that $\overline{H} > 0$ provided d is large enough, which amounts to shifting the curve far enough away from the axis of rotation. By continuity the average of the mean curvature of Γ_ρ is therefore also positive provided $\rho \in W$ is small enough. Finally, it is clear that by construction Σ contains a flat $(n-1)$-dimensional annulus U twice, with opposite orientations, and that Γ_ρ is in fact embedded in $\mathbb{R}^n$ provided $\rho < 0$ on the annulus.

We let $U^\pm$ be the two components of $i^{-1}(U)$, where $i : \Sigma \to \mathbb{R}^n$ is the immersion under consideration, and U is the flat annulus from above. We now choose $\sigma \in h^{2+\alpha}(\Sigma) \cap W$ with $\sigma \equiv 0$ on $U^\pm$ (in fact we could choose $\sigma \equiv 0$ on Σ so that $\Gamma_\sigma = \Sigma$), then by (3.2)

$$G(\sigma)|_{U^\pm} = \overline{H}_{\Gamma_\sigma} > 0 \, . \tag{3.4}$$

Let $\rho(\cdot, \sigma)$ be the unique solution of (3.1) with initial value $\rho(0) = \sigma$ and note that

$$\rho(\cdot, \sigma) \in C([0, T_0], h^{2+\alpha}(\Sigma)) \cap C^1([0, T_0], h^\alpha(\Sigma)) \tag{3.5}$$

due to Proposition 3.1. As in Section 2 we let $\rho^\pm(t, \sigma)$ denote the restriction of $\rho(t, \sigma)$ on $U^\pm$. We set $d^\pm(t) := \rho^\pm(t, \sigma)(s_0, \omega)$ with s_0 any fixed point on the line segment that generated U and any fixed $\omega \in S^{n-2}$. It follows that $d^\pm \in C^1[0, T_0]$ and it is easy to see that $d^\pm$ solves the equation

$$(d^\pm)'(t) = G(\rho(t, \sigma))|_{U^\pm}(s_0, \omega) \quad \text{for} \quad 0 \le t \le T_0 \, , \qquad d^\pm(0) = 0 \, . \tag{3.6}$$

Equations (3.4)–(3.6) and the mean value theorem show that

$$d^{\pm}(t) = Mt + \Big(\int_0^1 \big((d^{\pm})'(\tau t) - (d^{\pm})'(0)\big)\, d\tau \Big) t$$

with $M := \overline{H}_{\Gamma_\sigma}$. Using Proposition 3.1(b) we can choose a function σ_0 in W_0 such that Γ_{σ_0} is embedded and such that $\Gamma_{\rho(t,\sigma_0)}$ ceases to be embedded on a time interval $(t_0, t_1) \subset (0, T_0]$. The idea is that the time derivative of $d^{\pm}$ is positive, and hence so will be $d^{\pm}$ for some later time even if it was initially negative, see Section 2 for more details. According to Proposition 3.1(b) the same behavior will still prevail for ρ_0 in a small enough $h^{1+\beta_0}(\Sigma)$–neighborhood $W(\sigma_0) \subset W_0$ of σ_0. Theorem 2 now follows by setting $U_0 := j^{-1}(W(\sigma_0))$ with $j := C^{1+\beta}(\Sigma) \to h^{1+\beta_0}(\Sigma)$ the natural injection. $\square$

Remark 3.2. The following is the essence of the construction: Γ_σ is an immersed hypersurface such that its image contains an $(n-1)$-dimensional flat piece twice, with opposite orientation, and Γ_σ has a positive average of its mean curvature. The symmetry of Γ_σ is irrelevant. Locally we can compute the initial velocity of Γ_σ, and it is such as to create an overlapping near the two flat pieces of the surface. We can find embedded hypersurfaces as close to Γ_σ as we want. A continuity argument then guarantees the same behavior for nearby embedded hypersurfaces.

References

[1] H. Amann, *Linear and quasilinear parabolic problems. Vol. I*, Birkhäuser, Basel, 1995, *Vol. II,III*, in preparation.

[2] S.B. Angenent, *Nonlinear analytic semiflows*, Proc. Roy. Soc. Edinburgh Sect. A **115** (1990), 91–107.

[3] A.J. Bernoff, A.L. Bertozzi, and T.P. Witelski, *Axisymmetric surface diffusion: Dynamics and stability of self-similar pinch-off*, preprint.

[4] P. Baras, J. Duchon, and R. Robert, *Évolution d'une interface par diffusion de surface*, Comm. Partial Differential Equations **9** (1984), no. 4, 313–335.

[5] L. Bronsard and B. Stoth, *Volume-preserving mean curvature flow as a limit of a nonlocal Ginzburg-Landau equation*, SIAM J. Math. Anal. **28** (1997), 769–807.

[6] J.W. Cahn, C.M. Elliott, and A. Novick-Cohen, *The Cahn–Hilliard equation with a concentration dependent mobility: motion by minus the Laplacian of the mean curvature*, European J. Appl. Math. **7** (1996), 287–301.

[7] J.W. Cahn and J.E. Taylor, *Surface motion by surface diffusion*, Acta metall. mater. **42** (1994), 1045–1063.

[8] B.D. Coleman, R.S. Falk, and M. Moakher, *Space-time finite element methods for surface diffusion with applications to the theory of the stability of cylinders*, SIAM J. Sci. Comput. **17** (1996), 1434–1448.

[9] G. DaPrato and P. Grisvard, *Equations d'évolution abstraites nonlinéaires de type parabolique*, Ann. Mat. Pura Appl. (4) **120** (1979), 329–396.

[10] F. Davì and M.E. Gurtin, *On the motion of a phase interface by surface diffusion*, Z. Angew. Math. Phys. **41** (1990), 782–811.

[11] C.M. Elliott and H. Garcke, *Existence results for geometric interface models for surface diffusion*, Adv. Math. Sci. Appl. **7** (1997), 467–490.

[12] J. Escher, U.F. Mayer, and G. Simonett, *The surface diffusion flow for immersed hypersurfaces*, SIAM J. Math. Anal., to appear.

[13] J. Escher, U.F. Mayer, and G. Simonett, *On the surface diffusion flow*, Proc. Intern. Conf. on Navier–Stokes Equations and Related Problems, TEV/VSP, Vilnius/Utrecht, 1998.

[14] J. Escher and G. Simonett, *The volume preserving mean curvature flow near spheres*, Proc. Amer. Math. Soc., to appear.

[15] M. Gage, *On an area-preserving evolution equation for plane curves, Nonlinear Problems in Geometry*, D.M. DeTurck, editor, Contemp. Math. **51**, AMS, Providence (1986), 51–62.

[16] Y. Giga and K. Ito, *On pinching of curves moved by surface diffusion*, preprint, Hokkaido University, Japan (1997).

[17] G. Huisken, *The volume preserving mean curvature flow*, J. Reine Angew. Math. **382** (1987), 35-48.

[18] A. Lunardi, *Analytic semigroups and optimal regularity in parabolic problems*, Birkhäuser, Basel, 1995.

[19] U.F. Mayer, *Numerical simulations for the surface diffusion flow in three space dimensions*, preprint.

[20] W.W. Mullins, *Theory of thermal grooving*, J. Appl. Phys. **28** (1957), 333–339.

[21] A. Polden, *Curves and surfaces of least total curvature and forth-order flows*, Ph.D. dissertation, Universität Tübingen, Germany (1996).

Department of Mathematics, Vanderbilt University, Nashville, TN 37240, U.S.A.

mayer@math.vanderbilt.edu
simonett@math.vanderbilt.edu