

Sojourn Times of Brownian Sheet

PERIODICA MATH. HUNG., 41(1-2), 187-194 (2000)

D. Khoshnevisan*
University of Utah

R. Pemantle†
Ohio State University

Keyword and Phrases Brownian sheet, arcsine law, Feynman-Kac formula

AMS 1991 Subject Classification 60G60.

*This paper is dedicated to Professor Endré Csáki
on the occasion of his 65th birthday.*

1 Introduction

Let B denote the standard Brownian sheet. That is, B is a centered Gaussian process indexed by \mathbb{R}_+^2 with continuous trajectories and covariance structure

$$\mathbb{E}\{B_s B_t\} = \min\{s_1, t_1\} \times \min\{s_2, t_2\}, \quad s = (s_1, s_2), \quad t = (t_1, t_2) \in \mathbb{R}_+^2.$$

In a canonical way, one can think of B as “two-parameter Brownian motion”.

In this article, we address the following question: “Given a measurable function $v : \mathbb{R} \rightarrow \mathbb{R}_+$, what can be said about the distribution of $\int_{[0,1]^2} v(B_s) ds$?” The one-parameter variant of this question is both easy-to-state and well understood. Indeed, if b designates standard Brownian motion, the Laplace transform of $\int_0^1 v(b_s + x) ds$ often solves a Dirichlet eigenvalue problem (in x), as prescribed by the Feynman–Kac formula; cf. Revuz and Yor [6], for example. While analogues of Feynman–Kac for B are not yet known to hold, the following highlights some of the unusual behavior of $\int_{[0,1]^2} v(B_s) ds$ in case $v = \mathbf{1}_{[0,\infty)}$ and, anecdotally, implies that finding explicit formulæ may present a challenging task.

Theorem 1.1

There exists a $c_0 \in (0, 1)$, such that for all $0 < \varepsilon < \frac{1}{8}$,

$$\exp\left\{-\frac{1}{c_0} \log^2(1/\varepsilon)\right\} \leq \mathbb{P}\left\{\int_{[0,1]^2} \mathbf{1}_{\{B_s > 0\}} ds < \varepsilon\right\} \leq \exp\left\{-c_0 \log^2(1/\varepsilon)\right\}.$$

*Research supported in part by grants from NSF and NATO

†Research supported in part by NSF grant 98-03249

Remark 1.2

By the arcsine law, the one-parameter version of the above has the following simple form: given a linear Brownian motion b ,

$$\lim_{\varepsilon \rightarrow 0^+} \varepsilon^{-1/2} \mathbb{P} \left\{ \int_0^1 \mathbf{1}_{\{b_s > 0\}} ds < \varepsilon \right\} = \frac{2}{\pi};$$

see [6, Theorem 2.7, Ch. 6]. □

Remark 1.3

R. Pyke (personal communication) has asked whether $\int_{[0,1]^2} \mathbf{1}_{\{B_s > 0\}} ds$ has an arcsine-type law; see [5, Section 4.3.2] for a variant of this question in discrete time. According to Theorem 1.1, as $\varepsilon \rightarrow 0$, the cumulative distribution function of $\int_{[0,1]^2} \mathbf{1}_{\{B_s > 0\}} ds$ goes to zero faster than any power of ε . In particular, the distribution of time (in $[0, 1]^2$) spent positive does not have any simple extension of the arcsine law. □

Theorem 1.4

Let $v(x) := \mathbf{1}_{[-1,1]}(x)$, or $v(x) := \mathbf{1}_{(-\infty,1)}(x)$. Then, there exists a $c_1 \in (0, 1)$, such that for all $\varepsilon \in (0, \frac{1}{8})$,

$$\exp \left\{ - \frac{\log^3(1/\varepsilon)}{c_1 \varepsilon} \right\} \leq \mathbb{P} \left\{ \int_{[0,1]^2} v(B_s) ds < \varepsilon \right\} \leq \exp \left\{ - c_1 \frac{\log(1/\varepsilon)}{\varepsilon} \right\}.$$

For a refinement, see Theorem 2.2 below.

Remark 1.5

The one-parameter version of Theorem 1.4 is quite simple. For example, let $\Gamma = \int_0^1 \mathbf{1}_{[-1,1]}(b_s) ds$, where b is linear Brownian motion. In principle, one can compute the Laplace transform of Γ by means of Kac's formula and invert it to calculate its distribution function. However, direct arguments suffice to show that the two-parameter Theorem 1.4 is more subtle than its one-parameter counterpart:

$$-\infty < \liminf_{\varepsilon \rightarrow 0^+} \varepsilon \ln \mathbb{P} \{ \Gamma < \varepsilon \} \leq \limsup_{\varepsilon \rightarrow 0^+} \varepsilon \ln \mathbb{P} \{ \Gamma < \varepsilon \} < 0, \tag{1.1}$$

where \ln denotes the natural logarithm function. We will verify this later on in the Appendix. □

Remark 1.6

The arguments used to demonstrate Theorem 1.4 can be used to also estimate the distribution function of additive functionals of form, e.g., $\int_{[0,1]^2} v(B_s) ds$, as long as $\alpha \mathbf{1}_{[-r,r]} \leq v \leq \beta \mathbf{1}_{[-R,R]}$, where $0 < r \leq R$ and $0 < \alpha \leq \beta$. Other formulations are also possible. For instance, when $\alpha \mathbf{1}_{(-\infty,r]} \leq v \leq \beta \mathbf{1}_{(-\infty,R]}$. □

2 Proof of Theorems 1.1 and 1.4

Our proof of Theorem 1.1 rests on a lemma that is close in spirit to a Feynman-Kac formula of the theory of one-parameter Markov processes.

Proposition 2.1

There exists a finite and positive constant c_2 , such that for all measurable $D \subset \mathbb{R}$ and all $0 < \eta, \varepsilon < \frac{1}{8}$.

$$\mathbb{P}\left\{\int_{[0,1]^2} \mathbf{1}_{\{B_s \notin D\}} ds < \varepsilon\right\} \leq \mathbb{P}\{\forall s \in [0, 1]^2 : B_s \in D_{\varepsilon^{\frac{1}{4}-2\eta}}\} + \exp\{-c_2\varepsilon^{-\eta}\},$$

where D_δ denotes the δ -enlargement of D for any $\delta > 0$. That is,

$$D_\delta := \{x \in \mathbb{R} : \text{dist}(x; D) \leq \delta\},$$

where ‘dist’ denotes Hausdorff distance.

Proof For all $t \in [0, 1]^2$, let $|t| := \max\{t_1, t_2\}$. Then, it is clear that for any $\varepsilon, \delta > 0$, whenever there exists some $s_0 \in [0, 1]^2$ for which $B_{s_0} \notin D_\delta$, either

1. $\sup_{|t-s| \leq \varepsilon^{1/2}} |B_t - B_s| > \delta$, where the supremum is taken over all such choices of s and t in $[0, 1]^2$; or
2. for all $t \in [0, 1]^2$ with $|t-s_0| \leq \varepsilon^{1/2}$, $B_t \in D$, in which case, we can certainly deduce that $\int_{[0,1]^2} \mathbf{1}_{D^c}(B_t) dt > \varepsilon$.

Thus,

$$\begin{aligned} \mathbb{P}\{\exists s_0 \in [0, 1]^2 : B_{s_0} \notin D_\delta\} &\leq \mathbb{P}\{\sup_{|t-s| \leq \varepsilon^{1/2}} |B_t - B_s| > \delta\} + \\ &+ \mathbb{P}\{\int_{[0,1]^2} \mathbf{1}_{D^c}(B_t) dt > \varepsilon\}. \end{aligned}$$

By the general theory of Gaussian processes, there exists a universal positive and finite constant c_2 such that

$$\mathbb{P}\left\{\sup_{|t-s| \leq \varepsilon^{1/2}} |B_t - B_s| > \delta\right\} \leq \exp\{-c_2\delta^2\varepsilon^{-1/2}\}. \tag{2.1}$$

Although it is well known, we include a brief derivation of this inequality for completeness. Indeed, we recall C. Borell’s inequality from Adler [1, Theorem 2.1]: if $\{g_t; t \in T\}$ is a centered Gaussian process such that $\|g\|_T = \mathbb{E}\{\sup_{t \in T} |g_t|\} < \infty$ and whenever T is totally bounded in the metric $d(s, t) = \sqrt{\mathbb{E}\{(g_t - g_s)^2\}}$ ($s, t \in T$),

$$\mathbb{P}\{\sup_{t \in T} |g_t| \geq \lambda + \|g\|_T\} \leq 2 \exp\left\{-\frac{\lambda^2}{2\sigma_T^2}\right\},$$

where $\sigma_T^2 = \sup_{t \in T} \mathbb{E}\{g_t^2\}$. Eq. (2.1) follows from this by letting $T = \{(s, t) \in (0, 1)^2 \times (0, 1)^2 : |s - t| \leq \varepsilon^{1/2}\}$, $g_{t,s} = B_t - B_s$ and by making a few lines of standard calculations. Having derived (2.1), we can let $\delta := \varepsilon^{\frac{1}{4}-\frac{\eta}{2}}$ to obtain the proposition. \square

Proof of Theorem 1.1 Let $D = (-\infty, 0)$ and use Proposition 2.1 to see that

$$\mathbb{P}\left\{\int_{[0,1]^N} \mathbf{1}_{\{B_s > 0\}} < \varepsilon\right\} \leq \mathbb{P}\left\{\sup_{s \in [0,1]^2} B_s \leq \varepsilon^{\frac{1}{4} - 2\eta}\right\} + \exp\{-c_2 \varepsilon^{-\eta}\}.$$

Thus, the upper bound of Theorem 1.1 follows from Li and Shao [4], which states that

$$\limsup_{\varepsilon \rightarrow 0^+} \frac{1}{\log^2(1/\varepsilon)} \log \mathbb{P}\left\{\sup_{s \in [0,1]^2} B_s \leq \varepsilon\right\} < -\infty.$$

(An earlier, less refined version, of this estimate can be found in Csáki et al. [2].) To prove the lower bound, we note that

$$\begin{aligned} & \mathbb{P}\left\{\int_{[0,1]^2} \mathbf{1}_{\{B_s > 0\}} ds < 2\varepsilon - \varepsilon^2\right\} \\ & \geq \mathbb{P}\left\{\sup_{s \in [\varepsilon, 1]^2} B_s < 0\right\} \\ & = \mathbb{P}\left\{\forall(u, v) \in [0, \ln(\frac{1}{\varepsilon})]^2 : e^{(u+v)/2} B(e^{-u}, e^{-v}) < 0\right\}, \end{aligned}$$

and observe that the stochastic process $(u, v) \mapsto B(e^{-u}, e^{-v})/e^{-(u+v)/2}$ is the 2-parameter Ornstein–Uhlenbeck sheet. All that we need to know about the latter process is that it is a stationary, positively correlated Gaussian process whose law is supported on the space of continuous functions on $[0, 1]^2$. We define $c_3 > 0$ via the equation

$$e^{-c_3} := \mathbb{P}\left\{\forall(u, v) \in [0, 1]^2 : \frac{B(e^{-u}, e^{-v})}{e^{-(u+v)/2}} < 0\right\}.$$

By the support theorem, $0 < c_3 < \infty$; this is a consequence of the Cameron–Martin theorem on Gauss space; cf. Janson [3, Theorem 14.1]. Moreover, by stationarity and by Slepian’s inequality (cf. [1, Corollary 2.4]),

$$\begin{aligned} & \mathbb{P}\left\{\int_{[0,1]^2} \mathbf{1}_{\{B_s < 0\}} ds < \varepsilon\right\} \\ & \geq \prod_{0 \leq i, j \leq \ln(1/\varepsilon)+1} \mathbb{P}\left\{\forall(u, v) \in [i, i+1] \times [j, j+1] : \frac{B(e^{-u}, e^{-v})}{e^{-(u+v)/2}} < 0\right\} \\ & = \exp\left\{-c_3 \ln^2(e^2/\varepsilon)\right\}. \end{aligned}$$

This proves the theorem. □

Next, we prove Theorem 1.4.

Proof of Theorem 1.4 Let \mathcal{D}_ε denote the collection of all points $(s, t) \in [0, 1]^2$, such that $st \leq \varepsilon$. Note that

1. Lebesgue’s measure of \mathcal{D}_ε is at least $\varepsilon \ln(1/\varepsilon)$; and
2. if $\sup_{s \in \mathcal{D}_\varepsilon} |B_s| \leq 1$, then $\int_{[0,1]^2} \mathbf{1}_{(-1,1)}(B_s) ds > \varepsilon \ln(1/\varepsilon)$.

Thus,

$$\mathbb{P}\left\{\int_{[0,1]^2} \mathbf{1}_{(-1,1)}(B_s) ds < \varepsilon \ln(1/\varepsilon)\right\} \leq \mathbb{P}\left\{\sup_{s \in \mathcal{D}_\varepsilon} |B_s| > 1\right\}.$$

A basic feature of the set \mathcal{D}_ε is that whenever $s \in \mathcal{D}_\varepsilon$, then $\mathbb{E}\{B_s^2\} \leq \varepsilon$. Since $\mathbb{E}\{\sup_{s \in \mathcal{D}_\varepsilon} |B_s|\} \leq \mathbb{E}\{\sup_{s \in [0,1]^2} |B_s|\} < \infty$, we can apply Borell's inequality to deduce the existence of a finite, positive constant $c_4 < 1$, such that for all $\varepsilon > 0$, $\mathbb{P}\{\sup_{s \in \mathcal{D}_\varepsilon} |B_s| > 1/c_4\} \leq \exp\{-c_4/\varepsilon\}$. We apply Brownian scaling and possibly adjust c_4 to conclude that

$$\mathbb{P}\left\{\sup_{s \in \mathcal{D}_\varepsilon} |B_s| > 1\right\} \leq e^{-c_4/\varepsilon}.$$

Consequently, we can find a positive, finite constant c_5 , such that for all $\varepsilon \in (0, \frac{1}{8})$,

$$\mathbb{P}\{\Gamma < \varepsilon\} \leq \exp\left\{-c_5 \frac{\ln(1/\varepsilon)}{\varepsilon}\right\}. \tag{2.2}$$

This implies the upper bound in the conclusion of Theorem 1.4. For the lower bound, we note that for all $\varepsilon \in (0, \frac{1}{8})$, Lebesgue's measure of \mathcal{D}_ε is bounded above by $c_6\varepsilon \log(1/\varepsilon)$. Thus,

$$\mathbb{P}\left\{\int_{[0,1]^2} \mathbf{1}_{(-\infty,1)}(B_s) ds < c_6\varepsilon \log(1/\varepsilon)\right\} \geq \mathbb{P}\left\{\inf_{s \in [0,1]^2 \setminus \mathcal{D}_\varepsilon} B_s > 1\right\}.$$

On the other hand, whenever $s \in [0,1]^2 \setminus \mathcal{D}_\varepsilon$, $s_1s_2 \geq \varepsilon$. Thus,

$$\begin{aligned} \mathbb{P}\left\{\int_{[0,1]^2} \mathbf{1}_{(-\infty,1)}(B_s) ds < c_6\varepsilon \log(1/\varepsilon)\right\} &\geq \mathbb{P}\left\{\inf_{s \in [0,1]^2 \setminus \mathcal{D}_\varepsilon} \frac{B_s}{\sqrt{s_1s_2}} > \frac{1}{\sqrt{\varepsilon}}\right\} \\ &= \mathbb{P}\left\{\inf_{\substack{u,v \geq 0: \\ u+v \leq \ln(1/\varepsilon)}} O_{u,v} > \varepsilon^{-1/2}\right\}, \end{aligned}$$

where $O_{u,v} := B(e^{-u}, e^{-v})/e^{-(u+v)/2}$ is an Ornstein-Uhlenbeck sheet. Consequently,

$$\mathbb{P}\left\{\int_{[0,1]^2} \mathbf{1}_{(-\infty,1)}(B_s) ds < c_6\varepsilon \log(1/\varepsilon)\right\} \geq \mathbb{P}\left\{\inf_{0 \leq u,v \leq \ln(1/\varepsilon)} O_{u,v} > \varepsilon^{-1/2}\right\},$$

By appealing to Slepian's inequality and to the stationarity of O , we can deduce that

$$\begin{aligned} &\mathbb{P}\left\{\int_{[0,1]^2} \mathbf{1}_{(-\infty,1)}(B_s) ds < c_3\varepsilon \log(1/\varepsilon)\right\} \\ &\geq \prod_{0 \leq i,j \leq \ln(1/\varepsilon)} \mathbb{P}\left\{\inf_{i \leq u \leq i+1} \inf_{j \leq v \leq j+1} O_{u,v} > \varepsilon^{-1/2}\right\} \\ &= \left[\mathbb{P}\left\{\inf_{0 \leq u,v \leq 1} O_{u,v} > \varepsilon^{-1/2}\right\}\right]^{\ln^2(e/\varepsilon)}. \end{aligned} \tag{2.3}$$

On the other hand, recalling the construction of O , we have

$$\begin{aligned} & \mathbb{P}\left\{\inf_{0 \leq u, v \leq 1} O_{u, v} > \varepsilon^{-1/2}\right\} \\ & \geq \mathbb{P}\left\{\inf_{1 \leq s, t \leq e} B_{s, t} \geq e \varepsilon^{-1/2}\right\} \\ & \geq \mathbb{P}\left\{B_{1, 1} \geq 2e \varepsilon^{-1/2}, \sup_{1 \leq s_1, s_2 \leq e} |B_s - B_{1, 1}| \leq e \varepsilon^{-1/2}\right\} \\ & = \mathbb{P}\left\{B_{1, 1} \geq 2e \varepsilon^{-1/2}\right\} \cdot \mathbb{P}\left\{\sup_{1 \leq s_1, s_2 \leq e} |B_s - B_{1, 1}| \leq e \varepsilon^{-1/2}\right\} \\ & \geq c_7 \mathbb{P}\left\{B_{1, 1} \geq 2e \varepsilon^{-1/2}\right\}, \end{aligned}$$

for some absolute constant c_7 that is chosen independently of all $\varepsilon \in (0, \frac{1}{8})$. Therefore, by picking c_8 large enough, we can insure that for all $\varepsilon \in (0, \frac{1}{8})$,

$$\mathbb{P}\left\{\inf_{0 \leq u, v \leq 1} O_{u, v} > \varepsilon^{-1/2}\right\} \geq \exp\{-c_8 \varepsilon^{-1}\}.$$

Plugging this in to Eq. (2.3), we obtain

$$\mathbb{P}\left\{\int_{[0, 1]^2} \mathbf{1}_{(-\infty, 1)}(B_s) ds < c_6 \varepsilon \log(1/\varepsilon)\right\} \geq \exp\left\{-c_8 \frac{\ln^2(1/\varepsilon)}{4\varepsilon}\right\}. \quad (2.4)$$

The lower bound of Theorem 1.4 follows from replacing ε by $\varepsilon/\ln(1/\varepsilon)$. \square

The methods of this proof go through with few changes to derive the following extension of Theorem 1.4.

Theorem 2.2

Suppose $\varphi : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ is a measurable function such that (a) as $r \downarrow 0$, $\varphi(r) \uparrow \infty$; and (b) there exists a finite constant $\gamma > 0$, such that for all $r \in (0, \frac{1}{2})$, $\varphi(2r) \geq \gamma\varphi(r)$. Define $J_\varphi = \int_{[0, 1]^2} \mathbf{1}_{\{|B_s| \leq \sqrt{s_1 s_2} \varphi(s_1 s_2)\}} ds$. Then, there exist a finite constant $c_9 > 1$, such that for all $\varepsilon \in (0, \frac{1}{2})$,

$$\exp\left\{-c_9 \varphi^2\left(\frac{\varepsilon}{\log(1/\varepsilon)}\right) \log^2(1/\varepsilon)\right\} \leq \mathbb{P}\{J_\varphi < \varepsilon\} \leq \exp\left\{-\frac{1}{c_9} \varphi^2\left(\frac{\varepsilon}{\log(1/\varepsilon)}\right)\right\}.$$

Appendix: On Remark 1.5

In this appendix, we include a brief verification of the exponential form of the distribution function of Γ ; cf. Eq. (1.1). Given any $\lambda > \frac{1}{2}$ and for $\zeta = (2\lambda)^{-1/2}$, we have

$$\begin{aligned} \mathbb{E}\{e^{-\lambda\Gamma}\} & \leq \mathbb{E}\left\{\exp\left(-\lambda \int_0^\zeta v(b_s) ds\right)\right\} \\ & \leq e^{-\lambda\zeta} + \mathbb{P}\left\{\sup_{0 \leq s \leq \zeta} |b_s| > 1\right\} \\ & \leq e^{-\lambda\zeta} + e^{-1/(2\zeta)} \\ & = 2e^{-\sqrt{\lambda/2}}. \end{aligned} \quad (2.5)$$

$$(2.6)$$

By Chebyshev's inequality, $\mathbb{P}\{\int_0^1 v(b_s) ds < \varepsilon\} \leq 2 \inf_{\lambda > 1} e^{-\sqrt{\lambda/2} + \lambda\varepsilon}$. Choose $\lambda = \frac{1}{8}\varepsilon^{-2}$ to obtain the following for all $\varepsilon \in (0, \frac{1}{2})$:

$$\mathbb{P}\{\Gamma < \varepsilon\} \leq 2e^{-1/(8\varepsilon)}. \tag{2.7}$$

Conversely, we can choose $\delta = (2\lambda)^{-1/2}$ and $\eta \in (0, \frac{1}{100})$ to see that

$$\begin{aligned} \mathbb{E}\{e^{-\lambda\Gamma}\} &\geq \mathbb{E}\left\{\exp\left(-\lambda\int_0^\delta v(b_s) ds\right); \inf_{\delta \leq s \leq 1} |b_s| > 1\right\} \\ &\geq e^{-\lambda\delta} \mathbb{P}\{|b_\delta| > 1 + \eta, \sup_{\delta < s < 1+\delta} |b_s - b_\delta| < \eta\}. \end{aligned}$$

Thus, we can always find a positive, finite constant c_{10} that only depends on η and such that

$$\mathbb{E}\{e^{-\lambda\Gamma}\} \geq c_{10} \exp\left\{-\sqrt{\frac{\lambda}{2}} [1 + (1 + \eta)^2(1 + \psi_\delta)]\right\},$$

where $\lim_{\delta \rightarrow 0+} \psi_\delta = 0$, uniformly in $\eta \in (0, \frac{1}{100})$. In particular, after negotiating the constants, we obtain

$$\liminf_{\lambda \rightarrow \infty} \lambda^{-1/2} \ln \mathbb{E}\{e^{-\lambda\Gamma}\} \geq -2^{1/2}. \tag{2.8}$$

Thus, for any $\varepsilon \in (0, \frac{1}{100})$,

$$e^{-\sqrt{2\lambda}(1+o_1(1))} \leq \mathbb{E}\{e^{-\lambda\Gamma}\} \leq \mathbb{P}\{\Gamma < \varepsilon\} + e^{-\lambda\varepsilon},$$

where $o_1(1) \rightarrow 0$, as $\lambda \rightarrow \infty$, uniformly in $\varepsilon \in (0, \frac{1}{100})$. In particular, if we choose $\varepsilon = (1 + \eta)\sqrt{2/\lambda}$, where $\eta > 0$, we obtain

$$\mathbb{P}\{\Gamma < (1 + \eta)\sqrt{2/\lambda}\} \geq e^{-\sqrt{2\lambda}(1+o_2(1))},$$

where $o_2(1) \rightarrow 0$, as $\lambda \rightarrow \infty$. This, Eq. (2.7) and a few lines of calculations, together imply Eq. (1.1). \square

References

- [1] R. J. Adler (1990). *An Introduction to Continuity, Extrema, and Related Topics for General Gaussian Processes*, Institute of Mathematical Statistics, Lecture Notes–Monograph Series, Volume 12, Hayward, California.
- [2] E. Csáki, D. Khoshnevisan and Z. Shi (2000). Boundary crossings and the distribution function of the maximum of Brownian sheet. *Stochastic Processes and Their Applications* (To appear).
- [3] S. Janson (1997). *Gaussian Hilbert Spaces*, Cambridge Tracts in Mathematics, Cambridge University Press, Cambridge, UK

- [4] W. V. Li and Q.-M. Shao (2000) Lower tail probabilities of Gaussian processes. Preprint.
- [5] R. Pyke (1973). Partial sums of matrix arrays, and Brownian sheets. In *Stochastic Analysis*, 331–348, John Wiley and Sons, London, D. G. Kendall and E. F. Harding: Ed.'s.
- [6] D. Revuz and M. Yor (1991). *Continuous Martingales and Brownian Motion*, Second Edition, Springer-Verlag, Berlin.

DAVAR KHOSHNEVISAN
University of Utah
Department of Mathematics
155 S 1400 E JWB 233
Salt Lake City, UT 84112-0090
davar@math.utah.edu
<http://www.math.utah.edu/~davar>

ROBIN PEMANTLE
Ohio State University
Department of Mathematics
231 W. 18 Ave., Columbus, OH 43210
Columbus, OH 43210
pemantle@math.ohio-state.edu
<http://www.math.ohio-state.edu/~pemantle>