

## ANSWERS

1. Calculate the following limits.

(a)  $\lim_{n \rightarrow \infty} \left(\frac{n-1}{n}\right)^{2n} = e^{-2}$

(b)  $\lim_{x \rightarrow 0} (1 + 4x)^{1/x} = e^4$

2. Calculate the following.

(a)  $\frac{d}{dx} (\ln(\tanh x)) = (\sinh x \cosh x)^{-1}$

(b)  $\int \frac{z}{2z^2 + 8} dz = \frac{1}{4} \ln(2z^2 + 8) + C$

(c)  $\int \frac{\tan(\ln x)}{x} dx = \ln |\sec(\ln x)| + C,$

(d)  $\int \frac{dx}{x(1-x)} = \ln \left| \frac{x}{x-1} \right| + C,$

(e)  $\frac{dy}{dx}, \quad y = \frac{(x^2 + 3)^{2/3}(3x + 2)^2}{\sqrt{x+1}} \quad \frac{dy}{dx} = \left( \frac{4x}{3x^2 + 9} + \frac{6}{3x + 2} - \frac{1}{2x + 2} \right)$

(f)  $\int \frac{e^x}{1 + e^{2x}} = \arctan(e^x) + C$

3. Experiments show that the rate of change of the atmospheric pressure  $P(x)$  with altitude  $x$  is proportional to the pressure. Write down the resulting differential equation for  $P(x)$ , and solve it, assuming that the pressure at 6000 meters is half its value  $P_0$  at sea level.

$$\frac{dP}{dx} = -kP, \quad k > 0; \quad P(x) = P_0 e^{-kx}, \quad k = (\ln 2)/6000$$

4. Section 6.5 #18

5,565 years ago

5. Section 6.3

40.  $\ln |e^x - 1| + C$

46.  $V = 2\pi \int_0^1 x e^{-x^2} dx = \pi \left(1 - \frac{1}{e}\right)$

6. Section 6.6

2.  $y(x) = \frac{\frac{1}{3}x^3 - x}{x-1} + \frac{C}{x-1}$

4.  $y(x) = \sin x + C \cos x$

7. Stewart wants to become a millionaire after 10 years by buying \$5,000 worth of a company's stock, which he wants to choose carefully. What must the sustained, annualized growth rate of the stock be in order to achieve his goal?

$$1,000,000 = 10,000(1+r)^{20} \text{ or } r = 0.2588\dots, \text{ or about } 26\%$$

8. Newton's law of cooling states that the rate at which an object cools is proportional to the difference between the temperature  $\theta(t)$  of the object and the constant ambient temperature  $T$ ,

$$\frac{d\theta}{dt} = -k(\theta - T),$$

where  $k > 0$  is a constant depending on the object. A corpse is discovered at 2 pm, and its temperature is found to be 85°F, with the ambient air temperature being 68°F. Assuming  $k = 0.5 \text{ hr}^{-1}$ , find the time of death.

$\theta(t) = T + (\theta_0 - T)e^{-kt}$ . The corpse is discovered at  $t = 0$  (or 2 pm), and has a temperature  $\theta_0 = \theta(0) = 85^\circ\text{F}$ . Assume that  $t_d$  is the time of death and that  $\theta(t_d) = \theta_d = 98.6^\circ\text{F}$ . Thus

$$t_d = -\frac{1}{k} \ln \left( \frac{\theta_d - T}{\theta_0 - T} \right) = -1.176$$

or about 12:49 pm.

9. Section 6.8

45.  $x^2 \left[ \frac{xe^x}{1+e^{2x}} + 3 \arctan(e^x) \right]$

51.  $\frac{3(1+\arcsin(x))^2}{\sqrt{1-x^2}}$

67.  $\frac{1}{3} \arcsin\left(\frac{\sqrt{3}}{2}x\right) + C$

10. Section 6.9

25.  $2 \tanh(x) \cosh(2x) + \sinh(2x) \operatorname{sech}^2(x)$

33.  $\frac{1}{\sqrt{x^2-1} \cosh(x)}$

41.  $2 \cosh(2z^{\frac{1}{4}}) + C$

45.  $\frac{1}{4} [\ln(\sinh(x^2))]^2 + C$