

Twenty-nine

- ① Line integrals
- ② Idea of line integrals
- ③ Four useful lemmas
- ④ Independence of path
- ⑤ Independence of parametrization

① Line integrals

Computing line integrals

C parametrized by $x=x(t)$, $y=y(t)$, $a \leq t \leq b$

$$f: \mathbb{R}^2 \rightarrow \mathbb{R}$$

Computing line integrals

C parametrized by $x=x(t)$, $y=y(t)$, $a \leq t \leq b$

$$f: \mathbb{R}^2 \rightarrow \mathbb{R}$$

- $\int_C f(x,y) d\Delta = \int_a^b f(x(t), y(t)) \|(x'(t), y'(t))\| dt$

- $\int_C f(x,y) dx = \int_a^b f(x(t), y(t)) x'(t) dt$

- $\int_C f(x,y) dy = \int_a^b f(x(t), y(t)) y'(t) dt$

Computing line integrals

C parametrized by $x=x(t)$, $y=y(t)$, $a \leq t \leq b$

$f: \mathbb{R}^2 \rightarrow \mathbb{R}$


- $\int_C f(x,y) d\Delta = \int_a^b f(x(t), y(t)) \|(x'(t), y'(t))\| dt$

- $\int_C f(x,y) dx = \int_a^b f(x(t), y(t)) x'(t) dt$

- $\int_C f(x,y) dy = \int_a^b f(x(t), y(t)) y'(t) dt$

Example: C given by $x(t)=t$, $y(t)=t^2$, $1 \leq t \leq 3$.

Find $\int_C \frac{1}{\sqrt{4y+1}} d\Delta$.




Example: C given by $x(t)=t$, $y(t)=t^2$, $1 \leq t \leq 3$.

Find $\int_C \frac{1}{\sqrt{4y+1}} d\Delta$.

Solution:

$$\int_C \frac{1}{\sqrt{4y+1}} d\Delta = \int_1^3 \frac{1}{\sqrt{4t^2+1}} \|(x'(t), y'(t))\| dt$$




Example: C given by $x(t)=t$, $y(t)=t^2$, $1 \leq t \leq 3$.

Find $\int_C \frac{1}{\sqrt{4y+1}} d\Delta$.


Solution:

$$\begin{aligned} \int_C \frac{1}{\sqrt{4y+1}} d\Delta &= \int_1^3 \frac{1}{\sqrt{4t^2+1}} \|(x'(t), y'(t))\| dt \\ &= \int_1^3 \frac{1}{\sqrt{4t^2+1}} \|(1, 2t)\| dt \end{aligned}$$



Example: C given by $x(t)=t$, $y(t)=t^2$, $1 \leq t \leq 3$.

Find $\int_C \frac{1}{\sqrt{4y+1}} d\Delta$.



Solution:


$$\int_C \frac{1}{\sqrt{4y+1}} d\Delta = \int_1^3 \frac{1}{\sqrt{4t^2+1}} \|(x'(t), y'(t))\| dt$$

$$= \int_1^3 \frac{1}{\sqrt{4t^2+1}} \|(1, 2t)\| dt$$

$$= \int_1^3 \frac{1}{\sqrt{4t^2+1}} \sqrt{1^2 + (2t)^2} dt$$

Example: C given by $x(t)=t$, $y(t)=t^2$, $1 \leq t \leq 3$.

Find $\int_C \frac{1}{\sqrt{4y+1}} d\Delta$.



Solution:

$$\int_C \frac{1}{\sqrt{4y+1}} d\Delta = \int_1^3 \frac{1}{\sqrt{4t^2+1}} \|(x'(t), y'(t))\| dt$$


$$= \int_1^3 \frac{1}{\sqrt{4t^2+1}} \|(1, 2t)\| dt$$

$$= \int_1^3 \frac{1}{\sqrt{4t^2+1}} \sqrt{1^2 + (2t)^2} dt$$

$$= \int_1^3 dt$$

Example: C given by $x(t)=t$, $y(t)=t^2$, $1 \leq t \leq 3$.

Find $\int_C \frac{1}{\sqrt{4y+1}} d\Delta$.



Solution:

$$\int_C \frac{1}{\sqrt{4y+1}} d\Delta = \int_1^3 \frac{1}{\sqrt{4t^2+1}} \|(x'(t), y'(t))\| dt$$

$$= \int_1^3 \frac{1}{\sqrt{4t^2+1}} \|(1, 2t)\| dt$$

$$= \int_1^3 \frac{1}{\sqrt{4t^2+1}} \sqrt{1^2 + (2t)^2} dt$$

$$= \int_1^3 dt = 2$$

Definition:

$$\int_C (f(x,y)dx + g(x,y)dy)$$

$$= \int_C f(x,y)dx + \int_C g(x,y)dy$$

Definition:

$$\int_C (f(x,y) dx + g(x,y) dy)$$

$$= \int_C f(x,y) dx + \int_C g(x,y) dy$$

Consequently:


$$\int_C f(x,y) dx + g(x,y) dy$$

$$= \int_a^b (f(x(t), y(t)) x'(t) + g(x(t), y(t)) y'(t)) dt$$

Example: C_1 given by $x=t, y=t, z=t, 0 \leq t \leq 1$.

C_2 given by $x=t^3, y=t^2, z=t, 0 \leq t \leq 1$.

Find $\int_{C_i} yz dx + xz dy + xy dz$ for $i=1,2$.



Example: C_1 given by $x=t$, $y=t$, $z=t$, $0 \leq t \leq 1$.


C_2 given by $x=t^3$, $y=t^2$, $z=t$, $0 \leq t \leq 1$.

Find $\int_{C_i} yz dx + xz dy + xy dz$ for $i=1,2$.

Solution:

$$\int_{C_1} yz dx + xz dy + xy dz$$

$$= \int_0^1 t \cdot t \cdot 1 + t \cdot t \cdot 1 + t \cdot t \cdot 1 dt$$



Example: C_1 given by $x=t$, $y=t$, $z=t$, $0 \leq t \leq 1$.

C_2 given by $x=t^3$, $y=t^2$, $z=t$, $0 \leq t \leq 1$.


Find $\int_{C_i} yz dx + xz dy + xy dz$ for $i=1,2$.

Solution:

$$\int_{C_1} yz dx + xz dy + xy dz$$

$$= \int_0^1 t \cdot t \cdot 1 + t \cdot t \cdot 1 + t \cdot t \cdot 1 dt$$

$$= \int_0^1 3t^2 dt$$



Example: C_1 given by $x=t$, $y=t$, $z=t$, $0 \leq t \leq 1$.

C_2 given by $x=t^3$, $y=t^2$, $z=t$, $0 \leq t \leq 1$.

Find $\int_{C_i} yz dx + xz dy + xy dz$ for $i=1,2$.


Solution:

$$\int_{C_1} yz dx + xz dy + xy dz$$

$$= \int_0^1 t \cdot t \cdot 1 + t \cdot t \cdot 1 + t \cdot t \cdot 1 dt$$

$$= \int_0^1 3t^2 dt$$

$$= [t^3]_0^1$$



Example: C_1 given by $x=t$, $y=t$, $z=t$, $0 \leq t \leq 1$.

C_2 given by $x=t^3$, $y=t^2$, $z=t$, $0 \leq t \leq 1$.

Find $\int_{C_i} yz dx + xz dy + xy dz$ for $i=1,2$.


Solution:

$$\int_{C_1} yz dx + xz dy + xy dz$$

$$= \int_0^1 t \cdot t \cdot 1 + t \cdot t \cdot 1 + t \cdot t \cdot 1 dt$$

$$= \int_0^1 3t^2 dt$$

$$= [t^3]_0^1 = 1^3 - 0^3 = 1.$$




Example: C_1 given by $x=t, y=t, z=t, 0 \leq t \leq 1$.

C_2 given by $x=t^3, y=t^2, z=t, 0 \leq t \leq 1$.

Find $\int_{C_i} yz dx + xz dy + xy dz$ for $i=1,2$.

Solution:

$$\begin{aligned} & \int_{C_2} yz dx + xz dy + xy dz \\ &= \int_0^1 t^2 \cdot t \cdot 3t^2 + t^3 \cdot t \cdot 2t + t^3 \cdot t^2 \cdot 1 dt \end{aligned}$$



Example: C_1 given by $x=t, y=t, z=t, 0 \leq t \leq 1$.

C_2 given by $x=t^3, y=t^2, z=t, 0 \leq t \leq 1$.


Find $\int_{C_i} yz dx + xz dy + xy dz$ for $i=1,2$.

Solution:

$$\int_{C_2} yz dx + xz dy + xy dz$$

$$= \int_0^1 t^2 \cdot t \cdot 3t^2 + t^3 \cdot t \cdot 2t + t^3 \cdot t^2 \cdot 1 dt$$

$$= \int_0^1 6t^5 dt$$




Example: C_1 given by $x=t, y=t, z=t, 0 \leq t \leq 1$.

C_2 given by $x=t^3, y=t^2, z=t, 0 \leq t \leq 1$.

Find $\int_{C_i} yz dx + xz dy + xy dz$ for $i=1,2$.

Solution:

$$\begin{aligned} & \int_{C_2} yz dx + xz dy + xy dz \\ &= \int_0^1 t^2 \cdot t \cdot 3t^2 + t^3 \cdot t \cdot 2t + t^3 \cdot t^2 \cdot 1 dt \\ &= \int_0^1 6t^5 dt \\ &= [t^6]_0^1 \end{aligned}$$




Example: C_1 given by $x=t, y=t, z=t, 0 \leq t \leq 1$.

C_2 given by $x=t^3, y=t^2, z=t, 0 \leq t \leq 1$.


Find $\int_{C_i} yz dx + xz dy + xy dz$ for $i=1,2$.

Solution:

$$\begin{aligned} & \int_{C_2} yz dx + xz dy + xy dz \\ &= \int_0^1 t^2 \cdot t \cdot 3t^2 + t^3 \cdot t \cdot 2t + t^3 \cdot t^2 \cdot 1 dt \\ &= \int_0^1 6t^5 dt \\ &= [t^6]_0^1 = 1^6 - 0^6 = 1. \end{aligned}$$



Integration along piecewise paths



$$C = C_1 \cup C_2 \cup C_3$$

$$\int_C f(x,y) d* = \int_{C_1} f(x,y) d* + \int_{C_2} f(x,y) d* + \int_{C_3} f(x,y) d*$$


(whatever $*$ is)

Example: C_1 given by $x=t, y=t, z=t, 0 \leq t \leq 1$.

C_2 given by $x=t^3, y=t^2, z=t, 0 \leq t \leq 1$.

Find $\int_C yz dx + xz dy + xy dz$ for
the simple closed curve

$$C = C_1 \cup -C_2.$$



Example: C_1 given by $x=t, y=t, z=t, 0 \leq t \leq 1$.


C_2 given by $x=t^3, y=t^2, z=t, 0 \leq t \leq 1$.

Find $\int_C yz dx + xz dy + xy dz$ for
the simple closed curve

$$C = C_1 \cup -C_2.$$


Solution:

$$\begin{aligned} \int_C yz dx + xz dy + xy dz \\ = \int_{C_1} yz dx + xz dy + xy dz + \int_{-C_2} yz dx + xz dy + xy dz \end{aligned}$$




② Idea of
line integrals

Idea of $\int_C ds$



C : $a \leq t \leq b$
 $x = x(t)$
 $y = y(t)$


Idea of $\int_C ds$



$$\underline{C}: \begin{aligned} a \leq t \leq b \\ x = x(t) \\ y = y(t) \end{aligned}$$

- $\int_C ds =$


Idea of $\int_C ds$



$$\underline{C}: \begin{aligned} a \leq t \leq b \\ x = x(t) \\ y = y(t) \end{aligned}$$

- $\int_C ds = \int_a^b \|(x'(t), y'(t))\| dt$


Idea of $\int_c ds$



$$\underline{C}: \begin{aligned} a \leq t \leq b \\ x = x(t) \\ y = y(t) \end{aligned}$$

- $\int_c ds = \int_a^b \|(x'(t), y'(t))\| dt = \text{Arclength}(C)$


Idea of $\int_c ds$



$$\underline{C}: \begin{aligned} a \leq t \leq b \\ x = x(t) \\ y = y(t) \end{aligned}$$


- $\int_c ds = \int_a^b \|(x'(t), y'(t))\| dt = \text{Arclength}(C)$
- $\int_c ds = \int_{-c} ds$

Idea of $\int_C dx$



$$\underline{C}: \begin{aligned} a \leq t \leq b \\ x = x(t) \\ y = y(t) \end{aligned}$$


Idea of $\int_C dx$



$$\underline{C}: \begin{aligned} a \leq t \leq b \\ x = x(t) \\ y = y(t) \end{aligned}$$

- $\int_C dx =$


Idea of $\int_C dx$



$$\underline{C}: \begin{aligned} a &\leq t \leq b \\ x &= x(t) \\ y &= y(t) \end{aligned}$$

- $\int_C dx = \int_a^b x'(t) dt$


Idea of $\int_C dx$



$$\underline{C}: \begin{aligned} a \leq t \leq b \\ x = x(t) \\ y = y(t) \end{aligned}$$

- $\int_C dx = \int_a^b x'(t) dt = [x(t)]_a^b$


Idea of $\int_C dx$



$$\underline{C}: \begin{aligned} a \leq t \leq b \\ x = x(t) \\ y = y(t) \end{aligned}$$

- $\int_C dx = \int_a^b x'(t) dt = [x(t)]_a^b = x(b) - x(a)$


Idea of $\int_C dx$



$$\begin{aligned} \underline{C}: \quad & a \leq t \leq b \\ & x = x(t) \\ & y = y(t) \end{aligned}$$

- $$\int_C dx = \int_a^b x'(t) dt = [x(t)]_a^b = \underbrace{x(b) - x(a)}_{\text{change in } x\text{-coordinate}}$$


Idea of $\int_C dx$



$$\underline{C}: \begin{aligned} a \leq t \leq b \\ x = x(t) \\ y = y(t) \end{aligned}$$

- $\int_C dx = \int_a^b x'(t) dt = [x(t)]_a^b = \underbrace{x(b) - x(a)}_{\text{change in } x\text{-coordinate}}$
- $\int_{-C} dx = - \int_C dx$

Idea of $\int_C dy$



$$\underline{C}: \begin{aligned} a \leq t \leq b \\ x = x(t) \\ y = y(t) \end{aligned}$$

- $\int_C dy = \int_a^b y'(t) dt = [y(t)]_a^b = \underbrace{y(b) - y(a)}_{\text{change in } y\text{-coordinate}}$
- $\int_{-C} dy = -\int_C dy$


Example:

C parametrized by

$$x = 2 + 4t$$

$$y = 7 - 6t$$

$$0 \leq t \leq 1.$$



Find $\int_C ds =$

$$\int_C dx =$$

$$\int_C dy =$$


Example:

C parametrized by

$$x = 2 + 4t$$

$$y = 7 - 6t$$

$$0 \leq t \leq 1.$$




$$\int_C ds = \sqrt{4^2 + 6^2} = 2\sqrt{13}$$

$$\int_C dx = 6 - 2 = 4$$

$$\int_C dy = 1 - 7 = -6$$

Example:

C parametrized by
 $x = \cos t,$
 $y = \sin t,$
 $0 \leq t \leq \pi.$




Find $\int_C ds =$

$$\int_C dx =$$

$$\int_C dy =$$

Example:

C parametrized by
 $x = \cos t,$
 $y = \sin t,$
 $0 \leq t \leq \pi.$




$$\int_C ds = \pi$$


$$\int_C dx = -2$$

$$\int_C dy = 0$$


$\int_c f(x,y) ds$ is arclength weighted by $f(x,y)$




$\int_C f(x,y) ds$ is arclength weighted by $f(x,y)$



$\int_C f(x,y) ds$ is arclength weighted by $f(x,y)$



$\int_C f(x,y) ds$ is arclength weighted by $f(x,y)$



• $\int_C f(x,y) ds = \int_{-C} f(x,y) ds$

Conversely,

$$\int_{-c} f(x,y)dx + g(x,y)dy = - \int_c f(x,y)dx + g(x,y)dy$$

Example: C_1 given by $x=t, y=t, z=t, 0 \leq t \leq 1$.


C_2 given by $x=t^3, y=t^2, z=t, 0 \leq t \leq 1$.

Find $\int_C yz dx + xz dy + xy dz$ for
the simple closed curve

$$C = C_1 \cup -C_2.$$

Solution:

$$\begin{aligned} \int_C yz dx + xz dy + xy dz \\ = \int_{C_1} yz dx + xz dy + xy dz + \int_{-C_2} yz dx + xz dy + xy dz \end{aligned}$$



Example: C_1 given by $x=t, y=t, z=t, 0 \leq t \leq 1$.

C_2 given by $x=t^3, y=t^2, z=t, 0 \leq t \leq 1$.

Find $\int_C yzdx + xzdy + xydz$ for
the simple closed curve


$$C = C_1 \cup -C_2.$$

Solution:

$$\int_C yzdx + xzdy + xydz$$

$$= \int_{C_1} yzdx + xzdy + xydz + \int_{-C_2} yzdx + xzdy + xydz$$

$$= \int_{C_1} yzdx + xzdy + xydz - \int_{C_2} yzdx + xzdy + xydz$$



Example: C_1 given by $x=t, y=t, z=t, 0 \leq t \leq 1$.

C_2 given by $x=t^3, y=t^2, z=t, 0 \leq t \leq 1$.

Find $\int_C yzdx + xzdy + xydz$ for
the simple closed curve

$$C = C_1 \cup -C_2.$$


Solution:

$$\int_C yzdx + xzdy + xydz$$

$$= \int_{C_1} yzdx + xzdy + xydz + \int_{-C_2} yzdx + xzdy + xydz$$

$$= \int_{C_1} yzdx + xzdy + xydz - \int_{C_2} yzdx + xzdy + xydz$$


$$= 1 - 1 = 0.$$



③ Four useful lemmas

Lemma:


If C is a vertical straight line in $\mathbb{R}^2$, then for any $f: \mathbb{R}^2 \rightarrow \mathbb{R}$, $\int_C f(x, y) dx = 0$.



Lemma:

If C is a vertical straight line in $\mathbb{R}^2$, then for any $f: \mathbb{R}^2 \rightarrow \mathbb{R}$, $\int_C f(x, y) dx = 0$.

proof:




Lemma:

If C is a vertical straight line in $\mathbb{R}^2$, then for any $f: \mathbb{R}^2 \rightarrow \mathbb{R}$, $\int_C f(x, y) dx = 0$.

proof:

$$C: x = x_0, y = y(t), a \leq t \leq b.$$




Lemma:

If C is a vertical straight line in $\mathbb{R}^2$, then for any $f: \mathbb{R}^2 \rightarrow \mathbb{R}$, $\int_C f(x, y) dx = 0$.

proof:


$$C: x = x_0, y = y(t), a \leq t \leq b.$$

$$\int_C f(x, y) dx =$$



Lemma:

If C is a vertical straight line in $\mathbb{R}^2$, then for any $f: \mathbb{R}^2 \rightarrow \mathbb{R}$, $\int_C f(x, y) dx = 0$.




proof:

$$C: x = x_0, y = y(t), a \leq t \leq b.$$

$$\int_C f(x, y) dx = \int_a^b f(x(t), y(t)) x'(t) dt$$

Lemma:

If C is a vertical straight line in $\mathbb{R}^2$, then for any $f: \mathbb{R}^2 \rightarrow \mathbb{R}$, $\int_C f(x, y) dx = 0$.




proof:

$$C: x = x_0, y = y(t), a \leq t \leq b.$$

$$\begin{aligned} \int_C f(x, y) dx &= \int_a^b f(x(t), y(t)) x'(t) dt \\ &= \int_a^b f(x(t), y(t)) 0 dt \end{aligned}$$

Lemma:

If C is a vertical straight line in $\mathbb{R}^2$, then for any $f: \mathbb{R}^2 \rightarrow \mathbb{R}$, $\int_C f(x, y) dx = 0$.




proof:

$$C: x = x_0, y = y(t), a \leq t \leq b.$$

$$\begin{aligned} \int_C f(x, y) dx &= \int_a^b f(x(t), y(t)) x'(t) dt \\ &= \int_a^b f(x(t), y(t)) 0 dt \\ &= 0. \end{aligned}$$

Lemma:

If C is a vertical straight line in $\mathbb{R}^2$, then for any $f: \mathbb{R}^2 \rightarrow \mathbb{R}$, $\int_C f(x, y) dx = 0$.



proof:

$$C: x = x_0, y = y(t), a \leq t \leq b.$$

$$\int_C f(x, y) dx = \int_a^b f(x(t), y(t)) x'(t) dt$$


$$= \int_a^b f(x(t), y(t)) 0 dt$$

$$= 0.$$

END

Lemma:


If C is a horizontal straight line in $\mathbb{R}^2$, then for any $f: \mathbb{R}^2 \rightarrow \mathbb{R}$, $\int_C f(x, y) dy = 0$.



Lemma:

If C is a curve
with a horizontal end,
and terminal point (x, y) ,
then

$$\frac{\partial}{\partial x} \left(\int_C f(x, y) dx + g(x, y) dy \right) = f(x, y).$$




Lemma:

If C is a curve with a horizontal end, and terminal point (x, y) , then

$$\frac{\partial}{\partial x} \left(\int_C f(x, y) dx + g(x, y) dy \right) = f(x, y).$$

proof:




Lemma:

If C is a curve with a horizontal end, and terminal point (x, y) , then

$$\frac{\partial}{\partial x} \left(\int_C f(x, y) dx + g(x, y) dy \right) = f(x, y).$$

proof:

$$\frac{\partial}{\partial x} \left(\int_C f(x, y) dx + g(x, y) dy \right) =$$




Lemma:

If C is a curve with a horizontal end, and terminal point (x, y) , then

$$\frac{\partial}{\partial x} \left(\int_C f(x, y) dx + g(x, y) dy \right) = f(x, y).$$

proof:

$$\frac{\partial}{\partial x} \left(\int_C f(x, y) dx + g(x, y) dy \right) =$$



Lemma:


If C is a curve with a horizontal end, and terminal point (x, y) , then

$$\frac{\partial}{\partial x} \left(\int_C f(x, y) dx + g(x, y) dy \right) = f(x, y).$$

proof:

$$\frac{\partial}{\partial x} \left(\int_C f(x, y) dx + g(x, y) dy \right) =$$

$$\frac{\partial}{\partial x} \left(\int_{C_1} f(x, y) dx + g(x, y) dy \right) + \frac{\partial}{\partial x} \left(\int_{C_2} f(x, y) dx + g(x, y) dy \right)$$




proof:

$$\frac{\partial}{\partial x} \left(\int_C f(x,y) dx + g(x,y) dy \right) =$$


$$= \frac{\partial}{\partial x} \left(\int_{C_1} f(x,y) dx + g(x,y) dy \right) + \frac{\partial}{\partial x} \left(\int_{C_2} f(x,y) dx + g(x,y) dy \right)$$

$$C = C_1 \cup C_2$$



proof:

$$\frac{\partial}{\partial x} \left(\int_C f(x,y) dx + g(x,y) dy \right) =$$




$$= \frac{\partial}{\partial x} \left(\int_{C_1} f(x,y) dx + g(x,y) dy \right) + \frac{\partial}{\partial x} \left(\int_{C_2} f(x,y) dx + g(x,y) dy \right)$$

$$= 0 +$$

proof:

$$\frac{\partial}{\partial x} \left(\int_C f(x,y) dx + g(x,y) dy \right) =$$




$$= \frac{\partial}{\partial x} \left(\int_{C_1} f(x,y) dx + g(x,y) dy \right) + \frac{\partial}{\partial x} \left(\int_{C_2} f(x,y) dx + g(x,y) dy \right)$$

$$= 0 + \frac{\partial}{\partial x} \left(\int_{C_2} f(x,y) dx \right) + \frac{\partial}{\partial x} \left(\int_{C_2} g(x,y) dy \right)$$

proof:

$$\frac{\partial}{\partial x} \left(\int_C f(x,y) dx + g(x,y) dy \right) =$$




$$= \frac{\partial}{\partial x} \left(\int_{C_1} f(x,y) dx + g(x,y) dy \right) + \frac{\partial}{\partial x} \left(\int_{C_2} f(x,y) dx + g(x,y) dy \right)$$

$$= 0 + \frac{\partial}{\partial x} \left(\int_{C_2} f(x,y) dx \right) + \frac{\partial}{\partial x} \left(\int_{C_2} g(x,y) dy \right)$$

$$= \quad \quad \quad + 0$$

proof:

$$\frac{\partial}{\partial x} \left(\int_C f(x,y) dx + g(x,y) dy \right) =$$



$$= \frac{\partial}{\partial x} \left(\int_{C_1} f(x,y) dx + g(x,y) dy \right) + \frac{\partial}{\partial x} \left(\int_{C_2} f(x,y) dx + g(x,y) dy \right)$$


$$= 0 + \frac{\partial}{\partial x} \left(\int_{C_2} f(x,y) dx \right) + \frac{\partial}{\partial x} \left(\int_{C_2} g(x,y) dy \right)$$

$$= \quad \quad \quad + 0$$

$$\begin{aligned} \underline{C_2}: \quad & x_0 \leq t \leq x \\ & x = t \\ & y = y \end{aligned}$$

proof:

$$\frac{\partial}{\partial x} \left(\int_C f(x,y) dx + g(x,y) dy \right) =$$



$$= \frac{\partial}{\partial x} \left(\int_{C_1} f(x,y) dx + g(x,y) dy \right) + \frac{\partial}{\partial x} \left(\int_{C_2} f(x,y) dx + g(x,y) dy \right)$$


$$= 0 + \frac{\partial}{\partial x} \left(\int_{C_2} f(x,y) dx \right) + \frac{\partial}{\partial x} \left(\int_{C_2} g(x,y) dy \right)$$

$$= \frac{\partial}{\partial x} \left(\int_{x_0}^x f(t,y) dt \right) + 0$$

C₂: $x_0 \leq t \leq x$
 $x = t$
 $y = y$

proof:

$$\frac{\partial}{\partial x} \left(\int_C f(x,y) dx + g(x,y) dy \right) =$$



$$= \frac{\partial}{\partial x} \left(\int_{C_1} f(x,y) dx + g(x,y) dy \right) + \frac{\partial}{\partial x} \left(\int_{C_2} f(x,y) dx + g(x,y) dy \right)$$

$$= 0 + \frac{\partial}{\partial x} \left(\int_{C_2} f(x,y) dx \right) + \frac{\partial}{\partial x} \left(\int_{C_2} g(x,y) dy \right)$$

$$= \frac{\partial}{\partial x} \left(\int_{x_0}^x f(t,y) dt \right) + 0$$

$$= f(x,y).$$


$$\underline{C_2}: \quad x_0 \leq t \leq x$$

$$x = t$$

$$y = y$$

proof:

$$\frac{\partial}{\partial x} \left(\int_C f(x,y) dx + g(x,y) dy \right) =$$



$$= \frac{\partial}{\partial x} \left(\int_{C_1} f(x,y) dx + g(x,y) dy \right) + \frac{\partial}{\partial x} \left(\int_{C_2} f(x,y) dx + g(x,y) dy \right)$$

$$= 0 + \frac{\partial}{\partial x} \left(\int_{C_2} f(x,y) dx \right) + \frac{\partial}{\partial x} \left(\int_{C_2} g(x,y) dy \right)$$

$$= \frac{\partial}{\partial x} \left(\int_{x_0}^x f(t,y) dt \right) + 0$$

$$= f(x,y).$$

C_2 : $x_0 \leq t \leq x$

$$x = t$$


$$y = y$$

END

Lemma:


If C is a curve
with a vertical end,
and terminal point (x, y) ,
then


$$\frac{\partial}{\partial y} \left(\int_C f(x, y) dx + g(x, y) dy \right) = g(x, y).$$




④

Independence
of path







$$\int_{C_1} dx = b_1 - a_1$$



$$\int_{C_2} dx = b_1 - a_1$$



$$\int_{C_3} dx = b_1 - a_1$$



$$\int_{C_1} dx = b_1 - a_1 \quad \int_{C_2} dx = b_1 - a_1 \quad \int_{C_3} dx = b_1 - a_1$$


If C is any path with initial point $\vec{a}$ and terminal point $\vec{b}$, then $\int_C dx = b_1 - a_1$.

If

Independence of path

$$\int_{C_1} f(x,y)dx + g(x,y)dy$$

$$= \int_{C_2} f(x,y)dx + g(x,y)dy$$



whenever C_1 and C_2 have the same initial points and the same terminal points, we say


$\int_C f(x,y)dx + g(x,y)dy$ is independent of path.

If

Independence of path

$$\int_{C_1} f(x,y)dx + g(x,y)dy$$

$$= \int_{C_2} f(x,y)dx + g(x,y)dy$$




whenever C_1 and C_2 have the same initial points and the same terminal points, we say

$\int_C f(x,y)dx + g(x,y)dy$ is independent of path.

Example: $\int_C dx$ is independent of path.

Independence of path


If



- C is a curve with initial point $\vec{a}$ and terminal point $\vec{b}$, and

Independence of path


If



- C is a curve with initial point $\vec{a}$ and terminal point $\vec{b}$, and
- $\int_C f(x,y)dx + g(x,y)dy$ is independent of path,

Independence of path

If



- C is a curve with initial point $\vec{a}$ and terminal point $\vec{b}$, and
- $\int_C f(x,y)dx + g(x,y)dy$ is independent of path,

then we write

$$\int_C f(x,y)dx + g(x,y)dy = \int_{\vec{a}}^{\vec{b}} f(x,y)dx + g(x,y)dy$$

Example: We'll see later that

$\int_C yz dx + xz dy + xy dz$ is independent of path.

Example: We'll see later that

$\int_C yz dx + xz dy + xy dz$ is independent of path.

So


$$\int_{C_i} yz dx + xz dy + xy dz$$

=

$$\int_{(0,0,0)}^{(1,1,1)} yz dx + xz dy + xy dz$$


=

1




⑤ Independence of
parametrization

Any curve has two
choices for **orientation** :
forwards or backwards.



Any curve has two choices for **orientation**: forwards or backwards.



As long as orientation is unchanged, line integrals are independent of parametrization.
(Using change of variables.)