

Thirty

- ① Line integrals and vector fields
- ② Line integrals and gradients
- ③ Theorem about line integrals, vector fields, and gradients

① Line integrals
and vector fields

C , a curve in $\mathbb{R}^3$.

$$\int_C x^2 y \, dx + (2x - z) \, dy + xy^3 z \, dz$$

C , a curve in $\mathbb{R}^3$.


$$\int_C x^2 y \, dx + (2x - z) \, dy + xy^3 z \, dz$$

C , a curve in $\mathbb{R}^3$.

$$\int_C x^2 y \, dx + (2x - z) \, dy + xy^3 z \, dz$$

C , a curve in $\mathbb{R}^3$.

$$\int_C x^2 y \, dx + (2x - z) \, dy + xy^3 z \, dz$$




$$x^2 y$$

$$2x - z$$

$$xy^3 z$$


C , a curve in $\mathbb{R}^3$.

$$\int_C x^2 y \, dx + (2x - z) \, dy + xy^3 z \, dz$$


$$(x^2 y, 2x - z, xy^3 z)$$

C , a curve in $\mathbb{R}^3$.

$$\int_C x^2 y \, dx + (2x - z) \, dy + xy^3 z \, dz$$


$$\vec{F}(x, y, z) = (x^2 y, 2x - z, xy^3 z)$$

C , a curve in $\mathbb{R}^3$.

$$\int_C x^2 y \, dx + (2x - z) \, dy + xy^3 z \, dz$$

$$\vec{F}(x, y, z) = (x^2 y, 2x - z, xy^3 z)$$

C , a curve in $\mathbb{R}^3$.

$$\vec{F}(x,y,z) = (x^2y, 2x-z, xy^3z)$$


C , a curve in $\mathbb{R}^3$.

$$\int_C dx + dy + dz$$

$$\vec{F}(x,y,z) = (x^2y, 2x-z, xy^3z)$$

C , a curve in $\mathbb{R}^3$.

$$\int_C x^2 y \, dx + (2x - z) \, dy + xy^3 z \, dz$$

$$\vec{F}(x, y, z) = (x^2 y, 2x - z, xy^3 z)$$


C , a curve in $\mathbb{R}^3$.

$$\int_C x^2 y \, dx + (2x - z) \, dy + xy^3 z \, dz$$

$$\vec{F}(x, y, z) = (x^2 y, 2x - z, xy^3 z)$$

C , a curve in $\mathbb{R}^3$.

Write the line integral
corresponding to C and
the vector field

$$\vec{F}(x,y,z) = (5xy, x-z, yz^2).$$

C , a curve in $\mathbb{R}^3$.

Write the line integral
corresponding to C and
the vector field

$$\vec{F}(x, y, z) = (5xy, x - z, yz^2).$$

$$\int_C 5xy \, dx + (x - z) \, dy + yz^2 \, dz$$

C , a curve in $\mathbb{R}^3$.

Write the line integral corresponding to C and the vector field

$$\vec{F}(x, y, z) = (5xy, x - z, yz^2).$$

$$\int_C 5xy \, dx + (x - z) \, dy + yz^2 \, dz$$

C , a curve in $\mathbb{R}^2$.

Write the vector field corresponding to the line integral

$$\int_C x^3 y^2 \, dx + (x^2 y - y^2 x) \, dy.$$

C , a curve in $\mathbb{R}^3$.

Write the line integral corresponding to C and the vector field

$$\vec{F}(x, y, z) = (5xy, x - z, yz^2).$$

$$\int_C 5xy \, dx + (x - z) \, dy + yz^2 \, dz$$

C , a curve in $\mathbb{R}^2$.

Write the vector field corresponding to the line integral

$$\int_C x^3 y^2 \, dx + (x^2 y - y^2 x) \, dy.$$

$$\vec{F}(x, y) = (x^3 y^2, x^2 y - y^2 x)$$

C , a curve in $\mathbb{R}^3$.

$$\int_C x^2 y \, dx + (2x - z) \, dy + xy^3 z \, dz$$

$$\vec{F}(x, y, z) = (x^2 y, 2x - z, xy^3 z)$$

C , a curve in $\mathbb{R}^3$.

$$\int_C x^2 y \, dx + (2x - z) \, dy + xy^3 z \, dz$$

$$\vec{F}(x, y, z) = (\underbrace{x^2 y}_{F_1}, \underbrace{2x - z}_{F_2}, \underbrace{xy^3 z}_{F_3})$$

C , a curve in $\mathbb{R}^3$.

$$\int_C \underset{F_1}{F_1} dx + \underset{F_2}{F_2} dy + \underset{F_3}{F_3} dz$$

$$\vec{F}(x,y,z) = (\underset{F_1}{x^2 y}, \underset{F_2}{2x-z}, \underset{F_3}{xy^3 z})$$

C , a curve in $\mathbb{R}^3$.

$$\int_C F_1 dx_1 + F_2 dx_2 + F_3 dx_3$$

$$\vec{F}(x,y,z) = (\underbrace{x^2 y}_{F_1}, \underbrace{2x-z}_{F_2}, \underbrace{xy^3 z}_{F_3})$$

C , a curve in $\mathbb{R}^3$.

$$\int_C F_1 dx_1 + F_2 dx_2 + F_3 dx_3 = \int_C \sum_{i=1}^3 F_i dx_i$$

$$\vec{F}(x,y,z) = (\underbrace{x^2 y}_{F_1}, \underbrace{2x-z}_{F_2}, \underbrace{xy^3 z}_{F_3})$$

If C , a curve in $\mathbb{R}^n$, and

$$\vec{F}: \mathbb{R}^n \rightarrow \mathbb{R}^n, \quad \vec{F} = (F_1, \dots, F_n),$$

then we can take the line integral

$$\int_C \sum_{i=1}^n F_i dx_i$$

If C , a curve in $\mathbb{R}^n$, and

$$\vec{F}: \mathbb{R}^n \rightarrow \mathbb{R}^n, \quad \vec{F} = (F_1, \dots, F_n),$$

then we can take the line integral

$$\int_C \sum_{i=1}^n F_i dx_i$$

Special case of above:

$$f: \mathbb{R}^n \rightarrow \mathbb{R}$$

$$\nabla f: \mathbb{R}^n \rightarrow \mathbb{R}^n$$

$$\nabla f = \left(\frac{\partial f}{\partial x_1}, \dots, \frac{\partial f}{\partial x_n} \right)$$

$$\int_C \sum_{i=1}^n \frac{\partial f}{\partial x_i} dx_i$$

If C , a curve in $\mathbb{R}^n$, and

$$\vec{F}: \mathbb{R}^n \rightarrow \mathbb{R}^n, \quad \vec{F} = (F_1, \dots, F_n),$$

then we can take the line integral

$$\int_C \sum_{i=1}^n F_i dx_i = \int_C \vec{F} \cdot d\vec{r}$$

Special case of above:

$$f: \mathbb{R}^n \rightarrow \mathbb{R}$$

$$\nabla f: \mathbb{R}^n \rightarrow \mathbb{R}^n$$

$$\nabla f = \left(\frac{\partial f}{\partial x_1}, \dots, \frac{\partial f}{\partial x_n} \right)$$

$$\int_C \sum_{i=1}^n \frac{\partial f}{\partial x_i} dx_i = \int_C \nabla f \cdot d\vec{r}$$

If C , a curve in $\mathbb{R}^n$, and

$$\vec{F}: \mathbb{R}^n \rightarrow \mathbb{R}^n, \quad \vec{F} = (F_1, \dots, F_n),$$

then we can take the line integral

$$\int_C \sum_{i=1}^n F_i dx_i = \int_C \vec{F} \cdot d\vec{r}$$

Special case of above:

$$f: \mathbb{R}^n \rightarrow \mathbb{R}$$

$$\nabla f: \mathbb{R}^n \rightarrow \mathbb{R}^n$$

$$\nabla f = \left(\frac{\partial f}{\partial x_1}, \dots, \frac{\partial f}{\partial x_n} \right)$$


definitions

$$\int_C \sum_{i=1}^n \frac{\partial f}{\partial x_i} dx_i = \int_C \nabla f \cdot d\vec{r}$$

② Line integrals and gradients

Theorem:

C a smooth curve in $\mathbb{R}^n$,
with initial point $\vec{a} \in \mathbb{R}^n$, and
terminal point $\vec{b} \in \mathbb{R}^n$.




If $f: \mathbb{R}^n \rightarrow \mathbb{R}$, then

$$\int_C \sum_{i=1}^n \frac{\partial f}{\partial x_i} dx_i = [f]_{\vec{a}}^{\vec{b}} = f(\vec{b}) - f(\vec{a}).$$

Theorem:

C a smooth curve in $\mathbb{R}^n$,
with initial point $\vec{a} \in \mathbb{R}^n$, and
terminal point $\vec{b} \in \mathbb{R}^n$.



If $f: \mathbb{R}^n \rightarrow \mathbb{R}$, then


$$\int_C \sum_{i=1}^n \frac{\partial f}{\partial x_i} dx_i = [f]_{\vec{a}}^{\vec{b}} = f(\vec{b}) - f(\vec{a}).$$

$$\int_C \nabla f \cdot d\vec{r} = f(\vec{b}) - f(\vec{a})$$

Example: C_1 given by $x=t, y=t, z=t, 0 \leq t \leq 1$.

C_2 given by $x=t^3, y=t^2, z=t, 0 \leq t \leq 1$.

Find $\int_{C_i} yz dx + xz dy + xy dz$ for $i=1,2$.




Example: C_1 given by $x=t, y=t, z=t, 0 \leq t \leq 1$.

C_2 given by $x=t^3, y=t^2, z=t, 0 \leq t \leq 1$.

Find $\int_{C_i} yz dx + xz dy + xy dz$ for $i=1,2$.

$$f(x,y,z) = xyz.$$

$$\nabla f(x,y,z) = (yz, xz, xy).$$



Example: C_1 given by $x=t, y=t, z=t, 0 \leq t \leq 1$.


C_2 given by $x=t^3, y=t^2, z=t, 0 \leq t \leq 1$.

Find $\int_{C_i} yz dx + xz dy + xy dz$ for $i=1,2$.

$$f(x,y,z) = xyz.$$

$$\nabla f(x,y,z) = (yz, xz, xy).$$

$$\int_{C_i} yz dx + xz dy + xy dz$$



Example: C_1 given by $x=t, y=t, z=t, 0 \leq t \leq 1$.


C_2 given by $x=t^3, y=t^2, z=t, 0 \leq t \leq 1$.

Find $\int_{C_i} yz dx + xz dy + xy dz$ for $i=1,2$.

$$f(x,y,z) = xyz.$$

$$\nabla f(x,y,z) = (yz, xz, xy).$$

$$\begin{aligned} \int_{C_i} yz dx + xz dy + xy dz \\ = [xyz]_{(0,0,0)}^{(1,1,1)} \end{aligned}$$



Example: C_1 given by $x=t, y=t, z=t, 0 \leq t \leq 1$.


C_2 given by $x=t^3, y=t^2, z=t, 0 \leq t \leq 1$.

Find $\int_{C_i} yz dx + xz dy + xy dz$ for $i=1,2$.

$$f(x,y,z) = xyz.$$

$$\nabla f(x,y,z) = (yz, xz, xy).$$

$$\begin{aligned} \int_{C_i} yz dx + xz dy + xy dz &= [xyz]_{(0,0,0)}^{(1,1,1)} \\ &= 1 \cdot 1 \cdot 1 - 0 \cdot 0 \cdot 0 \end{aligned}$$



Example: C_1 given by $x=t, y=t, z=t, 0 \leq t \leq 1$.


C_2 given by $x=t^3, y=t^2, z=t, 0 \leq t \leq 1$.

Find $\int_{C_i} yz dx + xz dy + xy dz$ for $i=1,2$.

$$f(x,y,z) = xyz.$$


$$\nabla f(x,y,z) = (yz, xz, xy).$$

$$\begin{aligned} \int_{C_i} yz dx + xz dy + xy dz &= [xyz]_{(0,0,0)}^{(1,1,1)} \\ &= 1 \cdot 1 \cdot 1 - 0 \cdot 0 \cdot 0 \\ &= 1 \end{aligned}$$



Theorem:

C a smooth curve in $\mathbb{R}^n$,
with initial point $\vec{a} \in \mathbb{R}^n$, and
terminal point $\vec{b} \in \mathbb{R}^n$.



If $f: \mathbb{R}^n \rightarrow \mathbb{R}$, then

$$\int_C \sum_{i=1}^n \frac{\partial f}{\partial x_i} dx_i = [f]_{\vec{a}}^{\vec{b}} = f(\vec{b}) - f(\vec{a}).$$

proof:

Suppose C is parametrized as

$$\vec{r}(t) = (x_1(t), x_2(t), \dots, x_n(t)) \quad \text{with } 0 \leq t \leq 1.$$

proof:

Suppose C is parametrized as

$$\vec{r}(t) = (x_1(t), x_2(t), \dots, x_n(t)) \quad \text{with } 0 \leq t \leq 1.$$

$$\int_C \sum_{i=1}^n \frac{\partial f}{\partial x_i} dx_i =$$

proof:

Suppose C is parametrized as

$\vec{r}(t) = (x_1(t), x_2(t), \dots, x_n(t))$ with $0 \leq t \leq 1$.

$$\int_C \sum_{i=1}^n \frac{\partial f}{\partial x_i} dx_i = \int_0^1 \sum_{i=1}^n \frac{\partial f}{\partial x_i} x'_i(t) dt$$

proof:

Suppose C is parametrized as

$$\vec{r}(t) = (x_1(t), x_2(t), \dots, x_n(t)) \quad \text{with } 0 \leq t \leq 1.$$

$$\int_C \sum_{i=1}^n \frac{\partial f}{\partial x_i} dx_i = \int_0^1 \sum_{i=1}^n \frac{\partial f}{\partial x_i} x_i'(t) dt$$

$$= \int_0^1 \sum_{i=1}^n \frac{\partial f}{\partial x_i} \frac{dx_i}{dt} dt$$

proof:

Suppose C is parametrized as

$\vec{r}(t) = (x_1(t), x_2(t), \dots, x_n(t))$ with $0 \leq t \leq 1$.

$$\begin{aligned} \int_C \sum_{i=1}^n \frac{\partial f}{\partial x_i} dx_i &= \int_0^1 \sum_{i=1}^n \frac{\partial f}{\partial x_i} x'_i(t) dt \\ &= \int_0^1 \left(\sum_{i=1}^n \frac{\partial f}{\partial x_i} \frac{dx_i}{dt} \right) dt \end{aligned}$$

proof:

Suppose C is parametrized as

$$\vec{r}(t) = (x_1(t), x_2(t), \dots, x_n(t)) \quad \text{with } 0 \leq t \leq 1.$$

$$\int_C \sum_{i=1}^n \frac{\partial f}{\partial x_i} dx_i = \int_0^1 \sum_{i=1}^n \frac{\partial f}{\partial x_i} x'_i(t) dt$$

$$= \int_0^1 \left(\sum_{i=1}^n \frac{\partial f}{\partial x_i} \frac{dx_i}{dt} \right) dt$$

$$= \int_0^1 \frac{d(f \circ \vec{r})}{dt} dt$$

proof:

Suppose C is parametrized as

$$\vec{r}(t) = (x_1(t), x_2(t), \dots, x_n(t)) \quad \text{with } 0 \leq t \leq 1.$$

$$\int_C \sum_{i=1}^n \frac{\partial f}{\partial x_i} dx_i = \int_0^1 \sum_{i=1}^n \frac{\partial f}{\partial x_i} x'_i(t) dt$$

$$= \int_0^1 \left(\sum_{i=1}^n \frac{\partial f}{\partial x_i} \frac{dx_i}{dt} \right) dt$$

$$= \int_0^1 \frac{d(f \circ \vec{r})}{dt} dt$$

$$= [f \circ \vec{r}]'_0$$

proof:

Suppose C is parametrized as

$$\vec{r}(t) = (x_1(t), x_2(t), \dots, x_n(t)) \quad \text{with } 0 \leq t \leq 1.$$

$$\int_C \sum_{i=1}^n \frac{\partial f}{\partial x_i} dx_i = \int_0^1 \sum_{i=1}^n \frac{\partial f}{\partial x_i} x'_i(t) dt$$

$$= \int_0^1 \left(\sum_{i=1}^n \frac{\partial f}{\partial x_i} \frac{dx_i}{dt} \right) dt$$

$$= \int_0^1 \frac{d(f \circ \vec{r})}{dt} dt$$

$$= [f \circ \vec{r}]'_0$$

$$= f \circ \vec{r}(1) - f \circ \vec{r}(0)$$

proof:

Suppose C is parametrized as

$$\vec{r}(t) = (x_1(t), x_2(t), \dots, x_n(t)) \quad \text{with } 0 \leq t \leq 1.$$

$$\int_C \sum_{i=1}^n \frac{\partial f}{\partial x_i} dx_i = \int_0^1 \sum_{i=1}^n \frac{\partial f}{\partial x_i} x'_i(t) dt$$

$$= \int_0^1 \left(\sum_{i=1}^n \frac{\partial f}{\partial x_i} \frac{dx_i}{dt} \right) dt$$

$$= \int_0^1 \frac{d(f \circ \vec{r})}{dt} dt$$

$$= [f \circ \vec{r}]_0^1$$

$$= f \circ \vec{r}(1) - f \circ \vec{r}(0)$$

$$= f(\vec{b}) - f(\vec{a})$$

proof:

Suppose C is parametrized as

$$\vec{r}(t) = (x_1(t), x_2(t), \dots, x_n(t)) \quad \text{with } 0 \leq t \leq 1.$$

$$\int_C \sum_{i=1}^n \frac{\partial f}{\partial x_i} dx_i = \int_0^1 \sum_{i=1}^n \frac{\partial f}{\partial x_i} x'_i(t) dt$$

$$= \int_0^1 \left(\sum_{i=1}^n \frac{\partial f}{\partial x_i} \frac{dx_i}{dt} \right) dt$$

$$= \int_0^1 \frac{d(f \circ \vec{r})}{dt} dt$$

$$= [f \circ \vec{r}]'_0$$

$$= f \circ \vec{r}(1) - f \circ \vec{r}(0)$$

$$= f(\vec{b}) - f(\vec{a})$$


END

Corollary:

For any $f: \mathbb{R}^n \rightarrow \mathbb{R}$, and any curve C in $\mathbb{R}^n$ with initial point $\vec{a} \in \mathbb{R}^n$ and terminal point $\vec{b} \in \mathbb{R}^n$,

$$\int_C \sum_{i=1}^n \frac{\partial f}{\partial x_i} dx_i = [f]_{\vec{a}}^{\vec{b}}$$

is independent of path.




Corollary:

For any $f: \mathbb{R}^n \rightarrow \mathbb{R}$, and any curve C in $\mathbb{R}^n$ with initial point $\vec{a} \in \mathbb{R}^n$ and terminal point $\vec{b} \in \mathbb{R}^n$,

$$\int_C \sum_{i=1}^n \frac{\partial f}{\partial x_i} dx_i = [f]_{\vec{a}}^{\vec{b}}$$

is independent of path.




We write $\int_{\vec{a}}^{\vec{b}} \sum_{i=1}^n \frac{\partial f}{\partial x_i} dx_i$ for $\int_C \sum_{i=1}^n \frac{\partial f}{\partial x_i} dx_i$.

Problem:

Let C be a curve in $\mathbb{R}^2$ with initial point $(3,1)$ and terminal point $(2,4)$. Find


$$\int_C 2xy \, dx + x^2 \, dy$$



Problem:

Let C be a curve in $\mathbb{R}^2$ with initial point $(3,1)$ and terminal point $(2,4)$. Find

$$\int_C 2xy \, dx + x^2 \, dy$$




Solution:

$$\vec{F}(x,y) = (2xy, x^2) = \nabla f(x,y) \quad \text{where } f(x,y) = x^2y.$$

Problem:

Let C be a curve in $\mathbb{R}^2$ with initial point $(3,1)$ and terminal point $(2,4)$. Find

$$\int_C 2xy \, dx + x^2 \, dy$$



Solution:

$$\vec{F}(x,y) = (2xy, x^2) = \nabla f(x,y) \quad \text{where } f(x,y) = x^2y.$$

$$\text{So } \int_C 2xy \, dx + x^2 \, dy = [x^2y]_{(3,1)}^{(2,4)} = 2^2 \cdot 4 - 3^2 \cdot 1 = 7.$$

Problem:

Find $\int_{(0,\pi)}^{(1,0)} e^x \cos y \, dx - e^x \sin y \, dy$.

Problem:

Find $\int_{(0,\pi)}^{(1,0)} e^x \cos y \, dx - e^x \sin y \, dy$.

Solution:

Integral notation suggests
 $(e^x \cos y, -e^x \sin y)$ is conservative.

Problem:

Find $\int_{(0,\pi)}^{(1,0)} e^x \cos y \, dx - e^x \sin y \, dy$.

Solution:

Integral notation suggests
 $(e^x \cos y, -e^x \sin y)$ is conservative.

Find that the potential function is $e^x \cos y$.

Problem:

Find $\int_{(0,\pi)}^{(1,0)} e^x \cos y \, dx - e^x \sin y \, dy$.

Solution:

Integral notation suggests
 $(e^x \cos y, -e^x \sin y)$ is conservative.

Find that the potential function is $e^x \cos y$.

Then,

$$\int_{(0,\pi)}^{(1,0)} e^x \cos y \, dx - e^x \sin y \, dy = [e^x \cos y]_{(0,\pi)}^{(1,0)}$$

Problem:

Find $\int_{(0,\pi)}^{(1,0)} e^x \cos y \, dx - e^x \sin y \, dy$.

Solution:

Integral notation suggests $(e^x \cos y, -e^x \sin y)$ is conservative.

Find that the potential function is $e^x \cos y$.

Then,

$$\begin{aligned} \int_{(0,\pi)}^{(1,0)} e^x \cos y \, dx - e^x \sin y \, dy &= [e^x \cos y]_{(0,\pi)}^{(1,0)} \\ &= e^1 \cos 0 - e^0 \cos \pi \end{aligned}$$

Problem:

Find $\int_{(0,\pi)}^{(1,0)} e^x \cos y \, dx - e^x \sin y \, dy$.


Solution:

Integral notation suggests
 $(e^x \cos y, -e^x \sin y)$ is conservative.


Find that the potential function is $e^x \cos y$.

Then,


$$\begin{aligned} \int_{(0,\pi)}^{(1,0)} e^x \cos y \, dx - e^x \sin y \, dy &= [e^x \cos y]_{(0,\pi)}^{(1,0)} \\ &= e^1 \cos 0 - e^0 \cos \pi \\ &= e + 1. \end{aligned}$$



$$[f(x)]_a^b = \int_a^b f'(x) dx$$



$$[f(x)]_a^b = \int_c f'(x) dx$$




$$[f]_{\vec{a}}^{\vec{b}} = \int_C \sum_{i=1}^n \frac{\partial f}{\partial x_i} dx_i$$

③ Theorem about line integrals, vector fields, and gradients

$\int_C \sum_{i=1}^n F_i dx_i$ is independent of path if

$$\int_{C_1} \sum_{i=1}^n F_i dx_i = \int_{C_2} \sum_{i=1}^n F_i dx_i$$

whenever C_1 and C_2
have the same initial point
and the same terminal point.




If C is a simple closed curve,
it's common to write $\oint_C \sum_{i=1}^n F_i dx_i$
instead of $\int_C \sum_{i=1}^n F_i dx_i$.

If C is a simple closed curve,
it's common to write $\oint_C \sum_{i=1}^n F_i dx_i$
instead of $\int_C \sum_{i=1}^n F_i dx_i$.

Example:

C param. by
 $x = \cos t$,
 $y = \sin t$,
 $0 \leq t \leq 2\pi$.



$$\oint_C x^2 dx - xy dy$$

Theorem:

$$\vec{F}: \mathbb{R}^n \rightarrow \mathbb{R}^n, \quad \vec{F} = (F_1, \dots, F_n).$$

The following are equivalent:

① $\vec{F} = \nabla f.$

② $\frac{\partial F_i}{\partial x_j} = \frac{\partial F_j}{\partial x_i}$ for $1 \leq i, j \leq n.$

③ $\oint_C \sum_{i=1}^n F_i dx_i = 0$ for simple closed curves $C.$

④ $\int_C \sum_{i=1}^n F_i dx_i$ is independent of path.

$$\vec{F} = \nabla f$$

①



②

$$\frac{\partial F_i}{\partial x_j} = \frac{\partial F_j}{\partial x_i}$$



③

$$\oint_C \sum_{i=1}^n F_i dx_i = 0$$



④

$$\int_C \sum_{i=1}^n F_i dx_i$$

ind. of path

proof: $(① \Rightarrow ②)$

proof: (① $\Rightarrow$ ②)

Suppose $\vec{F} = \nabla f = \left(\frac{\partial f}{\partial x_1}, \frac{\partial f}{\partial x_2}, \dots, \frac{\partial f}{\partial x_n} \right)$.

proof: (① $\Rightarrow$ ②)

Suppose $\vec{F} = \nabla f = \left(\frac{\partial f}{\partial x_1}, \frac{\partial f}{\partial x_2}, \dots, \frac{\partial f}{\partial x_n} \right)$.

Note that $F_i = \frac{\partial f}{\partial x_i}$.

proof: (① $\Rightarrow$ ②)

Suppose $\vec{F} = \nabla f = \left(\frac{\partial f}{\partial x_1}, \frac{\partial f}{\partial x_2}, \dots, \frac{\partial f}{\partial x_n} \right)$.

Note that $F_i = \frac{\partial f}{\partial x_i}$.

Therefore,

$$\frac{\partial}{\partial x_j} F_i =$$

proof: (① $\Rightarrow$ ②)

Suppose $\vec{F} = \nabla f = \left(\frac{\partial f}{\partial x_1}, \frac{\partial f}{\partial x_2}, \dots, \frac{\partial f}{\partial x_n} \right)$.

Note that $F_i = \frac{\partial f}{\partial x_i}$.

Therefore,

$$\frac{\partial}{\partial x_j} F_i = \frac{\partial}{\partial x_j} \frac{\partial f}{\partial x_i}$$

proof: (① $\Rightarrow$ ②)

Suppose $\vec{F} = \nabla f = \left(\frac{\partial f}{\partial x_1}, \frac{\partial f}{\partial x_2}, \dots, \frac{\partial f}{\partial x_n} \right)$.

Note that $F_i = \frac{\partial f}{\partial x_i}$.

Therefore,

$$\frac{\partial}{\partial x_j} F_i = \frac{\partial}{\partial x_j} \frac{\partial f}{\partial x_i} = \frac{\partial}{\partial x_i} \frac{\partial f}{\partial x_j}$$

proof: (① $\Rightarrow$ ②)

Suppose $\vec{F} = \nabla f = \left(\frac{\partial f}{\partial x_1}, \frac{\partial f}{\partial x_2}, \dots, \frac{\partial f}{\partial x_n} \right)$.

Note that $F_i = \frac{\partial f}{\partial x_i}$.

Therefore,

$$\frac{\partial}{\partial x_j} F_i = \frac{\partial}{\partial x_j} \frac{\partial f}{\partial x_i} = \frac{\partial}{\partial x_i} \frac{\partial f}{\partial x_j} = \frac{\partial}{\partial x_i} F_j.$$

proof: (① $\Rightarrow$ ②)

Suppose $\vec{F} = \nabla f = \left(\frac{\partial f}{\partial x_1}, \frac{\partial f}{\partial x_2}, \dots, \frac{\partial f}{\partial x_n} \right)$.

Note that $F_i = \frac{\partial f}{\partial x_i}$.


Therefore,

$$\frac{\partial}{\partial x_j} F_i = \frac{\partial}{\partial x_j} \frac{\partial f}{\partial x_i} = \frac{\partial}{\partial x_i} \frac{\partial f}{\partial x_j} = \frac{\partial}{\partial x_i} F_j.$$

END

$$\vec{F} = \nabla f$$

①



②

$$\frac{\partial F_i}{\partial x_j} = \frac{\partial F_j}{\partial x_i}$$



③

$$\oint_C \sum_{i=1}^n F_i dx_i = 0$$



④

$$\int_C \sum_{i=1}^n F_i dx_i$$

ind. of path



proof: (③ $\Rightarrow$ ④)

proof: (③ $\Rightarrow$ ④)

Suppose $\oint_C \sum_{i=1}^n F_i dx_i = 0$ whenever C is
a simple closed curve.

proof: (3) $\Rightarrow$ (4)

Suppose $\oint_C \sum_{i=1}^n F_i dx_i = 0$ whenever C is a simple closed curve.

Suppose C_1 and C_2 have the same initial and terminal points.




proof: (③ $\Rightarrow$ ④)

Suppose $\oint_C \sum_{i=1}^n F_i dx_i = 0$ whenever C is a simple closed curve.

Suppose C_1 and C_2 have the same initial and terminal points.

Let $C = C_1 \cup -C_2$.



proof: (3) $\Rightarrow$ (4)

Suppose $\oint_C \sum_{i=1}^n F_i dx_i = 0$ whenever C is a simple closed curve.

Suppose C_1 and C_2 have the same initial and terminal points. Let $C = C_1 \cup -C_2$. Then,




proof: (3) $\Rightarrow$ (4)

Suppose $\oint_C \sum_{i=1}^n F_i dx_i = 0$ whenever C is a simple closed curve.

Suppose C_1 and C_2 have the same initial and terminal points.

Let $C = C_1 \cup -C_2$. Then,

$$0 = \int_C \sum_{i=1}^n F_i dx_i$$



proof: (3) $\Rightarrow$ (4)

Suppose $\oint_C \sum_{i=1}^n F_i dx_i = 0$ whenever C is a simple closed curve.

Suppose C_1 and C_2 have the same initial and terminal points.

Let $C = C_1 \cup -C_2$. Then,



$$0 = \int_C \sum_{i=1}^n F_i dx_i$$


$$= \int_{C_1} \sum_{i=1}^n F_i dx_i + \int_{-C_2} \sum_{i=1}^n F_i dx_i$$

proof: (3) $\Rightarrow$ (4)

Suppose $\oint_C \sum_{i=1}^n F_i dx_i = 0$ whenever C is a simple closed curve.

Suppose C_1 and C_2 have the same initial and terminal points.

Let $C = C_1 \cup -C_2$. Then,



$$0 = \int_C \sum_{i=1}^n F_i dx_i$$

$$= \int_{C_1} \sum_{i=1}^n F_i dx_i + \int_{-C_2} \sum_{i=1}^n F_i dx_i$$


$$= \int_{C_1} \sum_{i=1}^n F_i dx_i - \int_{C_2} \sum_{i=1}^n F_i dx_i.$$

proof: (3) $\Rightarrow$ (4)

Suppose $\oint_C \sum_{i=1}^n F_i dx_i = 0$ whenever C is a simple closed curve.

Suppose C_1 and C_2 have the same initial and terminal points.

Let $C = C_1 \cup -C_2$. Then,



$$0 = \int_C \sum_{i=1}^n F_i dx_i$$

$$= \int_{C_1} \sum_{i=1}^n F_i dx_i + \int_{-C_2} \sum_{i=1}^n F_i dx_i$$

$$= \int_{C_1} \sum_{i=1}^n F_i dx_i - \int_{C_2} \sum_{i=1}^n F_i dx_i.$$

So

$$\int_{C_1} \sum_{i=1}^n F_i dx_i = \int_{C_2} \sum_{i=1}^n F_i dx_i.$$

proof: (3) $\Rightarrow$ (4)

Suppose $\oint_C \sum_{i=1}^n F_i dx_i = 0$ whenever C is a simple closed curve.

Suppose C_1 and C_2 have the same initial and terminal points.



Let $C = C_1 \cup -C_2$. Then,

$$0 = \int_C \sum_{i=1}^n F_i dx_i$$

$$= \int_{C_1} \sum_{i=1}^n F_i dx_i + \int_{-C_2} \sum_{i=1}^n F_i dx_i$$

$$= \int_{C_1} \sum_{i=1}^n F_i dx_i - \int_{C_2} \sum_{i=1}^n F_i dx_i.$$


So

$$\int_{C_1} \sum_{i=1}^n F_i dx_i = \int_{C_2} \sum_{i=1}^n F_i dx_i.$$

END

$$\vec{F} = \nabla f$$

①



②

$$\frac{\partial F_i}{\partial x_j} = \frac{\partial F_j}{\partial x_i}$$



③

$$\oint_C \sum_{i=1}^n F_i dx_i = 0$$



④

$$\int_C \sum_{i=1}^n F_i dx_i$$


ind. of path



Lemma:

If C is a curve
with a horizontal end,
and terminal point (x, y) ,
then


$$\frac{\partial}{\partial x} \left(\int_C f(x, y) dx + g(x, y) dy \right) = f(x, y).$$



Lemma:

If C is a curve
with a vertical end,
and terminal point (x, y) ,
then


$$\frac{\partial}{\partial y} \left(\int_C f(x, y) dx + g(x, y) dy \right) = g(x, y).$$



Lemma:

If C is a curve with a horizontal end, and terminal point (x, y) , then


$$\frac{\partial}{\partial x} \left(\int_C F_1(x, y) dx + F_2(x, y) dy \right) = F_1(x, y).$$



Lemma:

If C is a curve with a vertical end, and terminal point (x, y) , then


$$\frac{\partial}{\partial y} \left(\int_C F_1(x, y) dx + F_2(x, y) dy \right) = F_2(x, y).$$



Lemma:

If C is a curve with a horizontal end, and terminal point (x, y) , then


$$\frac{\partial}{\partial x} \left(\int_C F_1 dx + F_2 dy \right) = F_1.$$



Lemma:

If C is a curve with a vertical end, and terminal point (x, y) , then

$$\frac{\partial}{\partial y} \left(\int_C F_1 dx + F_2 dy \right) = F_2.$$



proof: $(4 \Rightarrow 1)$ (For $n=2$.)


proof: (④ $\Rightarrow$ ①) (For $n=2$.)

Suppose $\int_C F_1 dx + F_2 dy$ is independent of path.

proof: (④ $\Rightarrow$ ①) (For $n=2$.)

Suppose $\int_C F_1 dx + F_2 dy$ is independent of path.

Let $f: \mathbb{R}^2 \rightarrow \mathbb{R}$ be $f(x, y) = \int_{(0,0)}^{(x,y)} F_1 dx + F_2 dy$.




proof: $(4) \Rightarrow (1)$ (For $n=2$.)

Suppose $\int_C F_1 dx + F_2 dy$ is independent of path.

Let $f: \mathbb{R}^2 \rightarrow \mathbb{R}$ be $f(x, y) = \int_{(0,0)}^{(x,y)} F_1 dx + F_2 dy$.

Note,

$$\frac{\partial}{\partial x} (f(x, y)) = \frac{\partial}{\partial x} \left(\int_{(0,0)}^{(x,y)} F_1 dx + F_2 dy \right)$$




proof: $(4) \Rightarrow (1)$ (For $n=2$.)

Suppose $\int_C F_1 dx + F_2 dy$ is independent of path.

Let $f: \mathbb{R}^2 \rightarrow \mathbb{R}$ be $f(x, y) = \int_{(0,0)}^{(x,y)} F_1 dx + F_2 dy$.

Note,

$$\begin{aligned} \frac{\partial}{\partial x} (f(x, y)) &= \frac{\partial}{\partial x} \left(\int_{(0,0)}^{(x,y)} F_1 dx + F_2 dy \right) \\ &= \frac{\partial}{\partial x} \left(\int_{C_1} F_1 dx + F_2 dy \right) \end{aligned}$$




proof: $(4) \Rightarrow (1)$ (For $n=2$.)

Suppose $\int_C F_1 dx + F_2 dy$ is independent of path.

Let $f: \mathbb{R}^2 \rightarrow \mathbb{R}$ be $f(x, y) = \int_{(0,0)}^{(x,y)} F_1 dx + F_2 dy$.

Note,

$$\begin{aligned} \frac{\partial}{\partial x} (f(x, y)) &= \frac{\partial}{\partial x} \left(\int_{(0,0)}^{(x,y)} F_1 dx + F_2 dy \right) \\ &= \frac{\partial}{\partial x} \left(\int_{C_1} F_1 dx + F_2 dy \right) \\ &= F_1(x, y). \end{aligned}$$



proof: (④ $\Rightarrow$ ①) (For $n=2$.)


Suppose $\int_C F_1 dx + F_2 dy$ is independent of path.

Let $f: \mathbb{R}^2 \rightarrow \mathbb{R}$ be $f(x, y) = \int_{(0,0)}^{(x,y)} F_1 dx + F_2 dy$.

Note,

$$\begin{aligned} \frac{\partial}{\partial x} (f(x, y)) &= \frac{\partial}{\partial x} \left(\int_{(0,0)}^{(x,y)} F_1 dx + F_2 dy \right) \\ &= \frac{\partial}{\partial x} \left(\int_{C_1} F_1 dx + F_2 dy \right) \\ &= F_1(x, y). \end{aligned}$$

Similarly, using C_2 ,




proof: $(4) \Rightarrow (1)$ (For $n=2$.)

Suppose $\int_C F_1 dx + F_2 dy$ is independent of path.

Let $f: \mathbb{R}^2 \rightarrow \mathbb{R}$ be $f(x, y) = \int_{(0,0)}^{(x,y)} F_1 dx + F_2 dy$.

Note,

$$\begin{aligned} \frac{\partial}{\partial x} (f(x, y)) &= \frac{\partial}{\partial x} \left(\int_{(0,0)}^{(x,y)} F_1 dx + F_2 dy \right) \\ &= \frac{\partial}{\partial x} \left(\int_{C_1} F_1 dx + F_2 dy \right) \\ &= F_1(x, y). \end{aligned}$$



Similarly, using C_2 ,

$$\nabla f = \left(\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y} \right)$$


proof: $(4) \Rightarrow (1)$ (For $n=2$.)

Suppose $\int_C F_1 dx + F_2 dy$ is independent of path.

Let $f: \mathbb{R}^2 \rightarrow \mathbb{R}$ be $f(x, y) = \int_{(0,0)}^{(x,y)} F_1 dx + F_2 dy$.

Note,

$$\begin{aligned} \frac{\partial}{\partial x} (f(x, y)) &= \frac{\partial}{\partial x} \left(\int_{(0,0)}^{(x,y)} F_1 dx + F_2 dy \right) \\ &= \frac{\partial}{\partial x} \left(\int_{C_1} F_1 dx + F_2 dy \right) \\ &= F_1(x, y). \end{aligned}$$



Similarly, using C_2 ,

$$\nabla f = \left(\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y} \right) = (F_1,)$$


proof: $(4) \Rightarrow (1)$ (For $n=2$.)

Suppose $\int_C F_1 dx + F_2 dy$ is independent of path.

Let $f: \mathbb{R}^2 \rightarrow \mathbb{R}$ be $f(x, y) = \int_{(0,0)}^{(x,y)} F_1 dx + F_2 dy$.

Note,

$$\begin{aligned} \frac{\partial}{\partial x} (f(x, y)) &= \frac{\partial}{\partial x} \left(\int_{(0,0)}^{(x,y)} F_1 dx + F_2 dy \right) \\ &= \frac{\partial}{\partial x} \left(\int_{C_1} F_1 dx + F_2 dy \right) \\ &= F_1(x, y). \end{aligned}$$



Similarly, using C_2 ,

$$\nabla f = \left(\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y} \right) = (F_1, F_2)$$


proof: $(4) \Rightarrow (1)$ (For $n=2$.)

Suppose $\int_C F_1 dx + F_2 dy$ is independent of path.

Let $f: \mathbb{R}^2 \rightarrow \mathbb{R}$ be $f(x, y) = \int_{(0,0)}^{(x,y)} F_1 dx + F_2 dy$.

Note,

$$\begin{aligned} \frac{\partial}{\partial x} (f(x, y)) &= \frac{\partial}{\partial x} \left(\int_{(0,0)}^{(x,y)} F_1 dx + F_2 dy \right) \\ &= \frac{\partial}{\partial x} \left(\int_{C_1} F_1 dx + F_2 dy \right) \\ &= F_1(x, y). \end{aligned}$$



Similarly, using C_2 ,

$$\nabla f = \left(\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y} \right) = (F_1, F_2) = \vec{F}.$$


proof: (④ $\Rightarrow$ ①) (For $n=2$.)

Suppose $\int_C F_1 dx + F_2 dy$ is independent of path.

Let $f: \mathbb{R}^2 \rightarrow \mathbb{R}$ be $f(x, y) = \int_{(0,0)}^{(x,y)} F_1 dx + F_2 dy$.

Note,

$$\begin{aligned} \frac{\partial}{\partial x} (f(x, y)) &= \frac{\partial}{\partial x} \left(\int_{(0,0)}^{(x,y)} F_1 dx + F_2 dy \right) \\ &= \frac{\partial}{\partial x} \left(\int_{C_1} F_1 dx + F_2 dy \right) \\ &= F_1(x, y). \end{aligned}$$



Similarly, using C_2 ,

$$\nabla f = \left(\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y} \right) = (F_1, F_2) = \vec{F}.$$

END

$$\vec{F} = \nabla f$$

①



②

$$\frac{\partial F_i}{\partial x_j} = \frac{\partial F_j}{\partial x_i}$$



③

$$\oint_C \sum_{i=1}^n F_i dx_i = 0$$



④

$$\int_C \sum_{i=1}^n F_i dx_i$$

ind. of path

Theorem:

$$\vec{F}: \mathbb{R}^n \rightarrow \mathbb{R}^n, \quad \vec{F} = (F_1, \dots, F_n).$$

The following are equivalent:

① $\vec{F} = \nabla f.$

② $\frac{\partial F_i}{\partial x_j} = \frac{\partial F_j}{\partial x_i}$ for $1 \leq i, j \leq n.$

③ $\oint_C \sum_{i=1}^n F_i dx_i = 0$ for simple closed curves $C.$

④ $\int_C \sum_{i=1}^n F_i dx_i$ is independent of path.