


Mathematics of Frozen Seas

Ken Golden, University of Utah

Introduction to Sea Ice & its Microstructure



SLMath Summer School
UAF June 16, 2025

sea ice






semiconductor



invisibility cloak

Sea ice is a multiscale composite material.





micro


meso

macro



metamaterials




sea ice



human bone



cross-pollination



SEA ICE covers ~12% of Earth's ocean surface


- boundary between ocean and atmosphere
- mediates exchange of heat, gases, momentum
- global ocean circulation
- hosts rich ecosystem
- indicator of **climate change**



polar ice caps critical to global climate in reflecting incoming solar radiation



white snow and ice
reflect




dark water and land
absorb

$$\text{albedo } \alpha = \frac{\text{reflected sunlight}}{\text{incident sunlight}}$$

Arctic sea ice extent

September 15, 2020






*recent losses
in comparison to
the United States*

Perovich




ARCTIC summer sea ice loss



predictions require lots of math modeling

ANTARCTICA

southern cryosphere



**Weddell
Sea**

***East Antarctic
Ice Sheet***

***West Antarctic
Ice Sheet***

Ross Sea


sea ice

New Record Low for Antarctic Sea Ice

February 13, 2023

**Much of Antarctica
warmer than average**


Mean 2022 surface air temp
compared with 1991-2022 ($^{\circ}\text{C}$)



Source: ECMWF ERA5

BBC

**Minimum extent 1979-2023
(million sq km)**



Five-day rolling average of sea-ice extent


Source: National Snow and Ice Data Center (NSIDC)

BBC

sea ice formation








effect of Langmuir circulation
on grease and pancake ice






Polynyas

Size: 100 m - 1000 km


Two mechanisms can contribute to keeping polynyas open:

1. **Latent heat (or coastal) polynyas:** **Mertz Glacier Polynya**
Sea ice grows in open-water and is continually removed by winds and currents (e.g. katabatic winds)
 - latent heat released to the ocean during ice formation perpetuates the process
2. **Sensible heat (or open-ocean) polynyas:** **Weddell Polynya**
Upwelling warm waters, vertical heat diffusion, or convection may provide enough oceanic heat flux to maintain ice-free region





polynyas ice factories

Mertz Glacier Polynya, located in East Antarctica, covers only 0.001% of the overall Antarctic sea ice zone at its maximum winter extent, but is responsible for 1% of the total sea ice production in the Southern Ocean.




Buchanan Bay





Weddell Polynya



Antarctic Zone Flux Experiment (ANZFLUX) 1994

ocean swells propagating through a vast field of pancake ice


HOMOGENIZATION: long wave sees an effective medium, not individual floes, like long EM wave interacting with brine inclusion microstructure



pancake ice forming in a wave field in the Southern Ocean



pancake ice



“Dynamic” duo

- Fast
- Rough



- Slow
- Smooth



Dynamics



Thermodynamics

sea ice dynamics
plate tectonics on a fast time scale





measuring ice depth in ridges off Barrow, AK



dynamic sea ice

Dynamics



Momentum equation:

Ice acceleration = wind stress + ocean stress - Coriolis force
- sea surface tilt + internal ice stress


leads



heat flows directly from ocean to atmosphere




Thermodynamics: 4 ways to melt




Top, bottom, lateral, internal

Heat budgets



$$\text{Net shortwave} + \text{incoming longwave} + \text{outgoing longwave} + \text{sensible} + \text{evaporative} + \text{conduction} = \text{melt / freeze}$$



sea ice components of GCM's

What are the key ingredients -- or **governing equations** that need to be solved on grids using powerful computers?


1. Ice thickness distribution $g(x, y, h, t)$ evolution equation **dynamics** + **thermodynamics**
(Thorndike et al. 1975)

$$\frac{Dg}{Dt} = -g \nabla \cdot \mathbf{u} + \Psi(g) - \frac{\partial}{\partial h} (\tau g) + \mathcal{L}$$

**nonlinear PDE with
ice velocity field**

**ice growth
ice melting**

**mechanical redistribution
- ridging and opening**



2. Conservation of momentum, stress vs. strain relation (Hibler 1979)

$$m \frac{D\mathbf{u}}{Dt} = -m f \mathbf{k} \times \mathbf{u} + \boldsymbol{\tau}_a + \boldsymbol{\tau}_o - mg \nabla H + \mathbf{F}_{int} \quad \mathbf{F} = m\mathbf{a} \text{ for sea ice dynamics}$$

3. Heat equation of sea ice and snow

(Maykut and Untersteiner 1971)

$$\frac{\partial T}{\partial t} + \mathbf{u}_{br} \cdot \nabla T = \nabla \cdot k(T) \nabla T$$

thermodynamics

**+ balance of radiative and
thermal fluxes on interfaces**

sea ice and global ocean circulation


brine expulsion from freezing sea water results
in sinking of dense, saltier water to great depths

*melting sea ice inputs fresher water
into the upper ocean*




**deep-water formation drives
circulation in the world's oceans**

Thermohaline Circulation



GLOBAL THERMOHALINE CONVEYOR BELT

Global Climate Models



Climate models are systems of partial differential equations (PDE) derived from the basic laws of physics, chemistry, and fluid motion.

They describe the state of the ocean, ice, atmosphere, land, and their interactions.

The equations are solved on 3-dimensional grids of the air-ice-ocean-land system (with horizontal grid size ~ 50 km), using very powerful computers.

key challenge :

incorporating sub - grid scale processes

linkage of scales

