

COPYRIGHT & LICENSE

THEOREM

*Copyright © 2007 Erin Chamberlain  
Some rights reserved.*

*Theorem 1*

REAL ANALYSIS I

REAL ANALYSIS I

THEOREM

THEOREM

*Theorem 2*

*Theorem 3*

REAL ANALYSIS I

REAL ANALYSIS I

THEOREM

THEOREM

*Theorem 4*

*Theorem 5*

REAL ANALYSIS I

REAL ANALYSIS I

THEOREM

THEOREM

*Theorem 6*

*Theorem 7*

REAL ANALYSIS I

REAL ANALYSIS I

THEOREM

THEOREM

*Theorem 8*

*Theorem 9*

REAL ANALYSIS I

REAL ANALYSIS I

These flashcards and the accompanying L<sup>A</sup>T<sub>E</sub>X source code are licensed under a Creative Commons Attribution–NonCommercial–ShareAlike 3.0 License. For more information, see [creativecommons.org](http://creativecommons.org).

File last updated on Friday 3<sup>rd</sup> August, 2007,  
at 00:50

**Theorem 1.** *Let  $f$  be a continuous function. If  $\int_0^1 f(x) dx \neq 0$ , then there exists a point  $x$  in the interval  $[0, 1]$  such that  $f(x) \neq 0$ .*

**Theorem 3.** *Let  $x$  be a real number. If  $x > 0$ , then  $\frac{1}{x} > 0$ .*

**Theorem 2.** *Let  $x$  be a real number. If  $x > 0$ , then  $\frac{1}{x} > 0$ .*

**Theorem 5.** *Let  $A$  and  $B$  be subsets of a universal set  $U$ . Then  $A \cap (U \setminus B) = A \setminus B$ .*

**Theorem 4.** *Let  $A$  be a set. Then  $\emptyset \subseteq A$ .*

**Theorem 7.** *If  $A$  and  $B$  are subsets of a set  $U$  and  $A^c$  and  $B^c$  are their complements in  $U$ , then*

1.  $(A \cup B)^c = A^c \cap B^c$ .
2.  $(A \cap B)^c = A^c \cup B^c$ .

**Theorem 6.** *Let  $A, B$ , and  $C$  be subsets of a universal set  $U$ . Then the following statements are true.*

1.  $A \cup (U \setminus A) = U$ .
2.  $A \cap (U \setminus A) = \emptyset$ .
3.  $U \setminus (U \setminus A) = A$ .
4.  $A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$ .
5.  $A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$ .
6.  $A \setminus (B \cup C) = (A \setminus B) \cap (A \setminus C)$ .
7.  $A \setminus (B \cap C) = (A \setminus B) \cup (A \setminus C)$ .

**Theorem 9.** *Let  $R$  be an equivalence relation on a set  $S$ . Then  $\{E_x : x \in S\}$  is a partition of  $S$ . The relation “belongs to the same piece as” is the same as  $R$ . Conversely, if  $\mathcal{T}$  is a partition of  $S$ , let  $R$  be defined by  $xRy$  iff  $x$  and  $y$  are in the same piece of the partition. Then  $R$  is an equivalence relation and the corresponding partition into equivalence classes is the same as  $\mathcal{T}$ .*

**Theorem 8.**  $(a, b) = (c, d)$  iff  $a = c$  and  $b = d$ .

THEOREM

*Theorem 10 (part 1)*

REAL ANALYSIS I

THEOREM

*Theorem 11*

REAL ANALYSIS I

THEOREM

*Theorem 13*

REAL ANALYSIS I

THEOREM

*Theorem 15*

REAL ANALYSIS I

THEOREM

*Theorem 17*

REAL ANALYSIS I

THEOREM

*Theorem 10 (part 2)*

REAL ANALYSIS I

THEOREM

*Theorem 12*

REAL ANALYSIS I

THEOREM

*Theorem 14*

REAL ANALYSIS I

THEOREM

*Theorem 16*

REAL ANALYSIS I

THEOREM

*Theorem 18*

REAL ANALYSIS I

**Theorem 10.** Suppose that  $f : A \rightarrow B$ . Let  $C, C_1$  and  $C_2$  be subsets of  $A$  and let  $D, D_1$  and  $D_2$  be subsets of  $B$ . Then the following hold:

6.  $f^{-1}(D_1 \cap D_2) = f^{-1}(D_1) \cap f^{-1}(D_2)$ .
7.  $f^{-1}(D_1 \cup D_2) = f^{-1}(D_1) \cup f^{-1}(D_2)$ .
8.  $f^{-1}(B \setminus D) = A \setminus f^{-1}(D)$ .
9.  $f^{-1}(D_1 \setminus D_2) = f^{-1}(D_1) \setminus f^{-1}(D_2)$  if  $D_2 \subseteq D_1$ .

**Theorem 12.** Let  $f : A \rightarrow B$  and  $g : B \rightarrow C$ . Then

1. If  $f$  and  $g$  are surjective, then  $g \circ f$  is surjective.
2. If  $f$  and  $g$  are injective, then  $g \circ f$  is injective.
3. If  $f$  and  $g$  are bijective, then  $g \circ f$  is bijective.

**Theorem 14.** Let  $f : A \rightarrow B$  and  $g : B \rightarrow C$  be bijective. Then the composition  $g \circ f : A \rightarrow C$  is bijective and  $(g \circ f)^{-1} = f^{-1} \circ g^{-1}$ .

**Theorem 16.** Let  $S$  be a nonempty set. The following three conditions are equivalent:

1.  $S$  is countable.
2. There exists an injection  $f : S \rightarrow \mathbb{N}$ .
3. There exists a surjection  $f : \mathbb{N} \rightarrow S$ .

**Theorem 18.** Let  $S, T$  and  $U$  be sets.

1. If  $S \subseteq T$ , then  $|S| \leq |T|$ .
2.  $|S| \leq |S|$ .
3. If  $|S| \leq |T|$  and  $|T| \leq |U|$ , then  $|S| \leq |U|$ .
4. If  $m, n \in \mathbb{N}$  and  $m \leq n$ , then  $|\{1, 2, \dots, m\}| \leq |\{1, 2, \dots, n\}|$ .
5. If  $S$  is finite, then  $S < \aleph_0$ .

**Theorem 10.** Suppose that  $f : A \rightarrow B$ . Let  $C, C_1$  and  $C_2$  be subsets of  $A$  and let  $D, D_1$  and  $D_2$  be subsets of  $B$ . Then the following hold:

1.  $C \subseteq f^{-1}[f(C)]$ .
2.  $f[f^{-1}(D)] \subseteq D$ .
3.  $f(C_1 \cap C_2) \subseteq f(C_1) \cap f(C_2)$ .
4.  $f(C_1 \cup C_2) = f(C_1) \cup f(C_2)$ .
5.  $f(C_1) \setminus f(C_2) \subseteq f(C_1 \setminus C_2)$  if  $C_2 \subseteq C_1$ .

**Theorem 11.** Suppose that  $f : A \rightarrow B$ . Let  $C, C_1$  and  $C_2$  be subsets of  $A$  and let  $D$  be a subset of  $B$ . Then the following hold:

1. If  $f$  is injective, then  $f^{-1}[f(C)] = C$ .
2. If  $f$  is surjective, then  $f[f^{-1}(D)] = D$ .
3. If  $f$  is injective, then  $f(C_1 \cap C_2) = f(C_1) \cap f(C_2)$ .

**Theorem 13.** Let  $f : A \rightarrow B$  be bijective. Then

1.  $f^{-1} : B \rightarrow A$  is bijective.
2.  $f^{-1} \circ f = i_A$  and  $f \circ f^{-1} = i_B$ .

**Theorem 15.** Let  $S$  be a countable set and let  $T \subseteq S$ . Then  $T$  is countable.

**Theorem 17.** The set  $\mathbb{R}$  of real numbers is uncountable.

THEOREM

*Theorem 19*

REAL ANALYSIS I

THEOREM

*Theorem 21*

REAL ANALYSIS I

THEOREM

*Theorem 23*  
*The Binomial Formula*

REAL ANALYSIS I

THEOREM

*Theorem 25*

REAL ANALYSIS I

THEOREM

*Theorem 27*

REAL ANALYSIS I

THEOREM

*Theorem 20*  
*Principle of Mathematical Induction*

REAL ANALYSIS I

THEOREM

*Theorem 22*

REAL ANALYSIS I

THEOREM

*Theorem 24*

REAL ANALYSIS I

THEOREM

*Theorem 26*

REAL ANALYSIS I

THEOREM

*Theorem 28*

REAL ANALYSIS I

**Theorem 20.** (*Principle of Mathematical Induction*)  
 Let  $P(n)$  be a statement that is either true or false for each  $n \in \mathbb{N}$ . Then  $P(n)$  is true for all  $n \in \mathbb{N}$  provided that

1.  $P(1)$  is true, and
2. for each  $k \in \mathbb{N}$ , if  $P(k)$  is true, then  $P(k+1)$  is true.

**Theorem 22.**  $7^n - 4^n$  is a multiple of 3 for all  $n \in \mathbb{N}$ .

**Theorem 24.** Let  $m \in \mathbb{N}$  and let  $P(n)$  be a statement that is either true or false for each  $n \geq m$ . Then  $P(n)$  is true for all  $n \geq m$  provided that

1.  $P(m)$  is true, and
2. for each  $k \geq m$ , if  $P(k)$  is true, then  $P(k+1)$  is true.

**Theorem 26.** Let  $x, y \in \mathbb{R}$  such that  $x \leq y + \epsilon$  for every  $\epsilon > 0$ . Then  $x \leq y$ .

**Theorem 28.** Let  $m, n, p \in \mathbb{Z}$ . If  $p$  is a prime number and  $p$  divides the product  $mn$ , then  $p$  divides  $m$  or  $p$  divides  $n$ .

**Theorem 19.** For any set  $S$ , we have  $|S| < |\mathcal{P}(S)|$ .

**Theorem 21.**  $1+2+3+\dots+n = \frac{1}{2}n(n+1)$  for every natural number  $n$ .

**Theorem 23.** (*The Binomial Formula*) If  $x$  and  $y$  are real numbers and  $n \in \mathbb{N}$ , then

$$(x+y)^n = \sum_{k=0}^n \binom{n}{k} x^{n-k} y^k.$$

**Theorem 25.** Let  $x, y$ , and  $z$  be real numbers.

1. If  $x+z = y+z$ , then  $x = y$ .
2.  $x \cdot 0 = 0$ .
3.  $(-1) \cdot x = -x$ .
4.  $xy = 0$  iff  $x = 0$  or  $y = 0$ .
5.  $x < y$  iff  $-y < -x$ .
6. If  $x < y$  and  $z < 0$ , then  $xz > yz$ .

**Theorem 27.** Let  $x, y \in \mathbb{R}$  and let  $a \geq 0$ . Then

1.  $|x| \geq 0$ .
2.  $|x| \leq a$  iff  $-a \leq x \leq a$ .
3.  $|xy| = |x| \cdot |y|$ .
4.  $|x+y| \leq |x| + |y|$ . (*The triangle inequality*)

THEOREM

*Theorem 29*

REAL ANALYSIS I

THEOREM

*Theorem 31*

REAL ANALYSIS I

THEOREM

*Theorem 33*

REAL ANALYSIS I

THEOREM

*Theorem 35*

REAL ANALYSIS I

THEOREM

*Theorem 37*

REAL ANALYSIS I

THEOREM

*Theorem 30*

REAL ANALYSIS I

THEOREM

*Theorem 32*

REAL ANALYSIS I

THEOREM

*Theorem 34*

REAL ANALYSIS I

THEOREM

*Theorem 36*  
*Archimedean Property of  $\mathbb{R}$*

REAL ANALYSIS I

THEOREM

*Theorem 38*

REAL ANALYSIS I

**Theorem 30.** Every non-empty subset of  $\mathbb{R}$  that is bounded below has a greatest lower bound.

**Theorem 29.** Let  $p$  be a prime number. Then  $\sqrt{p}$  is not a rational number.

**Theorem 32.** Let  $A$  and  $B$  be non-empty subsets of  $\mathbb{R}$ . Then

1.  $\inf A \leq \sup A$ .
2.  $\sup(-A) = -\inf A$  and  $\inf(-A) = -\sup A$ .
3.  $\sup(A+B) = \sup(A) + \sup(B)$  and  $\inf(A+B) = \inf(A) + \inf(B)$ .
4.  $\sup(A-B) = \sup(A) - \inf(B)$ .
5. If  $A \subseteq B$ , then  $\sup A \leq \sup B$  and  $\inf B \leq \inf A$ .

**Theorem 34.** Let  $f$  and  $g$  be functions defined on a set containing  $A$  as a subset, and let  $c \in \mathbb{R}$  be a positive constant. Then

1.  $\sup_A cf = c \sup_A f$  and  $\inf_A cf = c \inf_A f$ .
2.  $\sup_A(-f) = -\inf_A f$ .
3.  $\sup_A(f+g) \leq \sup_A f + \sup_A g$  and  $\inf_A f + \inf_A g \leq \inf_A(f+g)$ .
4.  $\sup\{f(x) - f(y) : x, y \in A\} \leq \sup_A f - \inf_A f$ .

**Theorem 31.** Let  $A$  be a non-empty subset of  $\mathbb{R}$  and  $x$  an element of  $\mathbb{R}$ . Then

1.  $\sup A \leq x$  iff  $a \leq x$  for every  $a \in A$ .
2.  $x < \sup A$  iff  $x < a$  for some  $a \in A$ .

**Theorem 33.** Suppose that  $D$  is a nonempty set and that  $f : D \rightarrow \mathbb{R}$  and  $g : D \rightarrow \mathbb{R}$ . If for every  $x, y \in D$ ,  $f(x) \leq g(y)$ , then  $f(D)$  is bounded above and  $g(D)$  is bounded below. Furthermore,  $\sup f(D) \leq \sup g(D)$ .

**Theorem 36.** (Archimedean Property of  $\mathbb{R}$ ) The set  $\mathbb{N}$  of natural numbers is unbounded above in  $\mathbb{R}$ .

**Theorem 35.** The real number system  $\mathbb{R}$  is a complete ordered field.

**Theorem 38.** Let  $p$  be a prime number. Then there exists a positive real number  $x$  such that  $x^2 = p$ .

**Theorem 37.** Each of the following is equivalent to the Archimedean property.

1. For each  $z \in \mathbb{R}$ , there exists  $n \in \mathbb{N}$  such that  $n > z$ .
2. For each  $x > 0$  and for each  $y \in \mathbb{R}$ , there exists  $n \in \mathbb{N}$  such that  $nx > y$ .
3. For each  $x > 0$ , there exists  $n \in \mathbb{N}$  such that  $0 < \frac{1}{n} < x$ .



THEOREM

*Theorem 39*

REAL ANALYSIS I

THEOREM

*Theorem 41*

REAL ANALYSIS I

COROLLARY

*Corollary 1*

REAL ANALYSIS I

LEMMA

*Lemma 1*

REAL ANALYSIS I

THEOREM

*Theorem 45*  
*Bolzano–Weierstrass Theorem*

REAL ANALYSIS I

THEOREM

*Theorem 40*

REAL ANALYSIS I

THEOREM

*Theorem 42*

REAL ANALYSIS I

THEOREM

*Theorem 43*

REAL ANALYSIS I

THEOREM

*Theorem 44*  
*Heine–Borel Theorem*

REAL ANALYSIS I

THEOREM

*Theorem 46*

REAL ANALYSIS I

**Theorem 40.** *If  $x$  and  $y$  are real numbers with  $x < y$ , then there exists an irrational number  $w$  such that  $x < w < y$ .*

**Theorem 39.** *(Density of  $\mathbb{Q}$  in  $\mathbb{R}$ ) If  $x$  and  $y$  are real numbers with  $x < y$ , then there exists a rational number  $r$  such that  $x < r < y$ .*

**Theorem 42.**

1. *The union of any collection of open sets is an open set.*
2. *The intersection of any finite collection of open sets is an open set.*

**Theorem 41.**

1. *A set  $S$  is open iff  $S = \text{int } S$ . Equivalently,  $S$  is open iff every point in  $S$  is an interior point of  $S$ .*
2. *A set  $S$  is closed iff its complement  $\mathbb{R} \setminus S$  is open.*

**Theorem 43.** *Let  $S$  be a subset of  $\mathbb{R}$ . Then*

1.  *$S$  is closed iff  $S$  contains all of its accumulation points.*
2.  *$\text{cl } S$  is a closed set.*
3.  *$S$  is closed iff  $S = \text{cl } S$ .*

**Corollary 1.**

1. *The intersection of any collection of closed sets is closed.*
2. *The union of any finite collection of closed sets is closed.*

**Theorem 44.** *(Heine–Borel) A subset  $S$  of  $\mathbb{R}$  is compact iff  $S$  is closed and bounded.*

**Lemma 1.** *If  $S$  is a nonempty closed bounded subset of  $\mathbb{R}$ , then  $S$  has a maximum and a minimum.*

**Theorem 46.** *Let  $\mathcal{F} = \{K_\alpha : \alpha \in \mathcal{A}\}$  be a family of compact subsets of  $\mathbb{R}$ . Suppose that the intersection of any finite subfamily of  $\mathcal{F}$  is nonempty. Then  $\bigcap \{K_\alpha : \alpha \in \mathcal{A}\} \neq \emptyset$ .*

**Theorem 45.** *(Bolzano–Weierstrass) If a bounded subset  $S$  of  $\mathbb{R}$  contains infinitely many points, then there exists at least one point in  $\mathbb{R}$  that is an accumulation point of  $S$ .*

COROLLARY

THEOREM

*Corollary 2*  
*Nested Intervals Theorem*

*Theorem 47*

REAL ANALYSIS I

REAL ANALYSIS I

THEOREM

THEOREM

*Theorem 48*

*Theorem 49*

REAL ANALYSIS I

REAL ANALYSIS I

THEOREM

THEOREM

*Theorem 50*

*Theorem 51*

REAL ANALYSIS I

REAL ANALYSIS I

THEOREM

THEOREM

*Theorem 52*  
*The Squeeze Principle*

*Theorem 53*

REAL ANALYSIS I

REAL ANALYSIS I

THEOREM

COROLLARY

*Theorem 54*

*Corollary 3*

REAL ANALYSIS I

REAL ANALYSIS I

**Theorem 47.** Let  $(s_n)$  and  $(a_n)$  be sequences of real numbers and let  $s \in \mathbb{R}$ . If for some  $k > 0$  and some  $m \in \mathbb{N}$ , we have

$$|s_n - s| \leq k|a_n|, \text{ for all } n > m,$$

and if  $\lim a_n = 0$ , then it follows that  $\lim s_n = s$ .

**Corollary 2.** (Nested Intervals Theorem) Let  $\mathcal{F} = \{A_n : n \in \mathbb{N}\}$  be a family of closed bounded intervals in  $\mathbb{R}$  such that  $A_{n+1} \subseteq A_n$  for all  $n \in \mathbb{N}$ . Then  $\bigcap_{n=1}^{\infty} A_n \neq \emptyset$ .

**Theorem 49.** If a sequence converges, its limit is unique.

**Theorem 48.** Every convergent sequence is bounded.

**Theorem 51.** Let  $(s_n)$  be a sequence of real numbers such that  $\lim s_n = 0$ , and let  $(t_n)$  be a bounded sequence. Then  $\lim s_n t_n = 0$ .

**Theorem 50.** A sequence  $(s_n)$  converges to  $s$  iff for each  $\epsilon > 0$ , there are only finitely many  $n$  for which  $|s_n - s| \geq \epsilon$ .

**Theorem 53.** Suppose that  $(s_n)$  and  $(t_n)$  are convergent sequences with  $\lim s_n = s$  and  $\lim t_n = t$ . Then

1.  $\lim(s_n + t_n) = s + t$ .
2.  $\lim(k s_n) = ks$  and  $\lim(k + s_n) = k + s$  for any  $k \in \mathbb{R}$ .
3.  $\lim(s_n t_n) = st$ .
4.  $\lim\left(\frac{s_n}{t_n}\right) = \frac{s}{t}$ , provided that  $t_n \neq 0$  for all  $n$  and  $t \neq 0$ .

**Theorem 52.** (The Squeeze Principle) If  $(a_n)$ ,  $(b_n)$ , and  $(c_n)$  are sequences for which there is a number  $K$  such that  $b_n \leq a_n \leq c_n$  for all  $n > K$ , and if  $b_n \rightarrow a$  and  $c_n \rightarrow a$ , then  $a_n \rightarrow a$ .

**Corollary 3.** If  $(t_n)$  converges to  $t$  and  $t_n \geq 0$  for all  $n \in \mathbb{N}$ , then  $t \geq 0$ .

**Theorem 54.** Suppose that  $(s_n)$  and  $(t_n)$  are convergent sequences with  $\lim s_n = s$  and  $\lim t_n = t$ . If  $s_n \leq t_n$  for all  $n \in \mathbb{N}$ , then  $s \leq t$ .

THEOREM

*Theorem 55*  
*Ratio Test*

REAL ANALYSIS I

THEOREM

*Theorem 57*

REAL ANALYSIS I

THEOREM

*Theorem 59*

REAL ANALYSIS I

LEMMA

*Lemma 3*

THEOREM

*Theorem 61*

REAL ANALYSIS I

THEOREM

*Theorem 56*

REAL ANALYSIS I

THEOREM

*Theorem 58*  
*Monotone Convergence Theorem*

REAL ANALYSIS I

LEMMA

*Lemma 2*

REAL ANALYSIS I

THEOREM

*Theorem 60*  
*Cauchy Convergence Criterion*

REAL ANALYSIS I

REAL ANALYSIS I

THEOREM

*Theorem 62*  
*Bolzano–Weierstrass Theorem For Sequences*

REAL ANALYSIS I

**Theorem 56.** Suppose that  $(s_n)$  and  $(t_n)$  are sequences such that  $s_n \leq t_n$  for all  $n \in \mathbb{N}$ .

1. If  $\lim s_n = +\infty$ , then  $\lim t_n = +\infty$ .
2. If  $\lim t_n = -\infty$ , then  $\lim s_n = -\infty$ .

**Theorem 58.** (Monotone Convergence Theorem) A monotone sequence is convergent iff it is bounded.

**Lemma 2.** Every convergent sequence is a Cauchy sequence.

**Theorem 60.** (Cauchy Convergence Criterion) A sequence of real numbers is convergent iff it is a Cauchy sequence.

**Theorem 62.** (Bolzano–Weierstrass Theorem For Sequences) Every bounded sequence has a convergent subsequence.

**Theorem 55.** (Ratio Test) Suppose that  $(s_n)$  is a sequence of positive terms and that the limit  $L = \lim \left( \frac{s_{n+1}}{s_n} \right)$  exists. If  $L < 1$ , then  $\lim s_n = 0$ .

**Theorem 57.** Let  $(s_n)$  be a sequence of positive numbers. Then  $\lim s_n = +\infty$  iff  $\lim \left( \frac{1}{s_n} \right) = 0$ .

**Theorem 59.**

1. If  $(s_n)$  is an unbounded increasing sequence, then  $\lim s_n = +\infty$ .
2. If  $(s_n)$  is an unbounded decreasing sequence, then  $\lim s_n = -\infty$ .

**Lemma 3.** Every Cauchy sequence is bounded.

**Theorem 61.** If a sequence  $(s_n)$  converges to a real number  $s$ , then every subsequence of  $(s_n)$  also converges to  $s$ .

THEOREM

*Theorem 63*

REAL ANALYSIS I

THEOREM

*Theorem 65*

REAL ANALYSIS I

COROLLARY

*Corollary 4*

REAL ANALYSIS I

THEOREM

*Theorem 68*

REAL ANALYSIS I

THEOREM

*Theorem 70*

REAL ANALYSIS I

THEOREM

*Theorem 64*

REAL ANALYSIS I

THEOREM

*Theorem 66*

REAL ANALYSIS I

THEOREM

*Theorem 67*

REAL ANALYSIS I

THEOREM

*Theorem 69*

REAL ANALYSIS I

THEOREM

*Theorem 71*

REAL ANALYSIS I

**Theorem 64.** Let  $(s_n)$  be a sequence and suppose that  $m = \lim s_n$  is a real number. Then the following properties hold:

1. For every  $\epsilon > 0$  there exists  $N$  such that  $n > N$  implies that  $s_n < m + \epsilon$ .
2. For every  $\epsilon > 0$  and for every  $i \in \mathbb{N}$ , there exists an integer  $k > i$  such that  $s_k > m - \epsilon$ .

**Theorem 66.** Let  $f : D \rightarrow \mathbb{R}$  and let  $c$  be an accumulation point of  $D$ . Then  $\lim_{x \rightarrow c} f(x) = L$  iff for every sequence  $(s_n)$  in  $D$  that converges to  $c$  with  $s_n \neq c$  for all  $n$ , the sequence  $(f(s_n))$  converges to  $L$ .

**Theorem 67.** Let  $f : D \rightarrow \mathbb{R}$  and let  $c$  be an accumulation point of  $D$ . Then the following are equivalent:

- (a)  $f$  does not have a limit at  $c$ .
- (b) There exists a sequence  $(s_n)$  in  $D$  with each  $s_n \neq c$  such that  $(s_n)$  converges to  $c$ , but  $(f(s_n))$  is not convergent in  $\mathbb{R}$ .

**Theorem 69.** Let  $f : D \rightarrow \mathbb{R}$  and let  $c \in D$ . Then the following three conditions are equivalent:

- (a)  $f$  is continuous at  $c$ .
- (b) If  $(x_n)$  is any sequence in  $D$  such that  $(x_n)$  converges to  $c$ , then  $\lim f(x_n) = f(c)$ .
- (c) For every neighborhood  $V$  of  $f(c)$  there exists a neighborhood  $U$  of  $c$  such that  $f(U \cap D) \subseteq V$ .

Furthermore, if  $c$  is an accumulation point of  $D$ , then the above are all equivalent to

- (d)  $f$  has a limit at  $c$  and  $\lim_{x \rightarrow c} f(x) = f(c)$ .

**Theorem 71.** Let  $f$  and  $g$  be functions from  $D$  to  $\mathbb{R}$ , and let  $c \in D$ . Suppose that  $f$  and  $g$  are continuous at  $c$ . Then

- (a)  $f + g$  and  $fg$  are continuous at  $c$ ,
- (b)  $f/g$  is continuous at  $c$  if  $g(c) \neq 0$ .

**Theorem 63.** Every unbounded sequence contains a monotone subsequence that has either  $+\infty$  or  $-\infty$  as a limit.

**Theorem 65.** Let  $f : D \rightarrow \mathbb{R}$  and let  $c$  be an accumulation point of  $D$ . Then  $\lim_{x \rightarrow c} f(x) = L$  iff for each neighborhood  $V$  of  $L$  there exists a deleted neighborhood  $U$  of  $c$  such that  $f(U \cap D) \subseteq V$ .

**Corollary 4.** If  $f : D \rightarrow \mathbb{R}$  and if  $c$  is an accumulation point of  $D$ , then  $f$  can have only one limit at  $c$ .

**Theorem 68.** Let  $f : D \rightarrow \mathbb{R}$  and  $g : D \rightarrow \mathbb{R}$ , and let  $c$  be an accumulation point of  $D$ . If  $\lim_{x \rightarrow c} f(x) = L$ ,  $\lim_{x \rightarrow c} g(x) = M$ , and  $k \in \mathbb{R}$ , then  $\lim_{x \rightarrow c} (f+g)(x) = L + M$ ,  $\lim_{x \rightarrow c} (fg)(x) = LM$ , and  $\lim_{x \rightarrow c} (kf)(x) = kL$ .

**Theorem 70.** Let  $f : D \rightarrow \mathbb{R}$  and let  $c \in D$ . Then  $f$  is discontinuous at  $c$  iff there exists a sequence  $(x_n)$  in  $D$  such that  $(x_n)$  converges to  $c$  but the sequence  $(f(x_n))$  does not converge to  $f(c)$ .



THEOREM

*Theorem 72*

REAL ANALYSIS I

COROLLARY

*Corollary 5*

REAL ANALYSIS I

THEOREM

*Theorem 74*  
*Intermediate Value Theorem*

REAL ANALYSIS I

THEOREM

*Theorem 76*

REAL ANALYSIS I

THEOREM

*Theorem 78*

REAL ANALYSIS I

THEOREM

*Theorem 73*

REAL ANALYSIS I

LEMMA

*Lemma 4*

REAL ANALYSIS I

THEOREM

*Theorem 75*

REAL ANALYSIS I

THEOREM

*Theorem 77*

REAL ANALYSIS I

THEOREM

*Theorem 79*

REAL ANALYSIS I

**Theorem 73.** Let  $D$  be a compact subset of  $\mathbb{R}$  and suppose that  $f : D \rightarrow \mathbb{R}$  is continuous. Then  $f(D)$  is compact.

**Lemma 4.** Let  $f : [a, b] \rightarrow \mathbb{R}$  be continuous and suppose that  $f(a) < 0 < f(b)$ . Then there exists a point  $c$  in  $(a, b)$  such that  $f(c) = 0$ .

**Theorem 75.** Let  $I$  be a compact interval and suppose that  $f : I \rightarrow \mathbb{R}$  is a continuous function. Then the set  $f(I)$  is a compact interval.

**Theorem 77.** Let  $f : D \rightarrow \mathbb{R}$  be uniformly continuous on  $D$  and suppose that  $(x_n)$  is a Cauchy sequence in  $D$ . Then  $(f(x_n))$  is a Cauchy sequence.

**Theorem 79.** Let  $I$  be an interval containing the point  $c$  and suppose that  $f : I \rightarrow \mathbb{R}$ . Then  $f$  is differentiable at  $c$  iff, for every sequence  $(x_n)$  in  $I \setminus \{c\}$  that converges to  $c$ , the sequence

$$\left( \frac{f(x_n) - f(c)}{x_n - c} \right)$$

converges. Furthermore, if  $f$  is differentiable at  $c$ , then the sequence of quotients above will converge to  $f'(c)$ .

**Theorem 72.** Let  $f : D \rightarrow \mathbb{R}$  and  $g : E \rightarrow \mathbb{R}$  be functions such that  $f(D) \subseteq E$ . If  $f$  is continuous at a point  $c \in D$  and  $g$  is continuous at  $f(c)$ , then the composition  $g \circ f : D \rightarrow \mathbb{R}$  is continuous at  $c$ .

**Corollary 5.** Let  $D$  be a compact subset of  $\mathbb{R}$  and suppose that  $f : D \rightarrow \mathbb{R}$  is continuous. Then  $f$  assumes minimum and maximum values on  $D$ . That is, there exist points  $x_1$  and  $x_2$  in  $D$  such that  $f(x_1) \leq f(x) \leq f(x_2)$  for all  $x \in D$ .

**Theorem 74.** (Intermediate Value Theorem) Suppose that  $f : [a, b] \rightarrow \mathbb{R}$  is continuous. Then  $f$  has the intermediate value property on  $[a, b]$ . That is, if  $k$  is any value between  $f(a)$  and  $f(b)$  [i.e.  $f(a) < k < f(b)$  or  $f(b) < k < f(a)$ ], then there exists  $c \in [a, b]$  such that  $f(c) = k$ .

**Theorem 76.** Suppose that  $f : D \rightarrow \mathbb{R}$  is continuous on a compact set  $D$ . Then  $f$  is uniformly continuous on  $D$ .

**Theorem 78.** A function  $f : (a, b) \rightarrow \mathbb{R}$  is uniformly continuous on  $(a, b)$  iff it can be extended to a function  $\tilde{f}$  that is continuous on  $[a, b]$ .

THEOREM

*Theorem 80*

REAL ANALYSIS I

THEOREM

*Theorem 81 (part 2)*

REAL ANALYSIS I

THEOREM

*Theorem 83*

REAL ANALYSIS I

THEOREM

*Theorem 85  
Mean Value Theorem*

REAL ANALYSIS I

COROLLARY

*Corollary 6*

REAL ANALYSIS I

THEOREM

*Theorem 81 (part 1)*

REAL ANALYSIS I

THEOREM

*Theorem 82  
Chain Rule*

REAL ANALYSIS I

THEOREM

*Theorem 84  
Rolle's Theorem*

REAL ANALYSIS I

THEOREM

*Theorem 86*

REAL ANALYSIS I

THEOREM

*Theorem 87*

REAL ANALYSIS I

**Theorem 81.** Suppose that  $f : I \rightarrow \mathbb{R}$  and  $g : I \rightarrow \mathbb{R}$  are differentiable at  $c \in I$ . Then

- (a) If  $k \in \mathbb{R}$ , then the function  $kf$  is differentiable at  $c$  and  $(kf)'(c) = k \cdot f'(c)$ .
- (b) The function  $f + g$  is differentiable at  $c$  and  $(f + g)'(c) = f'(c) + g'(c)$ .

**Theorem 82.** (Chain Rule) Let  $I$  and  $J$  be intervals in  $\mathbb{R}$ , let  $f : I \rightarrow \mathbb{R}$  and  $g : J \rightarrow \mathbb{R}$ , where  $f(I) \subseteq J$ , and let  $c \in I$ . If  $f$  is differentiable at  $c$  and  $g$  is differentiable at  $f(c)$ , then the composite function  $g \circ f$  is differentiable at  $c$  and  $(g \circ f)'(c) = g'(f(c)) \cdot f'(c)$ .

**Theorem 84.** (Rolle's Theorem) Let  $f$  be a continuous function on  $[a, b]$  that is differentiable on  $(a, b)$  and such that  $f(a) = f(b) = 0$ . Then there exists at least one point  $c \in (a, b)$  such that  $f'(c) = 0$ .

**Theorem 86.** Let  $f$  be continuous on  $[a, b]$  and differentiable on  $(a, b)$ . If  $f'(x) = 0$  for all  $x \in (a, b)$ , then  $f$  is constant on  $[a, b]$ .

**Theorem 87.** Let  $f$  be differentiable on an interval  $I$ . Then

- (a) if  $f'(x) > 0$  for all  $x \in I$ , then  $f$  is strictly increasing on  $I$ , and
- (b) if  $f'(x) < 0$  for all  $x \in I$ , then  $f$  is strictly decreasing on  $I$ .

**Theorem 80.** If  $f : I \rightarrow \mathbb{R}$  is differentiable at a point  $c \in I$ , then  $f$  is continuous at  $c$ .

**Theorem 81.** Suppose that  $f : I \rightarrow \mathbb{R}$  and  $g : I \rightarrow \mathbb{R}$  are differentiable at  $c \in I$ . Then

- (c) (Product Rule) The function  $fg$  is differentiable at  $c$  and  $(fg)'(c) = f(c)g'(c) + f'(c)g(c)$ .
- (d) (Quotient Rule) If  $g(c) \neq 0$ , then the function  $f/g$  is differentiable at  $c$  and

$$\left(\frac{f}{g}\right)'(c) = \frac{g(c)f'(c) - f(c)g'(c)}{[g(c)]^2}.$$

**Theorem 83.** If  $f$  is differentiable on an open interval  $(a, b)$  and if  $f$  assumes its maximum or minimum at a point  $c \in (a, b)$ , then  $f'(c) = 0$ .

**Theorem 85.** (Mean Value Theorem) Let  $f$  be a continuous function on  $[a, b]$  that is differentiable on  $(a, b)$ . Then there exists at least one point  $c \in (a, b)$  such that  $f'(c) = \frac{f(b) - f(a)}{b - a}$ .

**Corollary 6.** Let  $f$  and  $g$  be continuous on  $[a, b]$  and differentiable on  $(a, b)$ . Suppose that  $f'(x) = g'(x)$  for all  $x \in (a, b)$ . Then there exists a constant  $C$  such that  $f = g + C$  on  $[a, b]$ .

THEOREM

*Theorem 88*  
*Intermediate Value Theorem for Derivatives*

REAL ANALYSIS I

THEOREM

*Theorem 90*  
*Cauchy Mean Value Theorem*

REAL ANALYSIS I

THEOREM

*Theorem 92*  
*L'Hospital's Rule*

REAL ANALYSIS I

THEOREM

*Theorem 94*

REAL ANALYSIS I

THEOREM

*Theorem 96*

REAL ANALYSIS I

THEOREM

*Theorem 89*  
*Inverse Function Theorem*

REAL ANALYSIS I

THEOREM

*Theorem 91*  
*L'Hospital's Rule*

REAL ANALYSIS I

THEOREM

*Theorem 93*  
*Taylor's Theorem*

REAL ANALYSIS I

THEOREM

*Theorem 95*

REAL ANALYSIS I

THEOREM

*Theorem 97*

REAL ANALYSIS I

**Theorem 89.** (Inverse Function Theorem) Suppose that  $f$  is differentiable on an interval  $I$  and  $f'(x) \neq 0$  for all  $x \in I$ . Then  $f$  is injective,  $f^{-1}$  is differentiable on  $f(I)$ , and  $(f^{-1})'(y) = \frac{1}{f'(x)}$ , where  $y = f(x)$ .

**Theorem 88.** (Intermediate Value Theorem for Derivatives) Let  $f$  be differentiable on  $[a, b]$  and suppose that  $k$  is a number between  $f'(a)$  and  $f'(b)$ . Then there exists a point  $c \in (a, b)$  such that  $f'(c) = k$ .

**Theorem 91.** (L'Hospital's Rule) Let  $f$  and  $g$  be continuous on  $[a, b]$  and differentiable on  $(a, b)$ . Suppose that  $c \in [a, b]$  and  $f(c) = g(c) = 0$ . Suppose also that  $g'(x) \neq 0$  for  $x \in U$ , where  $U$  is the intersection of  $(a, b)$  and some deleted neighborhood of  $c$ . If  $\lim_{x \rightarrow c} \frac{f'(x)}{g'(x)} = L$ , with  $L \in \mathbb{R}$ , then  $\lim_{x \rightarrow c} \frac{f(x)}{g(x)} = L$ .

**Theorem 90.** (Cauchy Mean Value Theorem) Let  $f$  and  $g$  be functions that are continuous on  $[a, b]$  and differentiable on  $(a, b)$ . Then there exists at least one point  $c \in (a, b)$  such that

$$[f(b) - f(a)]g'(c) = [g(b) - g(a)]f'(c)$$

**Theorem 93.** (Taylor's Theorem) Let  $f$  and its first  $n$  derivatives be continuous on  $[a, b]$  and differentiable on  $(a, b)$ , and let  $x_0 \in [a, b]$ . Then for each  $x \in [a, b]$  with  $x \neq x_0$  there exists a point  $c$  between  $x$  and  $x_0$  such that

$$f(x) = f(x_0) + f'(x_0)(x - x_0) + \frac{f''(x_0)}{2!}(x - x_0)^2 + \dots + \frac{f^{(n)}(x_0)}{n!}(x - x_0)^n + \frac{f^{(n+1)}(c)}{(n+1)!}(x - x_0)^{n+1}.$$

**Theorem 92.** (L'Hospital's Rule) Let  $f$  and  $g$  be differentiable on  $(b, \infty)$ . Suppose that  $\lim_{x \rightarrow \infty} f(x) = \lim_{x \rightarrow \infty} g(x) = \infty$ , and that  $g'(x) \neq 0$  for  $x \in (b, \infty)$ . If  $\lim_{x \rightarrow \infty} \frac{f'(x)}{g'(x)} = L$ , where  $L \in \mathbb{R}$ , then  $\lim_{x \rightarrow \infty} \frac{f(x)}{g(x)} = L$ .

**Theorem 95.** Let  $f$  be a bounded function on  $[a, b]$ . Then  $L(f) \leq U(f)$ .

**Theorem 94.** Let  $f$  be a bounded function on  $[a, b]$ . If  $P$  and  $Q$  are partitions of  $[a, b]$  and  $Q$  is a refinement of  $P$ , then  $L(f, P) \leq L(f, Q) \leq U(f, Q) \leq U(f, P)$ .

**Theorem 97.** Let  $f$  be a monotonic function on  $[a, b]$ . Then  $f$  is integrable.

**Theorem 96.** Let  $f$  be a bounded function on  $[a, b]$ . Then  $f$  is integrable iff for each  $\epsilon > 0$  there exists a partition  $P$  of  $[a, b]$  such that  $U(f, P) - L(f, P) < \epsilon$ .

THEOREM

*Theorem 98*

REAL ANALYSIS I

THEOREM

*Theorem 100*

REAL ANALYSIS I

COROLLARY

*Corollary 7*

REAL ANALYSIS I

THEOREM

*Theorem 103*  
*The Fundamental Theorem of Calculus II*

REAL ANALYSIS I

THEOREM

*Theorem 105*

REAL ANALYSIS I

THEOREM

*Theorem 99*

REAL ANALYSIS I

THEOREM

*Theorem 101*

REAL ANALYSIS I

THEOREM

*Theorem 102*  
*The Fundamental Theorem of Calculus I*

REAL ANALYSIS I

THEOREM

*Theorem 104*

REAL ANALYSIS I

THEOREM

*Theorem 106*  
*Cauchy Criterion for Series*

REAL ANALYSIS I

**Theorem 99.** Let  $f$  and  $g$  be integrable functions on  $[a, b]$  and let  $k \in \mathbb{R}$ . Then

- (a)  $kf$  is integrable and  $\int_a^b kf = k \int_a^b f$ , and  
(b)  $f + g$  is integrable and  $\int_a^b (f + g) = \int_a^b f + \int_a^b g$ .

**Theorem 101.** Suppose that  $f$  is integrable on  $[a, b]$  and  $g$  is continuous on  $[c, d]$ , where  $f([a, b]) \subseteq [c, d]$ . Then  $g \circ f$  is integrable on  $[a, b]$ .

**Theorem 102.** (The Fundamental Theorem of Calculus I) Let  $f$  be integrable on  $[a, b]$ . For each  $x \in [a, b]$  let  $F(x) = \int_a^x f(t) dt$ . Then  $F$  is uniformly continuous on  $[a, b]$ . Furthermore, if  $f$  is continuous at  $c \in [a, b]$ , then  $F$  is differentiable at  $c$  and  $F'(c) = f(c)$ .

**Theorem 104.** Suppose that  $\sum a_n = s$  and  $\sum b_n = t$ . Then  $\sum (a_n + b_n) = s + t$  and  $\sum (ka_n) = ks$ , for every  $k \in \mathbb{R}$ .

**Theorem 106.** (Cauchy Criterion for Series) The infinite series  $\sum a_n$  converges iff for each  $\epsilon > 0$  there exists a number  $N$  such that if  $n \geq m > N$ , then  $|a_m + a_{m+1} + \cdots + a_n| < \epsilon$ .

**Theorem 98.** Let  $f$  be a continuous function on  $[a, b]$ . Then  $f$  is integrable on  $[a, b]$ .

**Theorem 100.** Suppose that  $f$  is integrable on both  $[a, c]$  and  $[c, b]$ . Then  $f$  is integrable on  $[a, b]$ . Furthermore,  $\int_a^b f = \int_a^c f + \int_c^b f$ .

**Corollary 7.** Let  $f$  be integrable on  $[a, b]$ . The  $|f|$  is integrable on  $[a, b]$  and  $\left| \int_a^b f \right| \leq \int_a^b |f|$ .

**Theorem 103.** (The Fundamental Theorem of Calculus II) If  $f$  is differentiable on  $[a, b]$  and  $f'$  is integrable on  $[a, b]$ , then  $\int_a^b f' = f(b) - f(a)$ .

**Theorem 105.** If  $\sum a_n$  is a convergent series, then  $\lim a_n = 0$ .



THEOREM

*Theorem 107*  
*Comparison Test*

REAL ANALYSIS I

THEOREM

*Theorem 109*  
*Ratio Test*

REAL ANALYSIS I

THEOREM

*Theorem 111*  
*Integral Test*

REAL ANALYSIS I

THEOREM

*Theorem 113*

REAL ANALYSIS I

THEOREM

*Theorem 115*

REAL ANALYSIS I

THEOREM

*Theorem 108*

REAL ANALYSIS I

THEOREM

*Theorem 110*  
*Root Test*

REAL ANALYSIS I

THEOREM

*Theorem 112*  
*Alternating Series Test*

REAL ANALYSIS I

THEOREM

*Theorem 114*  
*Ratio Criterion*

REAL ANALYSIS I

THEOREM

*Theorem 116*  
*Weierstrass M-test*

REAL ANALYSIS I

**Theorem 108.** *If a series converges absolutely, then it converges.*

**Theorem 110.** *(Root Test) Given a series  $\sum a_n$ , let  $\alpha = \limsup |a_n|^{\frac{1}{n}}$ .*

1. *If  $\alpha < 1$ , then the series converges absolutely.*
2. *If  $\alpha > 1$ , then the series diverges.*
3. *Otherwise,  $\alpha = 1$  and the test gives no information about convergence or divergence.*

**Theorem 112.** *(Alternating Series Test) If  $(a_n)$  is a decreasing sequence of positive numbers and  $\lim a_n = 0$ , then the series  $\sum (-1)^{n+1} a_n$  converges.*

**Theorem 114.** *(Ratio Criterion) The radius of convergence  $R$  of a power series  $\sum a_n x^n$  is equal to  $\lim \left| \frac{a_n}{a_{n+1}} \right|$ , provided that this limit exists.*

**Theorem 116.** *(Weierstrass M-test) Suppose that  $(f_n)$  is a sequence of functions defined on  $S$  and  $(M_n)$  is a sequence of nonnegative numbers such that  $|f_n(x)| \leq M_n$  for all  $x \in S$  and all  $n \in \mathbb{N}$ . If  $\sum M_n$  converges, then  $\sum f_n$  converges uniformly on  $S$ .*

**Theorem 107.** *(Comparison Test) Let  $\sum a_n$  and  $\sum b_n$  be infinite series of nonnegative terms. That is,  $a_n \geq 0$  and  $b_n \geq 0$  for all  $n$ . Then*

1. *If  $\sum a_n$  converges and  $0 \leq b_n \leq a_n$  for all  $n$ , then  $\sum b_n$  converges.*
2. *If  $\sum a_n = +\infty$  and  $0 \leq a_n \leq b_n$  for all  $n$ , then  $\sum b_n = +\infty$ .*

**Theorem 109.** *(Ratio Test) Let  $\sum a_n$  be a series of nonzero terms.*

1. *If  $\limsup \left| \frac{a_{n+1}}{a_n} \right| < 1$ , then the series converges absolutely.*
2. *If  $\liminf \left| \frac{a_{n+1}}{a_n} \right| > 1$ , then the series diverges.*
3. *Otherwise,  $\liminf \left| \frac{a_{n+1}}{a_n} \right| \leq 1 \leq \limsup \left| \frac{a_{n+1}}{a_n} \right|$  and the test gives no information about convergence or divergence.*

**Theorem 111.** *(Integral Test) Let  $f$  be a continuous function defined on  $[0, \infty)$ , and suppose that  $f$  is positive and decreasing. That is, if  $x_1 < x_2$ , then  $f(x_1) \geq f(x_2) > 0$ . Then the series  $\sum (f(n))$  converges iff  $\lim_{n \rightarrow \infty} \left( \int_1^n f(x) dx \right)$  exists as a real number.*

**Theorem 113.** *Let  $\sum a_n x^n$  be a power series and let  $\alpha = \limsup |a_n|^{\frac{1}{n}}$ . Define  $R$  by*

$$R = \begin{cases} \frac{1}{\alpha} & \text{if } 0 < \alpha < +\infty \\ +\infty & \text{if } \alpha = 0 \\ 0 & \text{if } \alpha = +\infty \end{cases}.$$

*Then the series converges absolutely whenever  $|x| < R$  and diverges whenever  $|x| > R$ . (When  $R = +\infty$  we take this to mean that the series converges absolutely for all real  $x$ . When  $R = 0$  then the series converges only at  $x = 0$ .)*

**Theorem 115.** *Let  $(f_n)$  be a sequence of functions defined on a subset  $S$  of  $\mathbb{R}$ . There exists a function  $f$  such that  $(f_n)$  converges to  $f$  uniformly on  $S$  iff the following condition (called the Cauchy criterion) is satisfied:*

*For every  $\epsilon > 0$  there exists a number  $N$  such that  $|f_n(x) - f_m(x)| < \epsilon$  for all  $x \in S$  and all  $m, n > N$ .*

THEOREM

COROLLARY

*Theorem 117*

*Corollary 8*

REAL ANALYSIS I

REAL ANALYSIS I

THEOREM

COROLLARY

*Theorem 118*

*Corollary 9*

REAL ANALYSIS I

REAL ANALYSIS I

THEOREM

COROLLARY

*Theorem 119*

*Corollary 10*

REAL ANALYSIS I

REAL ANALYSIS I

THEOREM

THEOREM

*Theorem 120*

*Theorem 121*

REAL ANALYSIS I

REAL ANALYSIS I

THEOREM

COROLLARY

*Theorem 122*

*Corollary 11*

REAL ANALYSIS I

REAL ANALYSIS I

**Corollary 8.** Let  $\sum_{n=0}^{\infty} f_n$  be a series of functions defined on a set  $S$ . Suppose that each  $f_n$  is continuous on  $S$  and that the series converges uniformly to a function  $f$  on  $S$ . Then  $f = \sum_{n=0}^{\infty} f_n$  is continuous on  $S$ .

**Corollary 9.** Let  $\sum_{n=0}^{\infty} f_n$  be a series of functions defined on an interval  $[a, b]$ . Suppose that each  $f_n$  is continuous on  $[a, b]$  and that the series converges uniformly to a function  $f$  on  $[a, b]$ . Then  $\int_a^b f(x) dx = \sum_{n=0}^{\infty} \int_a^b f_n(x) dx$ .

**Corollary 10.** Let  $\sum_{n=0}^{\infty} f_n$  be a series of functions that converges to a function  $f$  on an interval  $[a, b]$ . Suppose that for each  $n$ ,  $f'_n$  exists and is continuous on  $[a, b]$  and that the series of derivatives  $\sum_{n=0}^{\infty} f'_n$  is uniformly convergent on  $[a, b]$ . Then  $f'(x) = \sum_{n=0}^{\infty} f'_n(x)$  for all  $x \in [a, b]$ .

**Theorem 121.** Let  $\sum a_n x^n$  be a power series with radius of convergence  $R$ , where  $0 < R \leq +\infty$ . If  $0 < K < R$ , then the power series converges uniformly on  $[-K, K]$ .

**Corollary 11.** Suppose that  $f(x) = \sum_{n=0}^{\infty} a_n x^n$  for  $x \in (-R, R)$ , where  $R > 0$ . Then for each  $k \in \mathbb{N}$ , the  $k$ th derivative  $f^{(k)}$  of  $f$  exists on  $(-R, R)$  and

$$\begin{aligned} f^{(k)}(x) &= \sum_{n=k}^{\infty} \frac{n!}{(n-k)!} a_n x^{n-k} \\ &= k! a_k + (k+1)! a_{k+1} x + \frac{(k+2)!}{2!} a_{k+2} x^2 + \dots \end{aligned}$$

Furthermore,  $f^{(k)}(0) = k! a_k$ .

**Theorem 117.** Let  $(f_n)$  be a sequence of continuous functions defined on a set  $S$  and suppose that  $(f_n)$  converges uniformly on  $S$  to a function  $f : S \rightarrow \mathbb{R}$ . Then  $f$  is continuous on  $S$ .

**Theorem 118.** Let  $(f_n)$  be a sequence of continuous functions defined on an interval  $[a, b]$  and suppose that  $(f_n)$  converges uniformly on  $[a, b]$  to a function  $f$ . Then  $\lim_{n \rightarrow \infty} \int_a^b f_n(x) dx = \int_a^b f(x) dx$ .

**Theorem 119.** Suppose that  $(f_n)$  converges to  $f$  on an interval  $[a, b]$ . Suppose also that each  $f'_n$  exists and is continuous on  $[a, b]$ , and that the sequence  $(f'_n)$  converges uniformly on  $[a, b]$ . Then  $\lim_{n \rightarrow \infty} f'_n(x) = f'(x)$  for each  $x \in [a, b]$ .

**Theorem 120.** There exists a continuous function defined on  $\mathbb{R}$  that is nowhere differentiable.

**Theorem 122.** Suppose that a power series converges to a function  $f$  on  $(-R, R)$ , where  $R > 0$ . Then the series can be differentiated term by term, and the differentiated series converges on  $(-R, R)$  to  $f'$ . That is, if  $f(x) = \sum_{n=0}^{\infty} a_n x^n$ , then  $f'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1}$ , and both series have the same radius of convergence.

COROLLARY

THEOREM

*Corollary 12*

*Theorem 123*

REAL ANALYSIS I

REAL ANALYSIS I

COROLLARY

*Corollary 13*

REAL ANALYSIS I

**Theorem 123.** Let  $\sum_{n=0}^{\infty} a_n x^n$  be a power series with a finite positive radius of convergence  $R$ . If the series converges at  $x = R$ , then it converges uniformly on the interval  $[0, R]$ . Similarly, if the series converges at  $x = -R$ , then it converges uniformly on  $[-R, 0]$ .

**Corollary 12.** If  $\sum_{n=0}^{\infty} a_n x^n = \sum_{n=0}^{\infty} b_n x^n$  for all  $x$  in some interval  $(-R, R)$ , where  $R > 0$ , then  $a_n = b_n$  for all  $n \in \mathbb{N} \cup \{0\}$ .

**Corollary 13.** Let  $f(x) = \sum_{n=0}^{\infty} a_n x^n$  have a finite positive radius of convergence  $R$ . If the series converges at  $x = R$ , then  $f$  is continuous at  $x = R$ . If the series converges at  $x = -R$ , then  $f$  is continuous at  $x = -R$ .