

**LEADING TERM CYCLES OF HARISH-CHANDRA MODULES
AND
PARTIAL ORDERS ON COMPONENTS OF THE SPRINGER FIBER**

PETER E. TRAPA

ABSTRACT. We use the geometry of characteristic cycles of Harish-Chandra modules for a real semisimple Lie group $G_{\mathbb{R}}$ to prove an upper triangularity relationship between two bases of each special representation of a classical Weyl group. One basis consists of Goldie rank polynomials attached to primitive ideals in the enveloping algebra of the complexified Lie algebra \mathfrak{g} ; the other consists of polynomials that measure the Euler characteristic of the restriction of an equivariant line bundle on the flag variety for \mathfrak{g} to an irreducible component of the Springer fiber. While these two bases are defined only using the structure of the complex Lie algebra \mathfrak{g} , the relationship between them is closely tied to the real group $G_{\mathbb{R}}$. More precisely the order leading to the upper triangularity result is a suborder of closure order for the orbits of the complexification of a maximal compact subgroup of $G_{\mathbb{R}}$ on the flag variety for \mathfrak{g} .

1. INTRODUCTION

The main result of this paper concerns a relationship between two natural bases of each special representation of a Weyl group. One basis arises in the study of primitive ideals in enveloping algebras, the other in the complex geometry of the Springer fiber. The relationship we uncover originates in the geometry of characteristic cycles of Harish-Chandra modules. As we detail below, it allows one to transport interesting information between the category of Harish-Chandra modules and the category of highest weight modules.

Before turning to applications, we formulate our main result in more detail. Let \mathfrak{g} denote a complex semisimple Lie algebra, let \mathfrak{B} denote its variety of Borel subalgebras, and fix a base point $\mathfrak{b}_1 = \mathfrak{h}_1 \oplus \mathfrak{n}_1$. (It would be more customary to call the basepoint simply \mathfrak{b} , but shortly \mathfrak{b} will denote an arbitrary Borel subalgebra.) Let W denote the Weyl group of \mathfrak{h}_1 in \mathfrak{g} , and write $\rho \in \mathfrak{h}_1^*$ for the half-sum of the roots of \mathfrak{h}_1 in \mathfrak{b}_1 . Let \mathcal{N}^* denote the nilpotent cone in \mathfrak{g}^* , and fix a complex nilpotent coadjoint orbit \mathcal{O} . Consider the set $\text{Prim}_{\rho}(\mathfrak{g}, \mathcal{O})$ of two-sided ideals in the universal enveloping algebra $U(\mathfrak{g})$ such that: (1) I is the annihilator of a simple $U(\mathfrak{g})$ module X (i.e. I is primitive); (2) I contains the augmentation ideal in the center of $U(\mathfrak{g})$ (i.e. I has infinitesimal character ρ); and (3) with respect to the degree filtration, the associated graded ideal $\text{gr}I$ in the symmetric algebra $S(\mathfrak{g})$ cuts out the closure $\overline{\mathcal{O}}$ of \mathcal{O} (i.e. the associated variety of I is $\overline{\mathcal{O}}$). This set is nonempty if and only if \mathcal{O} is special [BV]. To each primitive ideal I in $U(\mathfrak{g})$ (and hence to each element of $\text{Prim}_{\rho}(\mathfrak{g}, \mathcal{O})$), Joseph attaches a harmonic homogeneous polynomial $p_I \in S(\mathfrak{h}_1^*)$, the so-called Goldie rank polynomial ([J2]); see Section 2.2. Set

$$(1.1) \quad \text{Sp}(\mathcal{O}) := \text{span}_{\mathbb{C}}\{p_I \mid I \in \text{Prim}_{\rho}(\mathfrak{g}, \mathcal{O})\}.$$

Joseph proves that the elements p_I are independent, and that the above span of this basis is an irreducible representation of W ; it is called the special representation of W parametrized by \mathcal{O} .

Now we introduce another basis of $\text{Sp}(\mathcal{O})$. Let $T^*\mathfrak{B}$ denote the cotangent bundle to \mathfrak{B} , and write $\mu : T^*\mathfrak{B} \rightarrow \mathcal{N}^*$ for the moment map. Concretely we may write $T^*\mathfrak{B} = \{(\mathfrak{b}, \xi) \mid \xi \in (\mathfrak{g}/\mathfrak{b})^*\}$ and then

Key words and phrases. characteristic cycles, Harish-Chandra modules, Joseph polynomials.

2000 Mathematics Subject Classification: 22E47.

The author was partially supported by NSF grant DMS-0300106.

μ maps a point (\mathfrak{b}, ξ) to ξ (or more precisely to the image of ξ in \mathfrak{g}^* under the canonical inclusion of $(\mathfrak{g}/\mathfrak{b})^*$ in \mathfrak{g}^*). For $\xi \in \mathcal{N}^*$, let $\mu^{-1}(\xi)$ denote the corresponding fiber, the so-called Springer fiber. Spaltenstein established that $\mu^{-1}(\xi)$ is equidimensional, and in particular that its irreducible components all have the same dimension. Write $\text{Irr}(\mu^{-1}(\xi))$ for the set of irreducible components. Now fix $\lambda \in \mathfrak{h}_1^*$ dominant and integral for the roots of \mathfrak{h}_1 in \mathfrak{b}_1 , and let \mathbb{C}_λ denote the corresponding one-dimensional representation of B_1 . Consider the Borel-Weil line bundle $\mathcal{L}_\lambda := G \times_{B_1} \mathbb{C}_\lambda$, and for any subset $Z \subset \text{Irr}(\mu^{-1}(\xi))$, define $q_Z(\lambda)$ to be the Euler characteristic of \mathcal{L}_λ restricted to Z . The assignment $\lambda \mapsto q_Z(\lambda)$ extends to a homogeneous polynomial $q_Z \in \mathbb{S}(\mathfrak{h}_1)$ sometimes called the Joseph polynomial after its introduction in a different context in [J3]; see also [J4] and the references therein, especially [R]. The polynomials q_Z are not independent for trivial reasons. The component group $A_G(\xi)$ of the centralizer of ξ in G clearly acts on $\text{Irr}(\mu^{-1}(\xi))$. (Here G may be taken to be any complex connected Lie group with Lie algebra \mathfrak{g} ; the action is insensitive to isogeny.) If Z and Z' are actually irreducible components in the same $A_G(\xi)$ orbit, then $q_Z = q_{Z'}$. In fact these are the only dependence relations, however, and the polynomials q_C as C ranges over $A_G(\xi) \backslash \text{Irr}(\mu^{-1}(\xi))$ are independent. Moreover if we set $\mathcal{O} = G \cdot \xi$ and assume this orbit is special, we have

$$\text{Sp}(\mathcal{O}) = \text{span}_{\mathbb{C}}\{q_C \mid C \in A_G(\xi) \backslash \text{Irr}(\mu^{-1}(\xi))\}$$

with notation as in Equation (1.1). Our main problem of interest, first posed by Joseph, is to relate the basis of q_C 's to the basis of the p_I 's.

Initially it was conjectured that (up to scaling) the two bases actually coincided [J3], and Melnikov proved that is indeed the cases for $\mathfrak{sl}(n, \mathbb{C})$ [Me]. Earlier Tanisaki provided examples (outside of Type A of course) where they did not coincide [Ta]. He suggested the next possible alternative, namely that there is an upper triangular relation between the two. McGovern proved this in all classical cases [Mc3]. Our main result is a proof of this result under a technical (conjecturally superfluous) hypothesis which we subsequently verify in many cases (including all classical types).

Theorem 1.2. *Suppose \mathcal{O} is a special nilpotent orbit in \mathfrak{g}^* which satisfies Hypothesis (\star) . (This condition is described explicitly below. It holds if \mathfrak{g} is classical and is conjectured to always hold; see Section 5.) Fix $\xi \in \mathcal{O}$. Then there is a bijection from*

$$A_G(\xi) \backslash \text{Irr}(\mu^{-1}(\xi)) \longrightarrow \text{Prim}_\rho(\mathfrak{g}, \mathcal{O})$$

and a total order on these sets so that the matrix relating the p - and q -bases is upper triangular. More precisely, use the total order to enumerate $A_G(\xi) \backslash \text{Irr}(\mu^{-1}(\xi))$ as C_1, \dots, C_d and let I_j denote the image of C_j under the above bijection. Then there are positive numbers m_{ij} such that

$$q_{I_i} = \sum_{j \leq i} m_{ij} p_{C_j}.$$

The main technique of the current paper interprets the bijection, the total order, and the coefficients using characteristic cycles of Harish-Chandra modules for real groups. (The more precise Theorem 3.13 makes this clear.) Once Hypothesis (\star) is absorbed, the proofs rely only on a simple conceptual result of Chang [Ch] which, roughly speaking, says that computing leading term cycles of Harish-Chandra modules is equivalent to relating the p - and q -basis. Since there are (easy) triangularity results available for characteristic cycles one obtains the current theorem. As we remarked above, McGovern has established the conclusion of Theorem 1.2 for all classical cases (without assuming Hypothesis (\star)). However our proof of Theorem 1.2 is very different from McGovern's: the total order he provides (which is essentially combinatorial) bears no obvious relation to ours (which is essentially geometric).

To discuss Hypothesis (\star) we need some notation. Suppose $G_{\mathbb{R}}$ is a real linear reductive Lie group with Lie algebra $\mathfrak{g}_{\mathbb{R}}$. Let $K_{\mathbb{R}}$ denote its maximal compact subgroup, and write K the corresponding complexification. Let \mathfrak{g} denote complexification of $\mathfrak{g}_{\mathbb{R}}$, and let $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$ denote the complexified Cartan decomposition. K acts on \mathfrak{B} with a finite number of orbits. Given such an orbit, let $T_Q^* \mathfrak{B} \subset T^* \mathfrak{B}$

denote the conormal bundle to Q ; concretely, it is easy to see that

$$T_Q^*\mathfrak{B} = \{(\mathfrak{b}, \xi) \mid \mathfrak{b} \in Q \text{ and } \xi \in (\mathfrak{g}/\mathfrak{b} + \mathfrak{k})^*\}$$

Thus the moment map image $\mu(T_Q^*\mathfrak{B})$ is a subvariety of the nilpotent cone elements in $(\mathfrak{g}/\mathfrak{k})^*$, which we denote $\mathcal{N}_{\mathfrak{p}}^*$. (The notation reflects that the Killing form identifies the nilpotent cone in $(\mathfrak{g}/\mathfrak{k})^*$ with the nilpotent cone in \mathfrak{p} .) Note that $\mu(T_Q^*\mathfrak{B})$ is transparently invariant under the action of K , and since μ is proper and $T_Q^*\mathfrak{B}$ is irreducible, $\mu(T_Q^*\mathfrak{B})$ is irreducible as well¹. Since there are only a finite number of K orbits on $\mathcal{N}_{\mathfrak{p}}^*$, it follows that there exists such an orbit, say $\mu(Q)$, such that $\mu(Q)$ is dense in $\mu(T_Q^*\mathfrak{B})$. All such K orbits on $\mathcal{N}_{\mathfrak{p}}^*$ arise in this way, and we partition $K \backslash \mathfrak{B}$ accordingly by defining, for $\mathcal{O}^K \in K \backslash \mathcal{N}_{\mathfrak{p}}^*$,

$$\mu^{-1}(\mathcal{O}^K) = \{Q \in K \backslash \mathfrak{B} \mid \mu(Q) = \mathcal{O}^K\}.$$

Recall if \mathcal{O} is a complex nilpotent orbit in \mathfrak{g}^* , then $\mathcal{O} \cap (\mathfrak{g}/\mathfrak{k})^*$ (if nonempty) is an equidimensional union of K orbits [KR], and of course all nilpotent K orbits on $(\mathfrak{g}/\mathfrak{k})^*$ arise this way. We often implicitly assume $\mathcal{O} \cap (\mathfrak{g}/\mathfrak{k})^*$ is nonempty and write $K \backslash (\mathcal{O} \cap (\mathfrak{g}/\mathfrak{k})^*)$ for the K orbits on $\mathcal{O} \cap (\mathfrak{g}/\mathfrak{k})^*$.

We also need to recall some basic features of the geometry of the conormal bundle to an element of $\mu^{-1}(\mathcal{O}^K)$. Fix $\xi \in \mathcal{O}^K$, and define $A_K(\xi)$ to be the component group of the centralizer of ξ in K . Because K is a subgroup of G , $A_K(\xi)$ maps to $A_G(\xi)$, and so the orbits of $A_G(\xi)$ on $\text{Irr}(\mu^{-1}(\xi))$ break into potentially smaller $A_K(\xi)$ orbits. This certainly happens and presents some minor technical complications detailed in Definition 3.19 below. Now fix $Q \in \mu^{-1}(\mathcal{O}^K)$. According to Proposition 2.7 below, $T_Q^*\mathfrak{B} \cap \mu^{-1}(\xi)$ is dense in an $A_K(\xi)$ orbit, say $Z(Q)$, on $\text{Irr}(\mu^{-1}(\xi))$ and all such orbits may be written uniquely as $Z(Q)$ for $Q \in K \backslash \mathfrak{B}$.

Finally we need to recall two representation-theoretic constructions. Given an irreducible Harish-Chandra module X for $G_{\mathbb{R}}$, one may define an orbit $\text{supp}_o(X) \in K \backslash \mathfrak{B}$ by considering the support of the appropriate localization of X (Definition 3.1). By replacing X by an associated graded object and passing to its support, one defines the associated variety of X ; see Section 3. It transpires that $\text{AV}(X)$ is a finite union of closures of K orbits on $\mathcal{N}_{\mathfrak{p}}^*$. It follows easily from the definitions (see Proposition 3.10) that $\mu(T_{\text{supp}_o(X)}^*\mathfrak{B})$ is contained in $\text{AV}(X)$. Hypothesis (\star) is designed so that this inclusion is as close to an equality as possible, given that $\text{AV}(X)$ is generally reducible.

Hypothesis (\star) . *Assume for the purposes of the introduction that the $A_K(\xi)$ orbits and $A_G(\xi)$ orbits on $\text{Irr}(\mu^{-1}(\xi))$ coincide; see Section 3.3 for the general case. A complex nilpotent orbit \mathcal{O} for \mathfrak{g}^* is said to satisfy Hypothesis (\star) if there is a real group $G_{\mathbb{R}}$ with complexified Lie algebra \mathfrak{g} and an orbit \mathcal{O}_K of K on $(\mathcal{O} \cap (\mathfrak{g}/\mathfrak{k})^*)$ such that for all $Q \in \mu^{-1}(\mathcal{O}_K)$, there exists an irreducible Harish-Chandra module M_Q for $G_{\mathbb{R}}$ with trivial infinitesimal character with the property that:*

- (1) $\text{supp}_o(M_Q) = Q$; and
- (2) $\mu(T_Q^*\mathfrak{B})$ is dense in an irreducible component of $\text{AV}(M_Q)$.

We conjecture that the more general Hypothesis (\star) of Section 3.3 always holds for any special orbit \mathcal{O} . (By the above remarks, if Hypothesis (\star) holds, \mathcal{O} must indeed be special.) This is clearly a technical condition, but it doesn't seem inaccessible to general techniques. We have been unable to prove it in general however. Using less general techniques, we establish Hypothesis (\star) for all special orbits for all classical types, for G_2 , and for some cases in F_4 ; see Section 5. (As explained there, the Type D case is due to W. McGovern.)

In any event, we may now describe the bijection of Theorem 1.2. Retain the hypothesis of the theorem. In Section 3 (see Theorem 3.13), we prove that the map

$$(1.3) \quad A_G(\xi) \cdot Z(Q) \mapsto \text{Ann}(M_Q)$$

¹Actually, if K is disconnected, then $T_Q^*\mathfrak{B}$ and hence $\mu(T_Q^*\mathfrak{B})$ need not be irreducible. But this complication is harmless and we ignore it for the purposes of the introduction.

is a bijection from

$$A_G(\xi) \setminus \text{Irr}(\mu^{-1}(\xi)) \rightarrow \text{Prim}(\mathfrak{g}, \mathcal{O})_\rho,$$

as in the theorem. Since we have assumed (for the purposes of the introduction) that the $A_K(\xi)$ and $A_G(\xi)$ orbits coincide on $\text{Irr}(\mu^{-1}(\xi))$, the assignment $Q \in \mu^{-1}(\mathcal{O}_K)$ to $Z(Q) \in A_G(\xi) \setminus \text{Irr}(\mu^{-1}(\xi))$ described above is a bijection. Now $\mu^{-1}(\mathcal{O}_K)$ is a subset of K orbits on \mathfrak{B} and hence inherits the partial order arising from the closure order. Hence $A_G(\xi) \setminus \text{Irr}(\mu^{-1}(\xi))$ inherits this partial order and, according to the bijection in (1.3), so does $\text{Prim}(\mathfrak{g}, \mathcal{O})_\rho$. Fix any total order compatible with this partial order. This total order and the bijection of (1.3) are the ones that appear in Theorem 1.2.

The construction of this bijection is interesting for several reasons. Most obviously, the problem of relating the p - and q - bases seems like a problem that involves only the complex semisimple Lie algebra \mathfrak{g} , and yet the total order of the theorem is intimately related to the geometry of *real* groups. A second key observation is that for a given complex orbit \mathcal{O} , there may be several real groups $G_\mathbb{R}$, or several K -orbits \mathcal{O}_K for which Hypothesis (\star) holds. In extreme cases, for instance, by using two different components \mathcal{O}_K one can get recover two total orders that are the exact *opposite* of each other; the immediate conclusion is that the matrix in this case is actually *diagonal*. (Understanding the family of partial orders obtained in this way on $A_G(\xi) \setminus \text{Irr}(\mu^{-1}(\xi))$ might be very enlightening.) In Section 4 we interpret these phenomena as a means to transfer information about characteristic cycles of Harish-Chandra modules between two different real forms of the same complex group; see Remark 3.14. In Remark 5.47, we indicate how this transfer could be related to information about the conjectural automorphic spectrum of a real group.

The theorem indicates the importance of Hypothesis (\star) . As remarked above it holds for any special orbit \mathcal{O} for a classical Lie algebra. Section 5 gives a complete status report, but for the purposes of the introduction, we give one example. Suppose $\mathfrak{g} = \mathfrak{sp}(2n, \mathbb{C})$ and fix \mathcal{O} . Then \mathcal{O} is parametrized by a partition of $2n$ in which all odd parts occur an even number of times. Suppose also that each even part $2n_i$ occur an even number of times, say $2d_i$. This latter condition is equivalent to the existence of some p and q with $p+q=n$ such that if $G_\mathbb{R} = \text{Sp}(p, q)$, then $\mathcal{O} \cap (\mathfrak{g}/\mathfrak{k})^*$ is nonempty. In Theorem 5.2, we prove that Hypothesis (\star) holds for \mathcal{O} by choosing *any* irreducible component of $\mathcal{O} \cap (\mathfrak{g}/\mathfrak{k})^*$, and thus Theorem 1.2 applies. As we let p and q vary and the choice of irreducible component \mathcal{O}_K , one obtains roughly $2^{\sum_i d_i}$ possibilities satisfying Hypothesis (\star) . The partial orders corresponding for to each different choice are wildly different (although the bijections they define are the same). Piecing each of them together gives powerful information. See Section 5 for more details.

To conclude the introduction, it is worthwhile to detail the content of our techniques in the special case of complex groups or, by the Bernstein-Gelfand-Gelfand-Enright-Joseph equivalence of categories (e.g. [BB3]), the highest weight category. Fix a Borel subalgebra $\mathfrak{b}_1 \subset \mathfrak{g}$, write B_1 for the corresponding complex group, and let L denote an irreducible (\mathfrak{g}, B_1) -module with trivial infinitesimal character. We may write $L = L(w\rho)$ in the standard parametrization, and recall the associated variety $\text{AV}(L)$ (e.g. [J3]); see Section 4 below. Write $\overline{\mathcal{O}} = \text{AV}(\text{Ann}(L))$, and fix $\xi \in \mathcal{O}$. Then $\text{AV}(L)$ is a union of the closures of a subset of the irreducible components $\mathbb{O}_1, \dots, \mathbb{O}_n$ of $\mathcal{O} \cap (\mathfrak{g}/\mathfrak{b}_1)^*$. Spaltenstein ([Sp]) attached to each \mathbb{O}_i an orbit Z_i of $A_G(\xi)$ on $\text{Irr}(\mu^{-1}(\xi))$ (Proposition 2.3). If we apply the equivalence of categories between (\mathfrak{g}, B_1) -modules and the category of Harish-Chandra bimodules for the complex group G , and use the properties of characteristic cycles developed in Section 3, we may conclude that

$$(1.4) \quad \text{AV}(L) = \overline{\mathbb{O}}_1 \cup \dots \cup \overline{\mathbb{O}}_k \quad \text{if and only if} \quad q_{\text{Ann}(L(w^{-1})\rho)} = \sum_{i=1}^k c_i p_{Z_i} \quad \text{with each } c_i \neq 0.$$

This (given the computations of [H]) is the main result of [J3], and indeed the main motivation for his definition of the p -basis. (We sketch a simple proof of (1.4) using the techniques of this paper in Theorem 4.1 below.) Thus Theorem 1.2 implies that the geometry of characteristic cycles of real groups has implications in the highest weight category. Earlier in [T3, Theorem 1.3], using the computation of characteristic cycles of derived functor modules, we produced many new infinite

families of highest weight modules with irreducible associated varieties, and different real forms gave different families. The present ideas may be seen as a significant extension.

Acknowledgments. I thank W. McGovern for numerous helpful conversations as well as continued encouragement during the course of this project. His paper [Mc3] was the inspiration for the current paper. I thank him for allowing me to include Theorem 5.43 here. I also thank D. Milićić for helpful conversations related to [Ch].

2. BACKGROUND AND NOTATION

2.1. General Notation. Let G be a complex reductive group. Let $G_{\mathbb{R}}$ be a real form of G corresponding to a (complexified) Cartan involution θ . Let $K_{\mathbb{R}} = G_{\mathbb{R}}^{\theta}$ denote the maximal compact subgroup of $G_{\mathbb{R}}$. Write $\mathfrak{g}_{\mathbb{R}}$ for the Lie algebra of $G_{\mathbb{R}}$, \mathfrak{g} for that of G , and likewise $\mathfrak{k}_{\mathbb{R}}$ and \mathfrak{k} . Let K be the corresponding complex subgroup of G and write $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$ for the Cartan decomposition with respect to θ . Write \mathcal{N}^* for the cone of nilpotents in \mathfrak{g}^* and $\mathcal{N}_{\mathfrak{p}}^*$ for $\mathcal{N}^* \cap (\mathfrak{g}/\mathfrak{k})^*$. Let \mathfrak{B} denote the variety of Borel subalgebras in \mathfrak{g} , and now revert to the more customary notation for a basepoint, $\mathfrak{b} = \mathfrak{h} \oplus \mathfrak{n}$, so that $\mathfrak{B} = G/B$. Let $\mu : T^*\mathfrak{B} \rightarrow \mathcal{N}(\mathfrak{g}^*)$ denote the moment map for the G -action on $T^*\mathfrak{B}$. For $\xi \in \mathcal{N}^*$, write $\mu^{-1}(\xi)$ for the inverse image of ξ under μ . Let W denote the Weyl group of \mathfrak{h} in \mathfrak{g} .

Given a complex orbit \mathcal{O} , we write $K \backslash (\mathcal{O} \cap (\mathfrak{g}/\mathfrak{k})^*)$ for the set of K orbits on $\mathcal{O} \cap (\mathfrak{g}/\mathfrak{k})^*$. When this notation is used, we often implicitly assume $\mathcal{O} \cap (\mathfrak{g}/\mathfrak{k})^*$ is nonempty.

Given $\xi \in \mathcal{N}_{\mathfrak{p}}^*$, we let $A_G(\xi)$ denote the component group of $Z_G(\xi)$, the centralizer of ξ in G , and write $A_K(\xi)$ for the component group of $Z_K(\xi)$. The groups $A_G(\xi)$ are sensitive to the isogeny class of G ; but we will only study certain orbits of $A_G(\xi)$ which are insensitive to isogeny. (So, for instance, we could assume with little loss of generality that G was adjoint.) In any event, the natural inclusion

$$Z_K(\xi) \longrightarrow Z_G(\xi)$$

descends to a map

$$i_{\xi} : A_K(\xi) \longrightarrow A_G(\xi).$$

In general, i_{ξ} is neither injective nor surjective. (First examples: surjectivity fails for the nonzero nilpotent orbit for $\mathrm{Sp}(1, 1)$; injectivity fails for any orbit in $\mathrm{Sp}(4, \mathbb{R})$ with Jordan form 2^2 .) For an algebraic variety Z , we let $\mathrm{Irr}(Z)$ denote the set of its components. Note that $A_G(\xi)$ and $A_K(\xi)$ act on $\mathrm{Irr}(\mu^{-1}(\xi))$, and, in particular, the orbits will coincide exactly when i_{ξ} is surjective.

Given $Q \in K \backslash \mathfrak{B}$ and an irreducible K -equivariant local system ψ supported on Q , we write $X(Q, \psi)$ for the Harish-Chandra module for $G_{\mathbb{R}}$ with trivial infinitesimal character corresponding to ψ via localization as in [Mi, Theorem H.5.3]. If $\psi = \mathbb{1}$, the trivial local system supported on Q , then we often write $X(Q)$ instead of $X(Q, \mathbb{1})$.

For $w \in W$, we write $L(w)$ for the simple highest weight module for \mathfrak{g} with trivial infinitesimal character parameterized by w . We adopt the standard convention so that $L(e)$ is an irreducible Verma module, while $L(w_o)$ is finite-dimensional (where w_o is the long element of W).

2.2. Primitive Ideals. Let $\mathrm{Prim}(U(\mathfrak{g}))$ denote the set of primitive ideals in $U(\mathfrak{g})$. For each element $I \in \mathrm{Prim}(U(\mathfrak{g}))$, let $\mathrm{gr}I$ denote the ideal in $S(\mathfrak{g})$ obtained from the grading of I by degree in the enveloping algebra. Define $\mathrm{AV}(I)$, the associated variety of I , to be the subvariety of \mathfrak{g}^* cut out by $\mathrm{gr}I$. It is well-known that $\mathrm{AV}(I) = \overline{\mathcal{O}}$ for some nilpotent orbit $\mathcal{O} \subset \mathcal{N}^*$ [BB1].

It is easy to see that I contains a unique maximal ideal of the center of $U(\mathfrak{g})$. Write $\chi \in \mathfrak{h}^*$ for the dominant weight parameterizing this ideal via the Harish-Chandra isomorphism and write $I \in \mathrm{Prim}(U(\mathfrak{g}))_{\chi}$. For any dominant integral $\lambda \in \mathfrak{h}^*$, write $I(\chi + \lambda) \in \mathrm{Prim}(U(\mathfrak{g}))_{\chi + \lambda}$ for the primitive ideal obtained from I using the translation functor from χ to $\chi + \lambda$. Finally recall the Goldie rank polynomial $q_I \in S(\mathfrak{h})$ attached to I . This polynomial is characterized by the condition that its value at $\chi + \lambda$ is the Goldie rank of the primitive quotient $U(\mathfrak{g})/I(\chi + \lambda)$; see [J2].

Primitive ideals with trivial infinitesimal character in type B, C, or D, are parametrized by standard domino tableaux of special shape, where the notion of specialness depends on the type [G].

2.3. Fiber polynomials. Fix $\xi \in \mathcal{N}(\mathfrak{g}^*)$ and $Z \subset \text{Irr}(\mu^{-1}(\xi))$. Fix a Borel B and consider the function, ϕ_Z say, which assigns to each antidominant character χ of B , the Euler characteristic of the restriction of the Borel-Weil line bundle $G \times_B \mathbb{C}_\chi$ to Z . All cohomology groups are in fact finite, and thus ϕ_Z is an integer-valued function on the lattice of dominant weights in \mathfrak{h}^* . The following result may be extracted from [J4].

Theorem 2.1. (1) *The function ϕ_Z extends to a W -harmonic polynomial on \mathfrak{h}^* , which we denote p_Z .*
(2) *If $C, C' \in \text{Irr}(\mu^{-1}(\xi))$ are in the same $A_G(\xi)$ orbit, then $p_C = p_{C'}$.*
(3) *The relations in (2) are the only dependence relations among the polynomials $\{p_C \mid C \in \text{Irr}(\mu^{-1}(\xi))\}$. In particular, the the polynomials $\{p_Z \mid Z \in A_G(\xi) \setminus \text{Irr}(\mu^{-1}(\xi))\}$ are independent. Moreover,*

$$\text{Span}\{p_Z \mid Z \in A_G(\xi) \setminus \text{Irr}(\mu^{-1}(\xi))\}$$

is W -invariant and isomorphic as a W -representation to $\text{Sp}(G \cdot \xi)$, the representation attached to the trivial local system on $G \cdot \xi$ by the Springer correspondence.

Remark 2.2. Joseph originally defined the p_C in a different way; again see [J4].

2.4. Orbital varieties and the Springer fiber. We recall the following result of Spaltenstein. In its statement, we fix a Borel B whose nilradical has Lie algebra \mathfrak{n} , write η for the projection of G onto G/B , and π_ξ for the map $G \rightarrow G \cdot \xi$ defined by $g \mapsto g \cdot \xi$.

Proposition 2.3. *Fix $\xi \in \mathcal{O} \in G \setminus \mathcal{N}(\mathfrak{g}^*)$. Then there is a bijection between $\text{Irr}(\mathcal{O} \cap (\mathfrak{g}/\mathfrak{b})^*)$ and $A_G(\xi) \setminus \text{Irr}(\mu^{-1}(\xi))$. Given Z in the latter set, $\pi_\xi(\eta^{-1}(Z))$ meets a unique element $\mathbb{O}(Z)$ in the former set densely, and the bijection maps Z to $\mathbb{O}(Z)$.*

2.5. Conormal bundles. Given an orbit $Q \in K \setminus \mathfrak{B}$, write $T_Q^* \mathfrak{B}$ for the conormal bundle of Q in $T^* \mathfrak{B}$. As above, let $\mu : T^* \mathfrak{B} \rightarrow \mathcal{N}(\mathfrak{g}^*)$ denote the moment map for the G -action on $T^* \mathfrak{B}$. As outlined in the introduction, the moment map image

$$(2.4) \quad \mu \left(\overline{T_Q^* \mathfrak{B}} \right)$$

is always the closure of a single K orbit on $\mathcal{N}_\mathfrak{p}^*$. To simplify notation as in the introduction,

$$(2.5) \quad \text{we write } \mu(Q) \text{ for the dense orbit of } K \text{ in } \mu \left(\overline{T_Q^* \mathfrak{B}} \right).$$

For a given $\mathcal{O}_K \in K \setminus \mathcal{N}_\mathfrak{p}^*$, define

$$(2.6) \quad \mu^{-1}(\mathcal{O}_K) := \{Q \in K \setminus \mathfrak{B} \mid \mu(Q) = \mathcal{O}_K\}.$$

Fix any $\xi \in \mathcal{O}_K$, and recall the notation $A_K(\xi)$ for the component group of the centralizer of ξ in K . I learned the following result from T. Springer.

Proposition 2.7 (e.g. [T2, Proposition 2.6.1]). *Retain the above notation. Then*

$$\mu^{-1}(\xi) \cap T_Q^* \mathfrak{B}$$

is dense in a unique orbit, say $Z(Q)$, of $A_K(\xi)$ on $\text{Irr}(\mu^{-1}(\xi))$. The map

$$Q \longrightarrow Z(Q)$$

is a bijection between $\mu^{-1}(\mathcal{O}_K)$ and $A_K(\xi) \setminus \text{Irr}(\mu^{-1}(\xi))$.

2.6. Notation for fiber polynomials. Given $Z \subset \text{Irr}(\mu^{-1}(\xi))$, recall the fiber polynomial p_Z of Section 2.3. If $Q \in K \backslash \mathfrak{B}$, we write p_Q for $p_{Z(Q)}$ where $Z(Q) = \mu^{-1}(\xi) \cap T_Q^* \mathfrak{B}$. If \mathbb{O} is an irreducible component of $\mathcal{O} \cap (\mathfrak{g}/\mathfrak{b})^*$ and $Z(\mathbb{O}) \subset \text{Irr}(\mu^{-1}(\xi))$ is the subset corresponding to \mathbb{O} by Proposition 2.3, we write $p_{\mathbb{O}}$ for $p_{Z(\mathbb{O})}$.

3. CHARACTERISTIC CYCLES OF HARISH-CHANDRA MODULES

We follow the notation of Section 2.1 and fix a real linear reductive group $G_{\mathbb{R}}$ with complexified Lie algebra \mathfrak{g} and complexified Cartan decomposition $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$. Let \mathcal{D} denote the sheaf of algebraic differential operators on the flag variety \mathfrak{B} of \mathfrak{g} .

Fix a finite length (\mathfrak{g}, K) module X with trivial infinitesimal character. Since the enveloping algebra $U(\mathfrak{g})$ acts by global differential operators on \mathfrak{B} , the localization $\mathcal{X} = \mathcal{D} \otimes_{U(\mathfrak{g})} X$ makes sense; it is a $(\mathcal{D}_{\mathfrak{B}}, K)$ module.

Definition 3.1. The support of a (\mathfrak{g}, K) module X with trivial infinitesimal character is defined to be the support of the localization \mathcal{X} . This is a K invariant subvariety of \mathfrak{B} . If X is irreducible, then the support of X is the closure of a unique K orbit on X (e.g. [Mi, Lemma H.5.1]). We denote this orbit by $\text{supp}_{\circ}(X)$.

3.1. Characteristic varieties. We need a microlocalization of the support construction. Choose a good K invariant filtration \mathcal{X}^j of \mathcal{X} compatible with the degree filtration on \mathcal{D} and pass to the associated graded object $\text{gr}\mathcal{X}$. This is a $(\text{gr}\mathcal{D}, K)$ module. Since the symbol calculus identifies $\text{gr}\mathcal{D}$ with functions on $T^*\mathfrak{B}$, we may view $\text{gr}\mathcal{X}$ as an $(\mathbb{O}_{T^*\mathfrak{B}}, K)$ module where $\mathbb{O}_{T^*\mathfrak{B}}$ denotes the structure sheaf of $T^*\mathfrak{B}$. Since X was assumed to have finite length (and hence to be finitely generated), it follows that we may identify $\text{gr}\mathcal{X}$ with a K -equivariant coherent sheaf on $T^*\mathfrak{B}$. Define the characteristic variety of X , denoted $\text{CV}(X)$, to be the support of the sheaf $\text{gr}\mathcal{X}$. This is transparently a K -equivariant subvariety of $T^*\mathfrak{B}$, but since \mathcal{X} is a special kind of \mathcal{D} module (arising as the localization of X), much more is true: there exists a subset $\text{cv}(X)$ of K orbits on \mathfrak{B} such that

$$\text{CV}(X) = \bigcup_{Q \in \text{cv}(X)} \overline{T_Q^* \mathfrak{B}};$$

here $\text{cv}(X)$ is a subset of $K \backslash \mathfrak{B}$. See [Mi, Proposition H.3.6] for more details. It is also useful to keep track of the rank of the sheaf $\text{gr}\mathcal{X}$ along each irreducible component. The resulting invariant is called the characteristic cycle and is denoted

$$\mathcal{CV}(X) = \sum_{Q \in \text{cv}(X)} m_Q \overline{T_Q^* \mathfrak{B}};$$

here each m_Q is a positive integer. Both invariants are independent of the filtration initially chosen. Clearly each construction is additive on short exact sequences and descends to the appropriate Grothendieck group.

There is no effective algorithm known to compute $\text{cv}(X)$. The following result provides some very weak information; its proof is quite easy and follows from the considerations around [Mi, Proposition H.3.6], for instance.

Proposition 3.2. *Let X be the Harish-Chandra module with trivial infinitesimal character. Recall its support (Definition 3.1). Then*

$$(3.3) \quad \text{supp}_{\circ}(X) \in \text{cv}(X) \text{ and } m_{\text{supp}_{\circ}(X)} = 1$$

and

$$(3.4) \quad \text{cv}(X) \subset \{Q \in K \backslash \mathfrak{B} \mid Q \subset \overline{\text{supp}_{\circ}(X)}\}.$$

Example 3.5. There is one important case where the computation of characteristic cycles is easy. Suppose $Q \in K \backslash \mathfrak{B}$ is closed. Then there is only one K -equivariant local system supported on Q , namely the trivial one. Recall the notation $X(Q) = X(Q, \mathbb{1})$ of Section 2.1. Then (3.3) and (3.4) taken together imply that

$$(3.6) \quad \mathcal{CV}(X(Q)) = 1 \cdot \overline{[T_Q^* \mathfrak{B}]}.$$

We may pursue this argument more generally by localizing on a partial flag variety. More precisely, suppose there exists a θ -stable parabolic \mathfrak{s} of \mathfrak{g} . (The notation \mathfrak{q} would be more customary, but the letter ‘q’ already has been taken.) Write S for the corresponding subgroup of G . Fix a Borel subalgebra $\mathfrak{b} \subset \mathfrak{s}$ and write B for the corresponding subgroup of S . Write π for the projection from G/B to G/S . Suppose $Q \in K \backslash \mathfrak{B}$ is dense in the preimage of the (closed) orbit of \mathfrak{s} under K on G/Q . Then $X(Q)$ is a derived functor module of the form $A_{\mathfrak{s}}$ (in the terminology of [VZ]). It follows that

$$(3.7) \quad \mathcal{CV}(X(Q)) = 1 \cdot \overline{[T_Q^* \mathfrak{B}]},$$

which generalizes (3.6). This will be important in Example 3.21 below.

We have assumed that X has trivial infinitesimal character. We could have just as easily worked with any infinitesimal character that differed from the trivial one by a weight of a finite-dimensional representation of \mathfrak{g} . In this setting, we localize using a sheaf of twisted differential operators on the flag variety, and define $\mathcal{CV}(X)$ just as above.

Proposition 3.8. *Assume for simplicity that $G_{\mathbb{R}}$ is connected. Suppose X is an irreducible Harish-Chandra module with trivial infinitesimal character. Choose a representative ρ of the infinitesimal character, or equivalently a system of positive roots Δ^+ for a fixed Cartan \mathfrak{h} in \mathfrak{g} . Suppose ν is a weight of \mathfrak{h} that is integral and dominant for Δ^+ . Define the translation functor $\psi_{\rho}^{\rho+\nu}$ from infinitesimal character ρ to $\nu + \rho$ (e.g. [KnV, Equation 7.141]) and set $X(\nu) = \psi_{\rho}^{\rho+\nu}(X)$. Then*

$$(3.9) \quad \mathcal{CV}(X) = \mathcal{CV}(X(\nu)).$$

(If $G_{\mathbb{R}}$ is disconnected, the extra technical requirements needed to define translation functors are treated in [KnV, Theorem 2.229]. Once this is done, the conclusion of (3.9) holds.)

3.2. Associated Varieties. The construction of the characteristic variety may be imitated without localizing; see [V5]. One chooses a good K -invariant filtration X^j on X compatible with the degree filtration on the enveloping algebra. The associated graded $\text{gr}X$ is a $(\text{gr}U(\mathfrak{g}), K)$ module. Of course $\text{gr}U(\mathfrak{g}) = S(\mathfrak{g})$, and since the filtration was chosen to be K -invariant, the action of $S(\mathfrak{g})$ factors to $S(\mathfrak{g}/\mathfrak{k})$. Thus $\text{gr}X$ is a finitely generated $(S(\mathfrak{g}/\mathfrak{k}), K)$ module, i.e. a K -equivariant coherent sheaf on $(\mathfrak{g}/\mathfrak{k})^*$. The associated variety of X , denoted $\text{AV}(X)$, is defined to be the support of this sheaf. It is a K -invariant subvariety of $(\mathfrak{g}/\mathfrak{k})^*$, but again since X is a special kind of $U(\mathfrak{g})$ module (annihilated by a central ideal of finite codimension) much more is true: there exists a subset $\text{av}(X)$ of K orbits on $\mathcal{N}_{(\mathfrak{g}/\mathfrak{k})^*}$ such that

$$\text{AV}(X) = \bigcup_{\mathcal{O}_K \in \text{av}(X)} \overline{\mathcal{O}_K}.$$

In fact, $\mathcal{O} := G \cdot \mathcal{O}_K$ is well-defined independent of the choice of $\mathcal{O}_K \in \text{av}(X)$ and, moreover, \mathcal{O} is dense in $\text{AV}(\text{Ann}(X))$. According to a theorem of Barbasch-Vogan, \mathcal{O} is a special orbit. Again we may keep track of the rank of the sheaf $\text{gr}X$ along each irreducible component and define the associated cycle

$$\mathcal{AV}(X) = \sum_{\mathcal{O}_K \in \text{av}(X)} m_{\mathcal{O}_K} \overline{\mathcal{O}_K},$$

where each $m_{\mathcal{O}_K}$ is a positive integer. Again both $\text{AV}(X)$ and $\mathcal{AV}(X)$ do not depend on the choice of filtration.

There is a simple relationship between the sets $\text{cv}(X)$ and $\text{av}(X)$.

Proposition 3.10. *Let X be a finite-length (\mathfrak{g}, K) module with trivial infinitesimal character. Let μ denote the moment map $T^*\mathfrak{B} \rightarrow \mathfrak{g}^*$ and recall the notation of (2.4). Then $\mu(\text{CV}(X)) = \text{AV}(X)$; that is*

$$\text{av}(X) = \{\mu(Q) \mid Q \in \text{cv}(X)\}.$$

Consequently, in the notation of Propositions 3.2 and 3.8,

$$\text{AV}(X) = \text{AV}(X(\nu)).$$

The relationship between the multiplicities in $\text{CV}(X)$ and $\text{AV}(X)$ is more subtle however, as the following result indicates must be the case.

Proposition 3.11 (e.g. proof of [V1, Corollary 4.4]). *Retain the notation of Proposition 3.8. Fix $\mathcal{O}_K \in \text{av}(X)$. By Proposition 3.10, $\mathcal{O}_K \in \text{av}(X(\nu))$, and so we may consider the assignment*

$$\nu \mapsto m_{\mathcal{O}_K}(\nu) := m_{\mathcal{O}_K}(X(\nu)).$$

Then $m_{\mathcal{O}_K}$ extends to a harmonic polynomial on \mathfrak{h} .

Roughly speaking, for suitably compatible choices of filtrations, $\text{gr}X$ is the pushforward of $\text{gr}\mathcal{X}$ via the moment map μ . So to compute the rank of $\text{gr}X$ along an irreducible component, we have to compute the integral over the fiber of μ , weighted by the appropriate rank of $\text{gr}\mathcal{X}$. To make that precise we need a little more notation. Given $\mathcal{O}_K \in \text{av}(X)$, define (using the notation of (2.6) and (2.5))

$$\text{cv}(X; \mathcal{O}_K) := \text{cv}(X) \cap \mu^{-1}(\mathcal{O}_K) = \{Q \in \text{cv}(X) \mid \mu(Q) = \mathcal{O}_K\},$$

and write

$$\text{CV}(X; \mathcal{O}_K) = \bigcup_{Q \in \text{cv}(X; \mathcal{O}_K)} \overline{T_Q^* \mathfrak{B}},$$

and

$$\text{CV}(X; \mathcal{O}_K) = \sum_{Q \in \text{cv}(X; \mathcal{O}_K)} m_Q \overline{T_Q^* \mathfrak{B}}.$$

We call this the *leading term of the characteristic cycle of X over \mathcal{O}_K* . The pushforward argument may be quantified precisely as follows.

Proposition 3.12 ([Ch, Corollary 2.5.6]). *Retain the notation of Proposition 3.8, and fix any $\mathcal{O}_K \in \text{av}(X) = \text{av}(X(\nu))$. Then*

$$m_{\mathcal{O}_K}(\nu) = \sum_{Q \in \text{cv}(X; \mathcal{O}_K)} m_Q p_Q(\nu)$$

with notation as in Section 2.6.

An argument reproduced in Section 1.6 of [Ch] shows that the multiplicity polynomial in Proposition 3.12 is in fact proportional to the Goldie rank polynomial of the annihilator of X . More precisely, we have the following result.

Theorem 3.13. *Suppose X has integral infinitesimal character. Fix any $\mathcal{O}_K \in \text{av}(X)$. Then there is a nonzero constant c such that*

$$q_{\text{Ann}(X)} = c \sum_{Q \in \text{cv}(X; \mathcal{O}_K)} m_Q p_Q,$$

with notation as in Sections 2.2 and 2.6.

Remark 3.14. It is important to note that the theorem holds for *any* $\mathcal{O}_K \in \text{av}(X)$. Recall (Theorem 2.1) that the polynomials p_Q are nearly independent. Thus Theorem 3.13 says (up to the slight potential dependence of the p_Q 's) that once one computes $\mathcal{CV}(X; \mathcal{O}_K)$ for some $\mathcal{O}_K \in \text{av}(X)$, then one may transfer this computation to compute $\mathcal{CV}(X; \mathcal{O}'_K)$ for any other $\mathcal{O}'_K \in \text{av}(X)$.

But even more is true: since the expression of Goldie rank polynomials in terms of fiber polynomials does not depend on the real form (apart from the minor complication posed by the difference between $A_K(\xi)$ and $A_G(\xi)$ orbit on $\text{Irr}(\mu^{-1}(\xi))$), one may transfer the computation of leading term cycles between *two different real forms of the same complex group*. This should have interesting and possibly deep applications. Some are proposed in Remark 5.47.

Theorem 3.13 has additional significant philosophical import. It says that the information of the annihilator of X is encoded in the leading term of the characteristic cycle of X over any irreducible component of $\text{AV}(X)$. This gives a geometric interpretation of the annihilator of a Harish-Chandra module. (Such ideas were implicitly present in the context of Harish-Chandra bimodules in [BB3].)

We have assembled the set of tools to prove Theorem 1.2 in the case the orbits of $A_K(\xi)$ and $A_G(\xi)$ coincide. In fact, we have the following more precise result.

Theorem 3.15. *Fix a special orbit \mathcal{O} and $\mathcal{O}_K \in K \setminus (\mathcal{O} \cap (\mathfrak{g}/\mathfrak{k})^*)$. Assume the orbits of $A_K(\xi)$ and $A_G(\xi)$ on $\text{Irr}(\mu^{-1}(\xi))$ coincide. Suppose that Hypothesis (\star) (described in the introduction) holds for \mathcal{O}_K . Enumerate $\mu^{-1}(\mathcal{O}_K)$ as Q_1, \dots, Q_d and the Harish-Chandra modules in Hypothesis (\star) as M_{Q_1}, \dots, M_{Q_d} .*

- (1) *Recall the notation of Proposition 2.7; so $\mu^{-1}(\xi) \cap T_{Q_i}^* \mathfrak{B}$ is dense in $Z(Q_i)$. The map*

$$\text{Ann}(M_{Q_i}) \longmapsto Z(Q_i)$$

defines a bijection

$$(3.16) \quad \text{Prim}(\mathfrak{g}, \mathcal{O})_\rho := \{I \in \text{Prim}(\text{U}(\mathfrak{g}))_\rho \mid \text{AV}(I) = \overline{\mathcal{O}}\} \longrightarrow A_G(\xi) \setminus \text{Irr}(\mu^{-1}(\xi)).$$

In particular, the primitive ideals $\text{Ann}(M_{Q_i})$ are distinct.

- (2) *Choose any total order on $\mu^{-1}(\mathcal{O}_K) \subset K \setminus \mathfrak{B}$ compatible with the closure order on $K \setminus \mathfrak{B}$, and reorder indices so that $i < j$ implies $Q_i < Q_j$ in this total order. Set $I_i = \text{Ann}(M_{Q_i})$, let $d = \dim(\text{Sp}(\mathcal{O}))$, and define a $d \times d$ matrix (M_{ij}) by*

$$q_{I_j} = \sum_i m_{ij} p_{Q_i}.$$

Then M is upper triangular.

Proof. Define the matrix M as indicated in part (2) of the theorem. Theorem (3.13) together with Proposition 3.2 imply that the matrix is upper triangular with nonzero entries on the diagonal. (The fact that the matrix is triangular follows from (3.4); the fact that the diagonal entries are nonzero follows from (3.3).) This proves (2). It also proves that the Goldie rank polynomials of the various primitive ideals I_j are independent. Hence the various primitive ideals I_j are distinct. This is the last assertion of (1). The theory of Goldie rank polynomials mentioned in the introduction implies that cardinality of $\text{Prim}(\mathfrak{g}, \mathcal{O})_\rho$ is d , and so the various I_j 's exhaust $\text{Prim}(\mathfrak{g}, \mathcal{O})_\rho$. Meanwhile, the number of $A_G(\xi)$ orbits on $\text{Irr}(\mu^{-1}(\xi))$ is d by Theorem 2.1, and so the two sets in (3.16) have the same cardinality. Finally, the map of Part (1) is injective by Proposition 2.7. Hence it is bijective and the theorem is proved. \square

3.3. Hypothesis (\star) and Theorem 1.2 in general. Theorem 3.15 and the discussion in the introduction assumed that the $A_G(\xi)$ and $A_K(\xi)$ orbits on $\text{Irr}(\mu^{-1}(\xi))$ coincided. In this section, we treat the mild complications that present themselves if that is not the case.

Fix $Q \in \mu^{-1}(\mathcal{O}_K)$ (with notation as in (2.6)) and $\xi \in \mathcal{O}_K$. Recall from Proposition 2.7 that $T_Q^* \mathfrak{B} \cap \mu^{-1}(\xi)$ is dense in an $A_K(\xi)$ orbit $Z(Q)$ on $\text{Irr}(\mu^{-1}(\xi))$. The map

$$(3.17) \quad \Phi : \mu^{-1}(\mathcal{O}_K) \longrightarrow A_G(\xi) \backslash \text{irr}(\mu^{-1}(\xi))$$

$$(3.18) \quad Q \longrightarrow A_G(\xi) \cdot Z(Q)$$

is surjective by Proposition 2.7 and the fact that each $A_G(\xi)$ orbit is a union of $A_K(\xi)$ orbits. Recall that $\mu^{-1}(\mathcal{O})$ is a partially ordered by the closure order on $K \backslash \mathfrak{B}$. Here is the definition we need.

Definition 3.19. A subset $\mu^{-1}(\mathcal{O}_K)' \subset \mu^{-1}(\mathcal{O}_K)$ is said to be of *minimal type* if the map Φ of (3.17) restricts to a bijection

$$\begin{aligned} \mu^{-1}(\mathcal{O}_K)' &\longrightarrow A_G(\xi) \backslash \text{irr}(\mu^{-1}(\xi)) \\ Q &\longrightarrow A_G(\xi) \cdot Z(Q) \end{aligned}$$

and if, furthermore, whenever if $Q' \in \mu^{-1}(\mathcal{O}_K)'$ and $Q \in \mu^{-1}(\mathcal{O}_K)$ are such that $\Phi(Q) = \Phi(Q')$, then $Q' \subset \overline{Q}$. Hence a subset $\mu^{-1}(\mathcal{O}_K)'$ of minimal type is constructed by picking an element Q' out of each fiber of Φ with the property that Q' is minimal in the closure order restricted to the fiber. Note that the cardinality of any subset of minimal type is simply that of $A_G(\xi) \backslash \text{Irr}(\mu^{-1}(\xi))$, the dimension of $\text{Sp}(\mathcal{O})$. Note also that it is obvious that if the $A_G(\xi)$ and $A_K(\xi)$ orbits on $\text{Irr}(\mu^{-1}(\xi))$ coincide, then the only subset of $\mu^{-1}(\mathcal{O}_K)$ which is of minimal type is $\mu^{-1}(\mathcal{O}_K)$ itself.

Here is the general version of Hypothesis (\star); when the orbits of $A_K(\xi)$ and $A_G(\xi)$ coincide it restricts to the version given in the introduction.

Hypothesis (\star). A complex nilpotent orbit \mathcal{O} for \mathfrak{g}^* is said to satisfy Hypothesis (\star) if there is a real group $G_{\mathbb{R}}$ with complexified Lie algebra \mathfrak{g} , an irreducible component \mathcal{O}_K of $K \backslash (\mathcal{O} \cap (\mathfrak{g}/\mathfrak{k})^*)$, and a subset $\mu^{-1}(\mathcal{O}_K)'$ of $\mu^{-1}(\mathcal{O}_K)$ of minimal type (Definition 3.19), such that for all $Q \in \mu^{-1}(\mathcal{O}_K)'$, there exists a Harish-Chandra module M_Q for $G_{\mathbb{R}}$ with trivial infinitesimal character with the property that:

- (1) $\text{supp}_{\mathfrak{o}}(M_Q) = Q$; and
- (2) $\mu(T_Q^* \mathfrak{B})$ is dense in an irreducible component of $\text{AV}(M_Q)$.

The following is the more general version of Theorem 3.15. Its proof follows in exactly the same way.

Theorem 3.20. Fix a special orbit \mathcal{O} and $\mathcal{O}_K \in K \backslash (\mathcal{O} \cap (\mathfrak{g}/\mathfrak{k})^*)$. Suppose that Hypothesis (\star) holds for \mathcal{O}_K and the subset $\mu^{-1}(\mathcal{O}_K)'$ of minimal type. Enumerate $\mu^{-1}(\mathcal{O}_K)'$ as Q_1, \dots, Q_d and the Harish-Chandra modules in Hypothesis (\star) as M_{Q_1}, \dots, M_{Q_d} .

- (1) Then map

$$\text{Ann}(M_{Q_i}) \longmapsto \mu^{-1}(\xi) \cap T_{Q_i}^* \mathfrak{B}$$

defines a bijection

$$\text{Prim}(\mathfrak{g}, \mathcal{O})_{\rho} := \{I \in \text{Prim}(\text{U}(\mathfrak{g}))_{\rho} \mid \text{AV}(I) = \overline{\mathcal{O}}\} \longrightarrow A_G(\xi) \backslash \text{Irr}(\mu^{-1}(\xi)).$$

In particular, the primitive ideals $\text{Ann}(M_{Q_i})$ are distinct.

- (2) Choose any total order on $\mu^{-1}(\mathcal{O}_K)' \subset K \backslash \mathfrak{B}$ compatible with the closure order on $K \backslash \mathfrak{B}$, and reorder indices so that $i < j$ implies $Q_i < Q_j$ in this total order. Set $I_i = \text{Ann}(M_{Q_i})$, let $d = \dim(\text{Sp}(\mathcal{O}))$, and define a $d \times d$ matrix (M_{ij}) by

$$q_{I_j} = \sum_i m_{ij} p_{Q_i}.$$

Then M is upper triangular.

Example 3.21. Recall Example 3.5 above. Fix \mathcal{O}^K and assume that \mathcal{O}_K is Richardson in the sense of Definition 5.26 below. Write $Q \in K \backslash \mathfrak{B}$ for the orbit attached to \mathfrak{s} as in Example 3.5. Then, in the notation of Section 2.1, $X(Q) = A_{\mathfrak{s}}$, and \mathcal{O}_K is dense in $\mu(T_Q^* \mathfrak{B})$. Hence it follows $M_Q := X(Q)$ satisfies conditions (1) and (2) of Hypothesis (\star) . This is the main trick we will use in verifying Hypothesis (\star) in Section 5: we start with a derived functor module where the verification of the conditions of the hypothesis is easy, and then use that information to define other (not necessarily derived functor) modules M_{Q_i} to fulfill the hypothesis.

4. CONNECTION WITH HIGHEST WEIGHT MODULES

Fix a Borel subgroup B in G with Lie algebra $\mathfrak{b} = \mathfrak{h} \oplus \mathfrak{n}$, and write W for the Weyl group of \mathfrak{h} in \mathfrak{g} . Let ρ denote the half-sum of the positive roots of \mathfrak{h} in \mathfrak{n} . Consider the category of finite-length (\mathfrak{g}, B) modules. The irreducible objects with trivial infinitesimal character are parametrized by W . If w_{\circ} denotes the long element in W , we arrange the parametrization so that $L(w)$ is the unique irreducible quotient of the Verma module $M(w) := \text{ind}_{\mathfrak{b}}^{\mathfrak{g}}(\mathbb{C}_{ww_{\circ}\rho-\rho})$.

We may easily transcribe the constructions of Section 3 from the category of (\mathfrak{g}, K) modules to (\mathfrak{g}, B) modules: one need only replace K with B in the discussion. In this case we write

$$\text{CV}(L) = \bigcup_{w \in \text{cv}(L)} \overline{T_w^* \mathfrak{B}} \quad \text{CV}(L) = \sum_{w \in \text{cv}(L)} m_w \overline{T_w^* \mathfrak{B}}.$$

Here $\text{cv}(L)$ is a subset of B orbits on \mathfrak{B} ; by the Bruhat decomposition, we identify $B \backslash \mathfrak{B}$ with W , and write $T_w^* \mathfrak{B}$ for the conormal bundle to the orbit parametrized by w . Analogously we write

$$\text{AV}(L) = \bigcup_{\mathbb{O} \in \text{av}(L)} \overline{\mathbb{O}} \quad \text{AV}(L) = \sum_{\mathbb{O} \in \text{av}(L)} m_{\mathbb{O}} \overline{\mathbb{O}}.$$

Let $\overline{\mathbb{O}}$ denote the associated variety of $\text{Ann}(L)$ (Section 2.2). Just as $\text{av}(X)$ for an irreducible (\mathfrak{g}, K) module consisted of K invariant Lagrangians of $\mathcal{O} \cap (\mathfrak{g}/\mathfrak{k})^*$, $\text{av}(L)$ consists of B invariant Lagrangians in $\mathcal{O} \cap (\mathfrak{g}/\mathfrak{b})^*$.

The following theorem is due to Joseph [J3]. His proof is algebraic in nature. A very simple geometric proof using Theorem 3.13 is indicated below.

Theorem 4.1. *Consider the simple (\mathfrak{g}, B) module $L(w)$ with trivial infinitesimal character. Then*

$$\text{AV}(L(w)) = \overline{\mathbb{O}_1} \cup \dots \cup \overline{\mathbb{O}_k}$$

if and only if there exist nonzero constants c_i such that

$$q_{\text{Ann}(L(w^{-1}))} = \sum_i c_i p_{\mathbb{O}_i},$$

with notation as in Sections 2.2 and 2.6.

Sketch. Let $X(w)$ denote the irreducible Harish-Chandra module for the complex group G parameterized by w ; this is a $(\mathfrak{g} \oplus \mathfrak{g}, \Delta(G))$ module where $\Delta(G)$ denotes the diagonal copy of G in $G \times G$. It is an easy consequence of the geometric equivalence of categories between Harish-Chandra modules for G and highest weight modules for \mathfrak{g} (e.g. Borho-Brylinski [BB3, Corollary 4.10]) that $\text{AV}(X(w)) = \text{AV}(\text{Ann}(L(w))) =: \overline{\mathbb{O}}$. (Here we are being a little sloppy: $\text{AV}(X(w))$ is closure of a nilpotent orbit of $\Delta(G)$ on $(\mathfrak{g} \oplus \mathfrak{g}/\Delta(\mathfrak{g}))^*$ where $\Delta(\mathfrak{g})$ denotes the diagonal copy of \mathfrak{g} in $\mathfrak{g} \oplus \mathfrak{g}$, but we simply identify \mathcal{O} with a nilpotent orbit of G on $\mathcal{N}(\mathfrak{g}^*)$.) Moreover it is easy to check that

$$\text{Ann}_{\text{U}(\mathfrak{g} \oplus \mathfrak{g})}(X(w)) = I(w^{-1}) \otimes \text{U}(\mathfrak{g}) + \text{U}(\mathfrak{g}) \otimes I(w),$$

from which one concludes that the Goldie rank polynomial of $\text{Ann}(X(w))$ factors as

$$q_{\text{Ann}(X(w))} = q_{I(w^{-1})} q_{I(w)}.$$

Let $T_x^*(\mathfrak{B} \times \mathfrak{B})$ denote the conormal bundle to $Q(x)$, the $\Delta(G)$ orbit on $\mathfrak{B} \times \mathfrak{B}$ parameterized by x . (Informally $Q(x)$ consists of flags in relative position x .) Suppose $x \in W$ such that $\mu(T_x^*(\mathfrak{B} \times \mathfrak{B})) = \overline{\mathbb{O}}$,

and fix $\xi \in \mathcal{O}$. Proposition 2.7 attaches an orbit $\Delta(A_G(\xi)) \cdot (C_l, C_r) \in \Delta(A_G(\xi)) \setminus \text{Irr}(\mu^{-1}(\xi) \times \mu^{-1}(\xi))$ to $Q(x)$; here $\Delta(A_G(\xi))$ denotes the diagonal copy of $\Delta(A_G(\xi))$ in $A_G(\xi) \times A_G(\xi)$. Write $Z_l(x) = A_G(\xi) \cdot C_l \in A_G(\xi) \setminus \text{Irr}(\mu^{-1}(\xi))$ and similarly for $Z_r(x)$. From the geometric equivalence of categories, one deduces that $\mathbb{O} \in \text{av}(L(w))$ if and only if there exists $Q(x) \in \text{cv}(X(w); \mathcal{O})$ such that $Z_l(x)$ corresponds to \mathbb{O} in the bijection of Proposition 2.3; after some unraveling, this follows, for instance, from [BB3, Theorem 4.8(a)]. According to Theorem 3.13 and the factorization of $q_{\text{Ann}(X(w))}$ mentioned above, we conclude $Q(x) \in \text{cv}(X(w); \mathcal{O})$ if and only if the expression of $q_{L(w^{-1})}$ in the basis $\{p_Z \mid Z \in A_G(\xi) \setminus \text{Irr}(\mu^{-1}(\xi))\}$ contains $p_{Z_l(x)}$ with nonzero coefficient. The last two sentences then give the conclusion of the theorem. \square

Using Theorem 3.13, we can thus transfer information between the Harish-Chandra category and the highest weight category.

Corollary 4.2. *Fix a special nilpotent orbit \mathcal{O} , a real form $G_{\mathbb{R}}$ for G , and $\mathcal{O}_K \in K \setminus (\mathcal{O} \cap (\mathfrak{g}/\mathfrak{k})^*)$. Suppose*

- (1) *X is a Harish-Chandra module for $G_{\mathbb{R}}$ with trivial infinitesimal character such that $\mathcal{O}_K \in \text{av}(X)$; and*
- (2) *$L(w)$ is a simple highest weight module for \mathfrak{g} with trivial infinitesimal character such that $\text{Ann}(L(w^{-1})) = \text{Ann}(X)$.*

Fix $Q \in \mu^{-1}(\mathcal{O}_K)$ (notation as in Section 2.5), and let $\mathbb{O}(Q)$ denote the orbital variety corresponding to $A_G(\xi) \cdot (\mu^{-1}(\xi) \cap T_Q^ \mathfrak{B})$ (Propositions 2.3 and 2.7). Then*

$$Q \in \text{cv}(X; \mathcal{O}_K) \implies \mathbb{O}(Q) \in \text{av}(L).$$

If we further assume that the $A_K(\xi)$ and $A_G(\xi)$ orbits on $\text{Irr}(\mu^{-1}(\xi))$ coincide (so the assignment $Q \mapsto \mathbb{O}(Q)$ is bijective by Propositions 2.3 and 2.7), then

$$Q \in \text{cv}(X; \mathcal{O}_K) \iff \mathbb{O}(Q) \in \text{av}(L).$$

Proof. According to Theorem 3.13, $Q \in \text{cv}(X, \mathcal{O}_K)$ implies p_Q appears with nonzero coefficient in the expression of $q_{\text{Ann}(X)}$ in terms of fiber polynomials. Since we have assumed $\text{Ann}(X) = \text{Ann}(L(w^{-1}))$, Theorem 4.1 implies that the condition of the previous sentence implies $\mathbb{O}(Q) \in \text{av}(L)$. If $A_K(\xi) \setminus \text{Irr}(\mu^{-1}(\xi)) = A_G(\xi) \setminus \text{Irr}(\mu^{-1}(\xi))$, then Theorem 2.1(2) implies $p_Q \neq p_{Q'}$ for $Q \neq Q'$ in $\mu^{-1}(\mathcal{O}_K)$. This gives the final conclusion of the theorem. \square

Notice that Corollary 4.2 quantifies how difficult it is to compute leading terms cycles of a Harish-Chandra module X . Roughly speaking one must compute the annihilator and associated variety of X and then the associated variety of a single simple highest weight module.

Remark 4.3. It seems likely that the statement of Corollary 4.2 can be deduced from a general functorial relationship between Harish-Chandra modules and highest weight modules. We would like to return to this elsewhere.

5. HYPOTHESIS (\star)

The purpose of this section is to place Hypothesis (\star) (given in Section 3.3) in a broader context, as well as establish it in all classical cases.

So fix notation as in Hypothesis (\star) . Fix $G_{\mathbb{R}}$ such that $\mathcal{O} \cap (\mathfrak{g}/\mathfrak{k})^*$ is nonempty and fix \mathcal{O}_K and $Q \in \mu^{-1}(\mathcal{O}_K)$. Then it is essentially obvious that we may find a *virtual* Harish-Chandra module M_Q satisfying the conditions of Hypothesis (\star) for \mathcal{O}_K . This is easy to arrange since the characteristic cycle construction is additive: we start with *any* M_Q supported on Q and subtract off appropriate Harish-Chandra modules. The techniques of Section 3 then related an algebraic basis of multiplicity polynomials of the virtual Harish-Chandra modules M_Q to the geometric basis arising in the Springer fiber. The change of basis matrix is upper triangular because of Proposition 3.2. As explained in Section 2, if M_Q is irreducible — that is, if Hypothesis (\star) holds — then the multiplicity polynomial

is a Goldie rank polynomial, and hence we obtain Theorem 1.2. It seems very likely that the appropriate virtual representations may be chosen so that their multiplicity polynomials are in fact proportional to Goldie rank polynomials. If this is the case, Theorem 1.2 would hold *without* Hypothesis (\star) . But we have been unable to see that the virtual Harish-Chandra modules may be chosen so that their multiplicity polynomials are in fact of the form p_I . Thus we are left with proving Hypothesis (\star) .

We begin with a discussion of Type A. Given [Me], this is less interesting from the point of view of Tanisaki's conjecture since [Me] already established that the matrix in Theorem 1.2 is in fact diagonal. Nonetheless the type case A is worthwhile to understand. For instance, using the real group $U(p, q)$ one may recover the main results of [Me]. The case of $U(p, q)$ will also be important in the proof of Theorem 5.32. The following is taken from [T1, Theorem 5.6]; as explained there, for $GL(n, \mathbb{C})$ the result follows by combining work of Steinberg [St] and Joseph [J1].

Theorem 5.1. *Suppose $G_{\mathbb{R}} = GL(n, \mathbb{C}), GL(n, \mathbb{H})$ or $U(p, q)$. (In these cases all A -group orbits on irreducible components of the Springer fiber are singletons.) If X is an irreducible Harish-Chandra module for $G_{\mathbb{R}}$ with trivial infinitesimal character, then*

$$\mu(\overline{T_{\text{supp}_o(X)}^* \mathfrak{B}}) = \text{AV}(X).$$

Consequently if \mathcal{O} is any orbit such that $\mathcal{O} \cap (\mathfrak{g}/\mathfrak{k})^$ is nonempty, and $\mathcal{O}_K \in K \setminus (\mathcal{O} \cap (\mathfrak{g}/\mathfrak{k})^*)$, Hypothesis (\star) may be verified for \mathcal{O}_K by taking, for each $Q \in \mu^{-1}(\mathcal{O}_K)$, M_Q to be any (in fact, the unique) irreducible Harish-Chandra module with trivial infinitesimal character supported on \overline{Q} .*

By contrast [T1, Remark 8.9] essentially provides an orbit Q for $GL(n, \mathbb{R})$ for which there is *no* Harish-Chandra module M_Q satisfying the requirements (1) and (2) of the hypothesis. This indicates the subtle nature of matters.

For the groups $Sp(p, q)$ and $SO^*(2n)$, the situation is very similar to that of Theorem 5.1.

Theorem 5.2. *Suppose $G_{\mathbb{R}} = Sp(p, q)$ or $SO^*(2n)$. (For these groups, the orbits of $A_K(\xi)$ on $\text{Irr}(\mu^{-1}(\xi))$ are singletons, while those of $A_G(\xi)$ are typically larger.) Suppose X is an irreducible Harish-Chandra module with trivial infinitesimal character. Then*

$$(5.3) \quad \mu(\overline{T_{\text{supp}_o(X)}^* \mathfrak{B}}) = \text{AV}(X).$$

Consequently if \mathcal{O} is complex nilpotent orbit for \mathfrak{g} such that $\mathcal{O} \cap (\mathfrak{g}/\mathfrak{k})^$ is nonempty, $\mathcal{O}_K \in K \setminus (\mathcal{O} \cap (\mathfrak{g}/\mathfrak{k})^*)$, and $\mu^{-1}(\mathcal{O}_K)'$ is any subset of $\mu^{-1}(\mathcal{O}_K)$ of minimal type, then Hypothesis (\star) may be verified for \mathcal{O}_K by taking, for each $Q \in \mu^{-1}(\mathcal{O}_K)'$, M_Q to be any (in fact, the unique) irreducible Harish-Chandra module supported on \overline{Q} .*

Proof. The proof is a rather elaborate counting argument using Theorems 6 and 10 from [Mc2]. Given a Harish-Chandra module X for $G_{\mathbb{R}}$, let $\mathcal{C}(X)$ denote the cell of Harish-Chandra modules containing X (e.g. [Mc2, Section 1]). Write G for the complexification of $G_{\mathbb{R}}$. For the groups in question, all Cartan subgroups are connected. So the only K -equivariant local system supported on a fixed orbit $Q \in K \setminus \mathfrak{B}$ is the trivial one. As in notation of Section 2.1, we set $X(Q) = X(Q, \mathbb{1})$.

The first ingredient we need is the inclusion

$$(5.4) \quad \overline{G \cdot \mu(Q)} \subset \text{AV}(\text{Ann}(X(Q)));$$

here $\mu(Q)$ is defined as in (2.4). The conclusion of (5.4) follows from (3.3), Proposition 3.10, and the well-known fact (e.g. [V5, Theorem 8.4]) that

$$(5.5) \quad G \cdot \text{AV}(X) = \text{AV}(\text{Ann}(X)).$$

Now fix $Q \in K \setminus \mathfrak{B}$, set $\mathcal{O}_K = \mu(Q)$ (Notation (2.4)), and fix $\xi \in \mathcal{O}_K$. By imitating the explicit centralizer calculations given in [CM, Section 6.1], one quickly concludes that the image of $A_K(\xi)$ in $A_G(\xi)$ is trivial. Hence by Proposition 2.7,

$$(5.6) \quad \#\mu^{-1}(\mathcal{O}_K) = \#\text{Irr}(\mu^{-1}(\xi)),$$

with notation as in Section 2.5. On the other hand, set $\mathcal{C}(Q) = \mathcal{C}(X(Q))$, the cell containing $X(Q)$. In the terminology of [Mc2], $\mathcal{C}(Q)$ is Springer in the sense that the cell affords the same Weyl group representations as the representation on the top homology of the entire Springer fiber; see Theorems 6 and 10 of [Mc2]. This implies that

$$(5.7) \quad \#\mathcal{C}(Q) = \#\text{Irr}(\mu^{-1}(\xi)),$$

and so

$$(5.8) \quad \#\mathcal{C}(Q) = \#\mu^{-1}(\mathcal{O}_K).$$

Let N denote this number. Set $\mathcal{O} = G \cdot \mathcal{O}_K$. For the groups in question, it follows from the partition classification (see Remark 5.25 below) that \mathcal{O} is special. Enumerate the orbits of K on $\mathcal{O} \cap (\mathfrak{g}/\mathfrak{k})^*$ as $\mathcal{O}_K^1, \dots, \mathcal{O}_K^k$. Since

$$(5.9) \quad \mu^{-1}(\mathcal{O}) := \{Q \mid G \cdot \mu(Q) = \mathcal{O}\} = \mu^{-1}(\mathcal{O}_K^1) \cup \dots \cup \mu^{-1}(\mathcal{O}_K^k),$$

Equation (5.6) implies

$$(5.10) \quad \#\{Q \mid G \cdot \mu(Q) = \mathcal{O}\} = kN.$$

One the other hand, consider the set of Harish-Chandra modules

$$\{X \mid \text{AV}(\text{Ann}(X)) = \overline{\mathcal{O}}\}.$$

From general principles, this set is a union of cells, say $\mathcal{C}_1, \dots, \mathcal{C}_l$, each of whose corresponding representation contains the special representation $\text{Sp}(\mathcal{O})$; moreover, each cells whose corresponding representation contains $\text{Sp}(\mathcal{O})$ is among the \mathcal{C}_i . McGovern proves that each cell contains $\text{Sp}(\mathcal{O})$ exactly once ([Mc2, Theorem 1]). Meanwhile, it is not difficult to compute the multiplicity of $\text{Sp}(\mathcal{O})$ in the full coherent continuation representation on the Grothendieck group of Harish-Chandra modules with trivial infinitesimal character. This is done in Theorems 6 and 10 of [Mc2], and the multiplicity turns out to be exactly $k = \#K \backslash (\mathcal{O} \cap (\mathfrak{g}/\mathfrak{k})^*)$. Combining the last few sentences implies that $k = l$. Consequently, $\{X \mid \text{AV}(\text{Ann}(X)) = \overline{\mathcal{O}}\}$ is a union of k cells, each of size N (by (5.8)), and we conclude from (5.10) that

$$(5.11) \quad \#\{Q \mid G \cdot \mu(Q) = \mathcal{O}\} = \#\{X \mid \text{AV}(\text{Ann}(X)) = \overline{\mathcal{O}}\} = kN.$$

As we have remarked above, all Cartan subgroups for the groups in question are connected, so the map $Q \mapsto X(Q)$ is a bijection, and we may recast (5.11) as

$$(5.12) \quad \#\{Q \mid G \cdot \mu(Q) = \mathcal{O}\} = \#\{Q \mid \text{AV}(\text{Ann}(X(Q))) = \overline{\mathcal{O}}\} = kN.$$

This equality will be important below.

We now seek to establish (for the groups in question) that equality actually holds in (5.4); that is, we now show

$$(5.13) \quad \overline{G \cdot \mu(Q)} = \text{AV}(\text{Ann}(X(Q))).$$

Suppose (5.13) fails. Let Q denote an orbit for which

$$(5.14) \quad \overline{G \cdot \mu(Q)} \subsetneq \text{AV}(\text{Ann}(X(Q))).$$

Assume $\mu(Q)$ has minimal dimension among all orbits for which the failure (5.14) holds. By assumption,

$$S_1 := \{Q' \mid G \cdot \mu(Q') = G \cdot \mu(Q)\}$$

is not contained in

$$S_2 := \{Q' \mid \text{AV}(\text{Ann}(X(Q'))) = \overline{G \cdot \mu(Q)}\}$$

since Q is contained in the former set but not the latter. But we have seen (in (5.12)) that S_1 and S_2 have the same size. Thus there must exist an element $Q'' \in S_2$ such that $Q'' \notin S_1$. Since Q'' is in S_2 ,

$$\overline{G \cdot \mu(Q)} = \text{AV}(\text{Ann}(X(Q''))).$$

Combined with (5.4) we conclude

$$(5.15) \quad G \cdot \mu(Q'') \subset \text{AV}(\text{Ann}(X(Q''))) = \overline{G \cdot \mu(Q)}.$$

Since Q'' does not belong to S_1 , we conclude that

$$(5.16) \quad G \cdot \mu(Q'') \text{ is properly contained in } \text{AV}(\text{Ann}(X(Q''))) = \overline{G \cdot \mu(Q)}.$$

But now (5.15) and (5.16) contradict the assumption that $\mu(Q)$ had minimal dimension among all orbit for which the failure (5.14) holds. This contradiction thus establishes (5.13).

We now turn to proving the first assertion of the theorem. Fix \mathcal{O} so that $K \setminus (\mathcal{O} \cap (\mathfrak{g}/\mathfrak{k})^*) = \{\mathcal{O}_K^1, \dots, \mathcal{O}_K^k\}$ is nonempty. Define $\mu^{-1}(\mathcal{O})$ as in (5.9). Now (5.5) and (5.13) imply that for any Q ,

$$(5.17) \quad \mu(Q) \text{ is dense in an irreducible component of } \text{AV}(X(Q)),$$

in the notation of Section 3.2. Now choose Q_1 of minimal dimension in $\mu^{-1}(\mathcal{O})$. We claim that indeed

$$(5.18) \quad \text{AV}(X(Q_1)) = \overline{\mu(Q_1)}.$$

Suppose Q'_1 is an element of $\text{cv}(X(Q_1))$ not equal to Q_1 . Then by Proposition 3.2, $Q'_1 \subsetneq \overline{Q_1}$. Since Q_1 has minimal dimension in $\mu^{-1}(\mathcal{O})$, $Q'_1 \notin \mu^{-1}(\mathcal{O})$. Now Proposition 3.10 implies (5.18).

Let \mathcal{C}_1 denote the cell containing $X(Q_1)$. Since the associated varieties of two elements in the same cell are the same (e.g. the proof of [BB1, Lemma 4.1]), it follows that $\text{AV}(X) = \overline{\mu(Q_1)}$ for all $X \in \mathcal{C}_1$, and indeed

$$\mathcal{C}_1 \subset \{X(Q) \mid \mu(Q) = \mu(Q_1)\}.$$

But in fact (5.8) implies that this inclusion must be an equality,

$$(5.19) \quad \mathcal{C}_1 = \{X(Q) \mid \mu(Q) = \mu(Q_1)\}.$$

Write \mathcal{O}_1 be the dense orbit in $\mu(Q_1)$. Combining (5.18) and (5.19) with the fact that associated varieties are constant on cells, we obtain

$$(5.20) \quad \mathcal{C}_1 = \{X \mid \text{AV}(X) = \overline{\mathcal{O}_1}\}$$

or after unwinding the notation

$$(5.21) \quad \text{for all } X \in \mathcal{C}_1, \quad \text{AV}(X) = \overline{\mu(T_{\text{supp}_o(X)}^* \mathfrak{B})},$$

the conclusion of the theorem for the cell \mathcal{C}_1 .

Next choose Q_2 of minimal dimension in

$$\mu^{-1}(\mathcal{O}) \setminus \mu^{-1}(\mathcal{O}_1).$$

Let $X_2 = X(Q_2)$ and write \mathcal{C}_2 for the cell containing it. Arguing as in (5.17) we conclude that $\overline{\mu(Q_2)}$ is an irreducible component of $\text{AV}(X_2)$ and arguing as in (5.18), we conclude that the only other possible irreducible component of $\text{AV}(X_2)$ is $\overline{\mu(Q_1)}$. Hence

$$\mathcal{C}_2 \subset \{X(Q) \mid \mu(Q) \in \{\mu(Q_1), \mu(Q_2)\}\},$$

and (5.19) then implies that

$$\mathcal{C}_2 \subset \mathcal{C}_1 \cup \{X(Q) \mid \mu(Q) = \mu(Q_2)\}.$$

Since cells are disjoint, we conclude

$$\mathcal{C}_2 = \{X(Q) \mid \mu(Q) = \mu(Q_2)\}.$$

Thus we may argue as in (5.20) and (5.21) to conclude

$$(5.22) \quad \mathcal{C}_2 = \{X \mid \text{AV}(X) = \overline{\mathcal{O}_2}\}$$

or after unwinding the notation

$$(5.23) \quad \text{for all } X \in \mathcal{C}_2, \text{AV}(X) = \overline{\mu(T_{\text{supp}_o(X)}^* \mathfrak{B})},$$

the conclusion of the theorem for the cell \mathcal{C}_1 .

Clearly we may continue in the this way and establish the analogs of (5.21) and (5.23) for any cell. Thus the first assertion of the theorem follows. The final assertion follows easily from the former. \square

Though it isn't relevant to Hypothesis (\star) , we isolate the following interesting consequence of the above proof.

Corollary 5.24. *Let $G_{\mathbb{R}} = \mathrm{Sp}(p, q)$ or $\mathrm{SO}^*(2n)$ and retain the notation introduced at the beginning of the proof of Theorem 5.2. Fix $\mathcal{O}_K \in K \backslash \mathcal{N}_{\mathfrak{p}}^*$. Then the set of irreducible Harish-Chandra modules X with trivial infinitesimal character such that $\mathrm{AV}(X) = \overline{\mathcal{O}_K}$ is a cell of Harish-Chandra modules. Moreover, all cells arise in this way. In particular, if X is an irreducible Harish-Chandra module for $G_{\mathbb{R}}$ with integral infinitesimal character, the associated variety of X is irreducible.*

Proof. The first two assertions follow from the general analogs of (5.20) and (5.22) established by the preceding proof. The final assertion is obvious for trivial infinitesimal character, and the general integral case follows from the final assertion of Proposition 3.10. \square

Remark 5.25. We make the hypothesis that $\mathcal{O} \cap (\mathfrak{g}/\mathfrak{k})^*$ be nonempty in Theorem 5.2 more explicit. As remarked in the introduction, nilpotent orbits for $G = \mathrm{Sp}(2n, \mathbb{C})$ are parametrized by partitions of n in which odd parts occur with even multiplicities. Fix such a partition λ , and write \mathcal{O} for the corresponding nilpotent orbit. Then there exists a real form $\mathrm{Sp}(p, q)$ of G such that $\mathcal{O} \cap (\mathfrak{g}/\mathfrak{k})^*$ is nonempty if and only if every part of λ occurs with even multiplicity. (In this case \mathcal{O} is automatically special since an orbit is special if it corresponds to a partition in which the number of even parts between consecutive odd part or greater than the largest odd part is even.) Meanwhile nilpotent orbits for $\mathrm{O}(2n, \mathbb{C})$ are parametrized by partitions of $2n$ in which even parts have even multiplicity. Fix such a partition λ and let \mathcal{O} for the corresponding orbit. Fix \mathfrak{p} arising from the real form $\mathrm{SO}^*(2n)$. Then $\mathcal{O} \cap (\mathfrak{g}/\mathfrak{k})^*$ if and only if each part of λ occurs with even multiplicity. (Again this implies \mathcal{O} is special.)

In view of Remark 5.25, Theorem 5.2 falls short of verifying Hypothesis (\star) for all orbits \mathcal{O} in types C and D. We now sketch a complete treatment of types B, C, and D using the groups $\mathrm{SO}(p, q)$ and $\mathrm{Sp}(2n, \mathbb{R})$. (The Type D case is due to W. McGovern.) As alluded to in Example 3.21, we need the following definition.

Definition 5.26. Fix $G_{\mathbb{R}}$ and the notation of Section 2.1. An orbit $\mathcal{O}_K \in K \backslash (\mathcal{O} \cap (\mathfrak{g}/\mathfrak{k})^*)$ is called *Richardson* if there exists a θ -stable parabolic $\mathfrak{s} = \mathfrak{l} \oplus \mathfrak{u}$ such that \mathcal{O}_K is dense in $K \cdot \mathfrak{u}$. (If $G_{\mathbb{R}}$ is a complex group, this latter condition reduces to the usual notion of complex Richardson orbits defined, for example, in [CM, Section 7.1].) As explained in the introduction of [T3], if \mathcal{O}_K is Richardson, then $G \cdot \mathcal{O}_K$ is automatically special.

We need to make this definition explicit for applications. We begin with $G_{\mathbb{R}} = \mathrm{Sp}(2n, \mathbb{R})$. Fix \mathcal{O} parametrized by a partition λ as in Remark 5.25. Then, according to [CM, Theorem 9.3.5], the elements of $K \backslash (\mathcal{O} \cap (\mathfrak{g}/\mathfrak{k})^*)$ are parametrized by diagrams obtained by filling the boxes of λ (viewed as a Young diagram) with plus and minus signs so as to alternate across rows and so that the number of plus signs which begin rows of each odd length $2k + 1$ coincides with the number of minus signs that begin rows of length $2k + 1$, modulo the equivalence of interchanging rows of equal length. Hence $\mathcal{O} \cap (\mathfrak{g}/\mathfrak{k})^*$ is always nonempty.

Proposition 5.27 ([T3, Corollary 4.2]). *Let $G_{\mathbb{R}} = \mathrm{Sp}(2n, \mathbb{R})$. Fix a complex special orbit \mathcal{O} . Consider $\mathcal{O}^K \in K \backslash (\mathcal{O} \cap (\mathfrak{g}/\mathfrak{k})^*)$ parametrized, as described above, by a signed diagram λ_{\pm} . Fix a chunk of even rows between consecutive odd rows or greater than the largest odd row and enumerate the distinct parts of this chunk as $2k_1, \dots, 2k_r$. Then the orbit \mathcal{O}^K is Richardson (Definition 5.26) if and only if the following two conditions are satisfied for each such maximal chunk of even rows*

- (1) *Each row of length $2k_i$ begins with the same sign ε_i ; and*

(2) There is a fixed sign ε (depending on the chunk of even rows being considered) such that

$$\varepsilon_i = \varepsilon(-1)^{k_i}.$$

Note that for any special orbit \mathcal{O} , there exists an element of $K \backslash (\mathcal{O} \cap (\mathfrak{g}/\mathfrak{k})^*)$ which is Richardson.

It is perhaps useful to give a slightly less combinatorial interpretation of the explicit Richardson condition appearing in the proposition. Fix an arbitrary orbit \mathcal{O}_K parametrized by λ_{\pm} , and let $Z(\mathcal{O}_K)$ denote the centralizer in K of a point of \mathcal{O}_K . (So $Z(\mathcal{O}_K)$ is well-defined up to isomorphism.) The group $Z(\mathcal{O}_K)$ factors as a direct product indexed by the distinct parts of the underlying partition λ . Suppose d is such a part that occurs with multiplicity m . If $m = 2l + 1$ is odd, then the part d contributes a factor $O(l, l + 1)$ to $Z(\mathcal{O}_K)$. If $m = 2k$ is even, write k_+ (resp. k_-) for the number of plus (resp. minus) signs at the beginning of rows of length $2k$ in λ_{\pm} . Then the part m contributes a factor $\mathrm{Sp}(k_+, k_-)$ to the centralizer $Z(\mathcal{O}_K)$. Hence, part (1) of the Richardson condition is that the centralizer of an element of \mathcal{O}_K is, in an appropriate sense, as compact as possible; part (2) is a requirement on the signatures of the compact factors of the centralizer.

We will ultimately verify Hypothesis (\star) for orbits \mathcal{O}_K which are a special kind of Richardson orbit. Here is the definition we need.

Definition 5.28. Let $G_{\mathbb{R}} = \mathrm{Sp}(2n, \mathbb{R})$. Fix a complex special orbit \mathcal{O} . Consider $\mathcal{O}^K \in K \backslash (\mathcal{O} \cap (\mathfrak{g}/\mathfrak{k})^*)$ parametrized, as described above, by a signed diagram λ_{\pm} . The orbit is called *relevant* if it is Richardson (Definition 5.26 and Proposition 5.27) and additionally if λ has only even parts, the condition of Proposition 5.27 applies to the chunk consisting of all its even parts. (This latter condition is of course a strengthening of the Richardson one.) Note that for any special orbit \mathcal{O} , there exists an element of $K \backslash (\mathcal{O} \cap (\mathfrak{g}/\mathfrak{k})^*)$ which is relevant.

Once again, one may provide a slightly more intrinsic definition of relevant by translating it into a condition on the compact part of the centralizer of a point in \mathcal{O}_K .

We need the analogous definitions for the groups $G_{\mathbb{R}} = O(p, q)$. (The combinatorics of the full (disconnected) orthogonal group, as usual, is slightly simpler, though we could easily work with $\mathrm{SO}(p, q)$ or its identity component instead.) This time complex nilpotent orbits for \mathfrak{g} are parametrized by partitions λ in which every even part occurs an even number of times. If $p + q$ is odd (resp. even), an orbit is special if its corresponding partition has an even number of odd parts between consecutive even parts and an odd (resp. even) number of odd parts greater than the largest even part.

Fix \mathcal{O} corresponding to λ . Then, according to [CM, Theorem 9.3.4], the elements of $K \backslash (\mathcal{O} \cap (\mathfrak{g}/\mathfrak{k})^*)$ are parametrized by diagrams obtained by filling the boxes of λ (viewed as a Young diagram) with p plus and q minus signs so as to alternate across rows and so that the number of plus signs which begin rows of each even length $2k$ coincides with the number of minus signs that begin rows of length $2k$, modulo the equivalence of interchanging rows of equal length. Hence, for the split groups $O(n, n + 1)$ and $O(n, n)$, $\mathcal{O} \cap (\mathfrak{g}/\mathfrak{k})^*$ is always nonempty.

Proposition 5.29 ([T3, Corollary 7.2]). *Let $G_{\mathbb{R}} = O(p, q)$. Fix a complex special orbit \mathcal{O} . Consider $\mathcal{O}^K \in K \backslash (\mathcal{O} \cap (\mathfrak{g}/\mathfrak{k})^*)$ parametrized, as described above, by a signed diagram λ_{\pm} . Fix a maximal chunk of odd rows between consecutive even parts, and enumerate the distinct parts of this chunk as $2k_1 + 1, \dots, 2k_r + 1$. The orbit \mathcal{O}^K is Richardson (Definition 5.26) if and only if the following two conditions are satisfied for each maximal chunk of odd rows*

- (1) Each row of length $2k_i + 1$ begins with the same sign ε_i ; and
- (2) There is a fixed sign ε (depending on the chunk of odd rows being considered) such that

$$\varepsilon_i = \varepsilon(-1)^{k_i}.$$

Note that for the split groups $O(n, n + 1)$ and $O(n, n)$, given any special orbit \mathcal{O} , there exists an element of $K \backslash (\mathcal{O} \cap (\mathfrak{g}/\mathfrak{k})^*)$ which is Richardson.

Once again, the definition translates into a condition on the compact part of the centralizer. We omit the details.

Definition 5.30. Let $G_{\mathbb{R}} = \mathrm{O}(p, q)$. Fix a complex special orbit \mathcal{O} . Consider $\mathcal{O}^K \in K \backslash (\mathcal{O} \cap (\mathfrak{g}/\mathfrak{k})^*)$ parametrized, as described above, by a signed diagram λ_{\pm} . The orbit \mathcal{O}^K is called *relevant* if it is Richardson (Definition 5.26 and Proposition 5.29) and additionally if λ has only odd parts, the condition of Proposition 5.29 applies to the chunk consisting of all its odd parts. Note that for any special orbit \mathcal{O} such that $\mathcal{O} \cap (\mathfrak{g}/\mathfrak{k})^*$ is nonempty (which is always the case if $G_{\mathbb{R}}$ is split), there exists an element of $K \backslash (\mathcal{O} \cap (\mathfrak{g}/\mathfrak{k})^*)$ which is relevant.

Now fix $G'_{\mathbb{R}} = \mathrm{SO}(p, q)$ and a complex special orbit \mathcal{O}' for \mathfrak{g}' . An orbit $\mathcal{O}'_K \in K' \backslash (\mathcal{O}' \cap (\mathfrak{g}'/\mathfrak{k}')^*)$ is called relevant for $\mathrm{SO}(p, q)$ if and only if $K \cdot \mathcal{O}'_K$ is a relevant orbit for $\mathrm{O}(p, q)$.

The next result simplifies matters slightly for the groups $\mathrm{Sp}(2n, \mathbb{R})$ and $\mathrm{O}(p, q)$.

Lemma 5.31. *Let $G_{\mathbb{R}} = \mathrm{Sp}(2n, \mathbb{R})$ or $\mathrm{O}(p, q)$ and fix a nilpotent element $\xi \in (\mathfrak{g}/\mathfrak{k})^*$. Then the natural map*

$$A_K(\xi) \longrightarrow A_G(\xi)$$

is always surjective. In particular, the orbits of $A_K(\xi)$ and $A_G(\xi)$ on $\mathrm{Irr}(\mu^{-1}(\xi))$ coincide.

Proof. This may be verified by a direct computation of centralizers along the lines of [CM, Chapter 6]. We omit the details. \square

Theorem 5.32. *Let $G_{\mathbb{R}} = \mathrm{Sp}(2n, \mathbb{R})$ or $\mathrm{SO}(p, q)$ with $p + q$ odd. Fix a complex special orbit \mathcal{O} such that $\mathcal{O} \cap (\mathfrak{g}/\mathfrak{k})^*$ is nonempty. (This is always the case if $G_{\mathbb{R}}$ is split.) Fix a relevant orbit $\mathcal{O}_K \in \backslash (\mathcal{O} \cap (\mathfrak{g}/\mathfrak{k})^*)$ (Definitions 5.28 and 5.30); as remarked above, if $\mathcal{O} \cap (\mathfrak{g}/\mathfrak{k})^*$ is nonempty, then such a relevant orbit exists. For any $Q \in \mu^{-1}(\mathcal{O}_K)$, write (as in Section 2.1) $X(Q)$ for the irreducible Harish-Chandra module attached to the trivial local system on Q ; so, in particular, $\mathrm{supp}_o(X(Q)) = Q$. Then*

$$\mu(\overline{T_Q^* \mathfrak{B}}) = \mathrm{AV}(X(Q)) = \overline{\mathcal{O}_K}.$$

Thus Hypothesis (\star) may be verified by taking $M_Q = X(Q)$ for each $Q \in \mu^{-1}(\mathcal{O}_K)$.

Remark 5.33. Example 3.5.2 in [T2] shows that the relevant hypothesis on \mathcal{O}_K is necessary in general.

Remark 5.34. As the proof below shows, each module M_Q appearing in the statement of the theorem lies in the same Harish-Chandra cell as a module of the form $A_{\mathfrak{s}}$ for a θ -stable parabolic $\mathfrak{s} = \mathfrak{l} \oplus \mathfrak{u}$ where \mathcal{O}_K is dense in $K \cdot \mathfrak{u}$.

Sketch of Theorem 5.32. Lemma 5.31 simplifies notation somewhat, e.g. $\mu^{-1}(\mathcal{O}_K)$ is the only subset of minimal type appearing in Hypothesis (\star) , and $\#\mu^{-1}(\mathcal{O}_K)$ coincides with the dimension of $\mathrm{Sp}(\mathcal{O})$ (by Proposition 2.7). We will prove the theorem for $\mathrm{Sp}(2n, \mathbb{R})$. The modifications necessary for $\mathrm{SO}(p, q)$ are straightforward.

Fix \mathcal{O} and choose $\mathcal{O}_K \in K \backslash (\mathcal{O} \cap (\mathfrak{g}/\mathfrak{k})^*)$ relevant. Since \mathcal{O}_K is relevant, it is Richardson, and there exists a θ -stable $\mathfrak{s} = \mathfrak{l} \oplus \mathfrak{u}$ with \mathcal{O}_K dense in $K \cdot \mathfrak{u}$. Let $Q \in K \backslash \mathfrak{B}$ be the orbit attached to \mathfrak{s} according to Example 3.5. Example 3.21 shows that we may take $M_Q := X(Q) = A_{\mathfrak{s}}$ to satisfy conditions (1) and (2) of (\star) for the orbit Q .

We will define the other modules M_{Q_i} by applying a sequence of operators of the form $T_{\alpha\beta}$ to $M_Q = X(Q)$. These operators are introduced in [V2, Section 3] on the level of primitive ideals, and their definitions easily extend to Harish-Chandra modules using [V3, Theorem 3.10]. See also [G], [Mc1], [Mc2], [GV] for further details. Their domains and ranges may be specified using the Borho-Jantzen-Duflo τ -invariant ([V3, Definition 3.3],[G]), which is defined for Harish-Chandra modules or primitive ideals with trivial infinitesimal character. They are single-valued (resp. potentially double valued) if α and β have equal (resp. unequal) length. If $Y \in T_{\alpha\beta}(X)$ and X and Y are Harish-Chandra

modules, then X and Y belong to the same cell and hence, as mentioned in the proof of Theorem 5.2, $\text{AV}(X) = \text{AV}(Y)$. In the course of verifying Hypothesis (\star) for the modules M_{Q_i} we will also calculate the annihilators and associated varieties of these modules, using algorithms introduced in [T2, Section 3].

Set $M_Q = X(Q)$, the derived functor module fixed above. Suppose $\mathbf{T} = T_{\alpha_k, \beta_k} \circ \cdots \circ T_{\alpha_1, \beta_1}$ is a sequence of $T_{\alpha\beta}$ operators so that $\mathbf{T}(M_Q)$ is defined. Write $\mathbf{I}(Q)$ for the set of primitive ideals obtained as annihilators of elements in $\mathbf{T}(M_Q)$. Suppose that

$$(5.35) \quad \forall I \in \mathbf{I}(Q) \text{ there exists } Q_1 \in \mu^{-1}(\mathcal{O}_K) \text{ s.t. } \text{Ann}(X(Q_1)) = I \text{ and } M_{Q_1} := X(Q_1) \in \mathbf{T}(M_Q)$$

with notation as in Section 2.1. Then since $\text{supp}_o(M_{Q_1}) = Q_1$ and since (as we remarked above) \mathbf{T} preserves associated varieties (so that $\text{AV}(M_{Q_1}) = \text{AV}(M_Q) = \overline{\mathcal{O}_K}$), it follows that M_{Q_1} satisfies conditions (1) and (2) of (\star) for the orbit Q_1 . If we can make the corresponding conclusion for all well-defined sequences \mathbf{T} , then it follows that Hypothesis (\star) holds for \mathcal{O}_K : it follows from general principles that the collection of modules M_{Q_1} obtained from all such sequences \mathbf{T} correspond to a submodule of the special representation $\text{Sp}(\mathcal{O})$; since $\text{Sp}(\mathcal{O})$ is irreducible, the submodule must be all of $\text{Sp}(\mathcal{O})$; and, finally, since $\#\mu^{-1}(\mathcal{O}_K)$ coincides with the dimension of $\text{Sp}(\mathcal{O})$ (as mentioned above), it follows that conditions (1) and (2) in (\star) hold for every orbit in $\mu^{-1}(\mathcal{O}_K)$ and hence Hypothesis (\star) holds. The remaining assertions in the theorem are then obvious.

Thus our task is to establish (5.35) for every possible sequence \mathbf{T} . Clearly by induction it suffices to do this for a single operator $T_{\alpha\beta}$. More precisely, if $Q \in \mu^{-1}(\mathcal{O}_K)$ is an orbit for which $X(Q)$ satisfies condition (2) of (\star) , write $I_{\alpha\beta}(Q)$ for the (one or two element) set of primitive ideals obtained as annihilators of elements of $T_{\alpha\beta}(X(Q))$ (when defined). Then we are to show

$$(5.36) \quad \forall I \in I_{\alpha\beta}(Q), \text{ there exists } Q_1 \in \mu^{-1}(\mathcal{O}_K) \text{ s.t. } \text{Ann}(X(Q_1)) = I \text{ and } M_{Q_1} := X(Q_1) \in T_{\alpha\beta}(M_Q),$$

for each $T_{\alpha\beta}$ for which $T_{\alpha\beta}(M_Q)$ is defined. (Here Q now denotes an arbitrary element of $\mu^{-1}(\mathcal{O}_K)$, not the orbit corresponding to A_s .)

We need to recall the computation of moment map images of conormal bundles given in [T2]. For this we need the group $G'_\mathbb{R} = \text{U}(n, n)$. Adopt the usual notation for $G'_\mathbb{R}$ with the appropriate addition of primes; for instance, we will consider \mathfrak{B}' , the flag variety for $\mathfrak{g}' \simeq \mathfrak{gl}(2n, \mathbb{C})$, with moment map μ' . Embed $G_\mathbb{R}$ in $G'_\mathbb{R}$ in the obvious way. Fix a Cartan involution θ' for $G'_\mathbb{R}$ and let θ denote its restriction to $G_\mathbb{R}$, a Cartan involution for $G_\mathbb{R}$. Then K is a subgroup of K' . Fix a Cartan subalgebra \mathfrak{h}' for \mathfrak{g}' and let $\mathfrak{h} = \mathfrak{h}' \cap \mathfrak{g}$; this is a Cartan subalgebra for \mathfrak{g} .

Since K is a subgroup of K' , to each orbit $Q \in K \backslash \mathfrak{B}$, we may consider the well-defined orbit

$$Q' := K' \cdot Q \in K' \backslash \mathfrak{B}'.$$

The (injective) map $Q \mapsto Q'$ is computed explicitly in [T3, Section 2.11]. In addition, [T2, Proposition 3.3.1] implies that

$$(5.37) \quad K' \cdot \mu(Q) = \mu'(Q'),$$

with notation as in (2.5).

Suppose α is a short root for \mathfrak{h} in \mathfrak{g} . There are two roots, say α'_1 and α'_2 , for \mathfrak{h}' in \mathfrak{g}' which coincide with α when restricted to \mathfrak{h} . Meanwhile, if γ is a long root of \mathfrak{h} in \mathfrak{g} , there is a unique root, say γ' , of \mathfrak{h}' in \mathfrak{g}' which restricts to γ . In terms of the obvious notation for standard coordinates, if $\alpha = e_i - e_{i+1}$ (with $1 \leq i \leq n-1$), then $\alpha'_1 = e'_i - e'_{i+1}$ and $\alpha'_2 = e'_{2n-i} - e'_{2n+1-i}$; and if $\gamma = 2e_i$, then $\gamma' = e'_i - e'_{2n+1-i}$.

Fix a system of positive roots for \mathfrak{h}' in \mathfrak{g}' . This restricts to a system of positive roots for \mathfrak{h} in \mathfrak{g} . Suppose α is a short simple root for \mathfrak{h} in \mathfrak{g} . We want to investigate the conditions under which

$$(5.38) \quad \alpha \in \tau(X(Q)) \text{ if and only if } \alpha'_1, \alpha'_2 \in \tau(X(Q')).$$

Using the explicit calculation of $Q \mapsto Q'$, the calculations of τ -invariants for $\text{U}(p, q)$ in [G', Section 1.13], and the analogous calculations for $\text{Sp}(2n, \mathbb{R})$, it is easy to see this is always the case *except*

in one family of cases. It can happen that α is a noncompact imaginary root for $X(Q)$ (and hence not in the τ -invariant of $X(Q)$), while both α'_i are complex roots for $X(Q')$ with $\theta'(\alpha'_i) < 0$ (and so they are both in the τ invariant of $X(Q')$). As a matter of terminology (for the purposes of this proof), we say that α is an exceptional noncompact imaginary root. The first case where such a root appears is in the rank-two group $\mathrm{Sp}(4, \mathbb{R})$ with Q the unique orbit attached to the Cartan subgroups attached to \mathbb{C}^\times which does not support a derived functor module. (This orbit has complex dimension three.) Then the short simple root in exceptional noncompact. In this case $\mu(Q)$ is the unique nilpotent orbit corresponding to the partition $4 = 2 + 2$ which is not relevant. It may happen, however, in higher rank examples that we may encounter an exceptional noncompact imaginary root α for $X(Q)$ while $\mu(Q)$ is still relevant. But the following always holds: if α is a simple exceptional noncompact root for $X(Q)$ and β is a simple root adjacent to α such that $\beta \in \tau(X(Q))$, then $\mu(Q)$ is not relevant. To prove this, one needs to make a detailed analysis of the combinatorial algorithm computing $\mu(Q)$ given in [T2, Section 3]. We omit this analysis. The conclusion is that *if \mathcal{O}_K is relevant, $Q \in \mu^{-1}(\mathcal{O}_K)$, and α is short, then (5.38) holds: whenever α is not exceptional noncompact imaginary; and whenever α is exceptional noncompact imaginary so that there is a short adjacent simple root $\beta \in \tau(X(Q))$.*

Now suppose γ is a long simple root. A similar analysis as discussed in the previous paragraph allows one to conclude that

$$(5.39) \quad \gamma \in \tau(X(Q)) \text{ if and only if } \gamma' \in \tau(X(Q')),$$

with no additional restrictions on Q .

The previous two paragraphs imply that if $Q \in \mu^{-1}(\mathcal{O}_K)$ with \mathcal{O}_K relevant, then $X(Q)$ is in the domain of an operator of the form $T_{\alpha\beta}$ with α and β short if and only if $X(Q')$ is in the domain of the operators $T_{\alpha'_1, \beta'_1}$ and $T_{\alpha'_2, \beta'_2}$. Meanwhile $X(Q)$ is in the domain of $T_{\alpha\gamma}$ with γ long if and only if $X(Q')$ is in the domain of each $T_{\alpha'_i, \gamma'}$. Fix an orbit $Q \in \mu^{-1}(\mathcal{O}_K)$ so that $X(Q)$ satisfies condition (2) of (\star) . Suppose α and β are two short simple roots for \mathfrak{h} in \mathfrak{g} that together span a root system of type A_2 . Fix $Q \in K \setminus \mathfrak{B}$ and suppose $X(Q)$ is in the domain of $T_{\alpha\beta}$. Using the computations in the proof of [V3, Theorem 3.10] distilled in [G', Definition 1.14.1], and the fact that the relevant hypothesis rules out the exception remarked upon above, one deduces that there exists an orbit Q_1 so that

$$T_{\alpha\beta}(X(Q)) = X(Q_1).$$

Recall that the previous paragraph implies that $X(Q')$ is in the domain of $T_{\alpha'_1, \beta'_1}$ and $T_{\alpha'_2, \beta'_2}$, since the roots involved in the two sets of operators are orthogonal, we can consider the composition of the two applied to $X(Q')$. (The result is independent of the order the composition is taken.) Again using the computation in [V3, Theorem 3.10], and the explicit computation of the map $Q \mapsto Q'$, we deduce

$$X(Q'_1) = T_{\alpha'_1, \beta'_1} \circ T_{\alpha'_2, \beta'_2}(X(Q')).$$

It now follows from (5.37) and Theorem 5.1 (for $U(n, n)$) that indeed $Q_1 \in \mu^{-1}(\mathcal{O}_K)$. Hence we have verified (5.36) whenever \mathbf{T} is a single operator $T_{\alpha\beta}$ with α and β short (and $I_{\alpha\beta}(Q)$ necessarily consists of one element).

We now turn to verifying (5.36) for an unequal-length operator. So let α be a short simple root and let γ be a long simple roots that together span a root system of type C_2 . Fix $Q \in \mu^{-1}(\mathcal{O}_K)$ and, as usual, assume \mathcal{O}_K is relevant. It transpires that the relevant hypothesis rules out that case that α is exceptional noncompact imaginary (in the terminology introduced above). This follows by exactly the same kind of detailed analysis of the combinatorial algorithm computing $\mu(Q)$ given in [T2, Section 3] that we needed above. We omit further details.

Assume now that $X(Q)$ is in the domain of $T_{\alpha\gamma}$. Let S_1 denote $T_{\alpha\gamma}(X(Q))$. This set has one or two elements, each of which is in the domain of $T_{\alpha\gamma}$. Let S_2 be the set obtained by applying $T_{\alpha\gamma}$ to each element of S_1 ; again S_2 has one or two elements, and $X(Q)$ is always contained in S_2 . Let $S = S_1 \cup S_2$. (More abstractly, S is the smallest subset of Harish-Chandra modules with trivial infinitesimal character containing $X(Q)$ consisting of modules in the domain of $T_{\alpha\gamma}$ that is actually

closed under application of $T_{\alpha\gamma}$.) A relatively short case-by-case analysis shows that S always has three elements and, moreover, since we have assumed that α is not exceptional noncompact imaginary, each element of S is of the form $X(Q_\circ)$ for some orbit Q_\circ . Hence we may write

$$S = \{X(Q_i), X(Q_{ii}), X(Q_{iii})\}.$$

The case-by-case analysis mentioned above in fact implies that we may arrange the notation so that $T_{\alpha\gamma}(X(Q_i)) = X(Q_{ii})$, $T_{\alpha\gamma}(X(Q_{iii})) = X(Q_{ii})$, and

$$T_{\alpha\gamma}(X(Q_{ii})) = \{X(Q_i), X(Q_{iii})\}.$$

We next implement (to the extent possible) the technique used to treat the equal-length operator situation. A case-by-case analysis shows that that technique always applies to show that either

$$(5.40) \quad \mu(Q_{ii}) = \mu(Q_i) \text{ or } \mu(Q_{ii}) = \mu(Q_{iii}).$$

More precisely, we can show that there is a well-defined composition, say \mathbf{T}' , of equal-length operators for $U(n, n)$ so that either

$$\mathbf{T}'(X(Q'_{ii})) = X(Q'_i) \text{ or } \mathbf{T}'(X(Q'_{ii})) = X(Q'_{iii}).$$

Arguing as above, this implies (5.40). After further relabeling we now assume the first case holds, namely

$$(5.41) \quad \mu(Q_{ii}) = \mu(Q_i).$$

Recall that we are assuming that $X(Q)$ satisfies condition (\star) for the orbit $Q \in \mu^{-1}(\mathcal{O}_K)$ and we are trying to establish (5.36). There are three cases to consider: $Q = Q_i$, $Q = Q_{ii}$, or $Q = Q_{iii}$. In all cases, we now argue that (5.36) is ultimately reduced to proving

$$(5.42) \quad \text{if } \text{Ann}(X(Q_i)) \neq \text{Ann}(X(Q_{iii})), \text{ then } \mu(Q_i) = \mu(Q_{iii}).$$

To see this, first assume $Q = Q_i$. Then (5.41) says that (5.36) holds whenever $\text{Ann}(X(Q_i)) = \text{Ann}(X(Q_{iii}))$; hence we are reduced to (5.42). Next assume $Q = Q_{ii}$. Then (5.41) again says that (5.36) holds whenever $\text{Ann}(X(Q_i)) = \text{Ann}(X(Q_{iii}))$; hence we are again reduced to (5.42). Finally assume $Q = Q_{iii}$. If $\mu(Q_i) \neq \mu(Q_{iii})$, a case-by-case analysis with the algorithm of [T2, Section 3] shows that $\mu(Q_{iii}) \subsetneq \mu(Q_i)$. Hence the GK-dimension of $X(Q_i)$ (which is weakly greater than the dimension of $\mu(Q_i)$) is strictly greater than the dimension of $\mu(Q_{iii})$ which, by the assumption that $Q = Q_{iii}$ satisfies condition (2) of (\star) , is the GK-dimension of $X(Q_{iii})$. Hence the GK-dimension of $X(Q_i)$ is strictly greater than that of $X(Q_{iii})$. But this contradicts the fact that $X(Q_i)$ can be obtained from applying the (GK-dimension preserving) operator $T_{\alpha\gamma}$ successively to $X(Q_{iii})$. This contradiction shows that if $Q = Q_{iii}$, then indeed $\mu(Q_i) = \mu(Q_{iii})$ and (5.36) holds automatically. Hence we really are reduced to (5.42) and we may assume $Q = Q_i$ or Q_{ii} . By (5.41), part (2) of (\star) holds for Q_i if and only if it holds for Q_{ii} . So indeed we may assume it holds for both of them.

Retain the setting of (5.42) and assume that $Q = Q_{ii}$. Recall (as mentioned in Section 2.2) that primitive ideals with trivial infinitesimal character in type C are in bijection to standard domino tableaux with special shape. Moreover there is an action of $T_{\alpha\gamma}$ on domino tableaux satisfying the appropriate condition compatible with this bijection [G]. Then the domino tableaux attached by [T2, Section 3] to Q_i and Q_{iii} , which parametrize the annihilators of $X(Q_i)$ and $X(Q_{iii})$, differ by moving through a single closed cycle in the sense of [G]. The construction of [T2, Section 3] then guarantees that Q_i and Q_{iii} lie in the same fiber of μ , as desired. □

Theorem 5.43 (McGovern). *The conclusion of Theorem 5.32 holds for the orthogonal groups $SO(p, q)$ with $p + q$ even.*

Sketch. The argument leading to Theorem 5.32 applies in type D with appropriate (and sometimes rather intricate) modifications. One needs to replace the unequal-length operators T_{α_γ} appearing in the proof of Theorem 5.32 with the more complicated ones T_D and S_D defined in [Mc1] using the main results in [GV]. By making further explicit calculations, one rules out a longer list of case-by-case exceptions that arise. We omit the details. \square

Remark 5.44. For $\mathrm{Sp}(2n, \mathbb{R})$ and $\mathrm{SO}(p, q)$, the proofs of Theorems 5.32 and 5.43 provide an effective way to compute the annihilator (and associated variety) of each Harish-Chandra module of the form $X(Q)$ for $Q \in \mu^{-1}(\mathcal{O}_K)$ and \mathcal{O}_K relevant. As McGovern has pointed out, this provides the as-of-yet most complete computation of annihilators and associated varieties for these groups. Because the unitary group $\mathrm{U}(n, n)$ (all of whose Cartan subgroups are connected) is used in the proofs, one cannot conclude much about computing annihilators of modules of the form $X(Q, \psi)$ with ψ nontrivial. Such nontrivial local systems present serious complications, and the problem of computing annihilators for all representations of $\mathrm{Sp}(2n, \mathbb{R})$ and $\mathrm{SO}(p, q)$ remains open. (Garfinkle has suggested that she has made progress on this problem.)

Remark 5.45. Since Theorems 5.32 and 5.43 verify Hypothesis (\star) for all complex special orbits in types B, C, and D, it may seem that Theorem 5.2 is superfluous. But this is not the case. For the kinds of complex orbits that appear in Theorem 5.2 (see Remark 5.25), the ordering giving the triangularity of the matrix in Theorem 3.20 provided by using $\mathrm{Sp}(p, q)$ or $\mathrm{SO}^*(2n)$ can be wildly different from the ordering provided by using $\mathrm{Sp}(2n, \mathbb{R})$ or $\mathrm{SO}(p, q)$. The orderings, which originate in the closure order for the orbits of two different reductive groups on the same flag variety, in fact bear no resemblance to each other whatsoever. This is a somewhat mysterious feature of our results.

Remark 5.46. For G_2 and the complex orbits relevant for the rank one form of F_4 , the hypothesis is easy to verify. We omit the details.

Remark 5.47. The main results of [ABV], especially Chapter 26, provide an algorithm to define a set of Arthur packets whose union conjecturally exhausts the automorphic spectrum of a group $G_{\mathbb{R}}$ arising as the real points of a connected reductive algebraic group defined over \mathbb{R} . (The automorphic spectrum consists of those (unitary) representation that appear in $L^2(\Gamma \backslash G_{\mathbb{R}})$ for a congruence subgroup Γ of $G_{\mathbb{R}}$.) The algorithm depends on the computation of $\mathcal{CV}(X)$ for Harish-Chandra modules X with trivial infinitesimal character, and as we mentioned in Section 3, there is no known effective way to compute these characteristic cycles. The main results of this paper give restrictions on the leading terms of such cycles. While these results are far from definitive, they lead to highly nontrivial conclusions in examples. Using [ABV], such conclusions should have applications to computing the smallest representations in an Arthur packet. (Because the duality of [V4] is involved in the definition of such packets, the computation of leading term cycles — the “largest” piece of the characteristic cycle — corresponds to information about the smallest representations in an Arthur packet. For many interesting packets (such as unipotent packets) all representations have the same size.) Remark 3.14 is particularly intriguing since it suggests a potential relationship between the computation of Arthur packets for different real forms of the same complex group. It would be interesting to make this more explicit.

REFERENCES

- [ABV] Adams, J., D. Barbasch, and D. A. Vogan, Jr., *The Langlands Classification and Irreducible Characters for Real Reductive Groups*, Progress in Math, Birkhäuser (Boston), **104**(1992).
- [BV] D. Barbasch, D. A. Vogan, Primitive ideals and orbital integrals in complex classical groups, *Math. Ann.*, **259** (1982), no. 2, 153–199; Primitive ideals and orbital integrals in complex exceptional groups, *J. Algebra* **80** (1983), no. 2, 350–382.
- [BB1] W. Borho, J.-L. Brylinski, Differential operators on homogeneous spaces I. Irreducibility of the associated variety for annihilators of induced modules, *Invent. Math.*, **69** (1982), no. 3, 437–476.
- [BB3] W. Borho, J.-L. Brylinski, Differential operators on homogeneous spaces III. Characteristic varieties of Harish-Chandra modules and of primitive ideals, *Invent. Math.*, **80** (1985), no. 1, 1–68.

- [Ch] J.-T. Chang, Asymptotics and characteristic cycles for representations of complex groups. *Compositio Math.*, **88** (1993), no. 3, 265–283.
- [CM] D. Collingwood and W. M. McGovern, *Nilpotent orbits in semisimple Lie algebras*, Van Nostrand Reinhold (New York), 1993.
- [G] D. Garfinkle, On the classification of primitive ideals for complex classical Lie algebras I. *Compositio Math.*, **75** (1990), no. 2, 135–169; On the classification of primitive ideals for complex classical Lie algebra II, *Compositio Math.*, **81** (1992), no. 3, 307–336; On the classification of primitive ideals for complex classical Lie algebras III, *Compositio Math.*, **88** (1993), no. 2, 187–234.
- [G'] D. Garfinkle, The annihilators of irreducible Harish-Chandra modules for $SU(p, q)$ and other type A_{n-1} groups, *Amer. J. Math.*, **115** (1993), no. 2, 305–369.
- [GV] D. Garfinkle and D. A. Vogan, On the structure of Kazhdan-Lusztig cells for branched Dynkin diagrams, *J. Algebra*, **153** (1992), 91–120.
- [H] R. Hotta, On Joseph's construction of Weyl group representations, *Tohoku Math. J. (2)*, **36** (1984), no. 1, 49–74.
- [J1] A. Joseph, A characteristic variety for the primitive spectrum of a semisimple Lie algebra; short version in *Non-commutative harmonic analysis (Actes Colloq., Marseille-Luminy, 1976)*, pp. 102–118, Lecture Notes in Math., **587**, Springer(Berlin), 1977.
- [J2] A. Joseph, Goldie rank in the enveloping algebra of a semisimple Lie algebra. I, *J. Algebra*, **65** (1980), no. 2, 269–283; A. Joseph, Goldie rank in the enveloping algebra of a semisimple Lie algebra. II, *J. Algebra*, **65** (1980), no. 2, 284–306.
- [J3] A. Joseph, On the variety of a highest weight module, *J. Algebra*, **88** (1984), no. 1, 238–278.
- [J4] A. Joseph, On the characteristic polynomials of orbital varieties, *Ann. Sci. cole Norm. Sup. (4)*, **22** (1989), no. 4, 569–603.
- [KnV] A. W. Knap and D. A. Vogan, Jr., *Cohomological Induction and Unitary Representations*, Princeton Mathematical Series, number 45, Princeton University Press, 1995.
- [KR] B. Kostant, S. Rallis, Orbits and representations associated with symmetric spaces, *Amer. J. Math.*, **93** (1971), 753–809.
- [Mc1] W. M. McGovern, Left cells and domino tableaux in classical Weyl groups, *Composito Math.*, **101**(1998), no. 1, 77–98.
- [Mc2] W. M. McGovern, Cells of Harish-Chandra modules for real classical groups, *Amer. J. Math.*, **120** (1998), no. 1, 211–228.
- [Mc3] W. M. McGovern, A triangularity result for associated varieties of highest weight modules, *Comm. Algebra*, **28** (2000), no. 4, 1835–1843.
- [Me] A. Melnikov, Irreducibility of the associated varieties of simple highest weight modules in $\mathfrak{sl}(n)$, *C. R. Acad. Sci. Paris Sr. I, Math*, **316** (1993), no. 1, 53–57.
- [Mi] D. Miličić, Localization and representation theory of reductive Lie groups, electronic preprint available at www.math.utah.edu/%7Emilicic/
- [R] W. Rossmann, Invariant eigendistributions on a semisimple Lie algebra and homology classes on the conormal variety II: Representations of Weyl groups, *J. Funct. Anal.*, **96** (1991), no. 1, 155–193.
- [Sp] N. Spaltenstein, On the fixed point set of a unipotent element on the variety of Borel subgroups, *Topology*, **16** (1977), no. 2, 203–204.
- [St] R. Steinberg, An occurrence of the Robinson-Schensted correspondence, *J. Algebra*, **113** (1988), no. 2, 523–528.
- [Ta] T. Tanisaki, Characteristic varieties of highest weight modules and primitive quotients, in *Representations of Lie groups, Kyoto, Hiroshima, 1986*, 1–30, Adv. Stud. Pure Math., **14**, Academic Press(Boston), 1988.
- [T1] P. E. Trapa, Generalized Robinson-Schensted algorithms for real Lie groups, *Internat. Math. Res. Notices*, **1999**, no. 15, 803–834.
- [T2] P. E. Trapa, Symplectic and Orthogonal Robinson-Schensted algorithms, *J. Algebra*, **286** (2005), 386–404.
- [T3] P. E. Trapa, Richardson orbits for real groups, *J. Algebra*, **286** (2005), 361–385.
- [V1] D. A. Vogan, Gelfand-Kirillov dimension for Harish-Chandra modules, *Inventiones Math.*, **48** (1978), 75–98.
- [V2] D. A. Vogan, A generalized τ -invariant for the primitive spectrum of a semisimple Lie algebra, *Math. Ann.*, **242** (1979), 209–224.
- [V3] D. A. Vogan, Irreducible characters of semisimple Lie groups I, *Duke Math. J.*, **46** (1979), no. 1, 61–108.
- [V4] D. A. Vogan, Irreducible characters of semisimple Lie groups IV. Character-multiplicity duality, *Duke Math. J.*, **49** (1982), no. 4, 943–1073.
- [V5] D. A. Vogan, Associated varieties and unipotent representations, in *Harmonic analysis on reductive groups (Brunswick, ME, 1989)*, 315–388, Progress in Mathematics, **101**, Birkhäuser(Boston), 1991.
- [VZ] D. A. Vogan, G. Zuckerman, Unitary representations with nonzero cohomology, *Compositio Math.*, **53** (1984), no. 1, 51–90.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF UTAH, SALT LAKE CITY, UT 84112
E-mail address: ptrapa@math.utah.edu